sometimes slow; and prompt and full dilatation occurs with mydriatics.

The cause of this phenomenon is unknown, and it is probably a manifestation of a disorder *sui generis*. Certainly the pupillary reactions differ essentially from those with which we are most familiar—namely, the Argyll Robertson phenomenon, fixed pupils, and ophthalmoplegia interna. The Argyll Robertson pupil is usually bilateral, the myotonic pupil unilateral; the first is typically small, the latter large; the one is always syphilitic in origin, the other never. Similar considerations distinguish it from fixed pupils and ophthalmoplegia interna. The iris of the myotonic pupil is not paralysed, and if the stimulus is intense enough or sufficiently prolonged the pupil contracts through a range wider than the normal; accommodation, too, may be slow, but the range is normal. The terms "internal ophthalmoplegia" and "iridoplegia" that have been used in the titles of papers dealing with myotonic pupils are inapplicable and confusing.

Mr. Foster Moore, to whom I am greatly indebted, has seen about twenty patients with myotonic pupils, and has allowed me to examine some of them. He applied the term "non-luetic Argyll Robertson pupil" to them, but is not satisfied with it. According to the view I am expressing here it is a contradiction in terms; moreover, there are many other non-luetic spurious Argyll Robertson pupils than the one he has described. As we must use linguistic shorthand, "myotonic" pupil is perhaps the best name, but you should think of the word myotonic as being in inverted commas, because the reaction is not the same as the myotonic reaction in Thomsen's disease. Its importance, whatever you choose to call it, is that it gives rise to an unfounded suspicion of syphilis.

I might mention other non-luetic conditions in which the pupils react on convergence but not to light, but I think I have said enough to convince you that pupils conforming to this definition occur in many dissimilar disorders, and that this definition is of little value; certainly it does not define what Argyll Robertson described. I hope, too, that your own experience will prove to you that the true Argyll Robertson pupil, as I have defined it, is, as near as may be in an imperfect world, an infallible sign of syphilis of the nervous system.

COMPLETE ERADICATION OF THE THYROGLOSSAL TRACT

BY

HAMILTON BAILEY, F.R.C.S.

SURGEON, ROYAL NORTHERN HOSPITAL

(With Special Plate)

In an adult it is usually a simple matter to tell at a glance if a thyroglossal fistula is an old-standing one. Fistulas originating during infancy—they are rarely congenital—have their orifices situated low down in the neck; furthermore, the crescentic appearance of an old fistula is characteristic (Fig. 1, Special Plate).

ANATOMICAL CONSIDERATIONS

The thyroglossal tract extends from the isthmus of the thyroid gland to the foramen caecum. The ramifications of a fistula connected with this tract can often be demonstrated by injecting lipiodol prior to a lateral radiogram, and the duct can sometimes be shown to pass right into the back of the tongue (Fig. 2, Special Plate). Unless the entire tract is removed, secreting epithelium is left behind, and after a short interval the fistula reappears (Fig. A). In the case of thyroglossal cysts the story of incomplete extirpation of the tract is much the same, for simple removal of the cyst is followed by a

fistula which continues to discharge. In a review of thirty cases¹ of thyroglossal fistula I found that seventeen followed an operation for a thyroglossal cyst.

OPERATIVE PROCEDURES

Sistrunk's method of ensuring that the whole tract is removed is an excellent one, which should be looked upon as the standard operation for dealing with these otherwise troublesome cases. It is only by completely eradicating the whole tract in every case of thyroglossal

FIG. A.—Section of a thyroglossal fistula in the region of the hyoid bone, showing variations in the type of epithelium which lines the tract. (Section by Dr. P. O. Ellison.)

cyst as well as in every thyroglossal fistula that a perfect end-result may be confidently expected. The details of the operation are as follows. Through an elliptical transverse incision about the orifice, the tract is dissected up to the level of the hyoid bone. There is nothing to disturb the even tenor of the dissection up to this point. The body of the hyoid bone is then cleared on either side and divided on the left and the right so as to free a little more than a quarter of an inch of the centre of the bone, to which the tract is attached. Incidentally, even in a child, the hyoid is a tough little bone, and full-sized bone-cutting forceps are necessary to divide it cleanly. The wedge out of the hyoid is lifted up and the thyrohyoid membrane displayed. Dissection is now continued in the direction of the foramen caecum, but this time without attempting to isolate the duct (Fig. B). The tissues are cored through, allowing about a third of an inch on every side of the duct. The foramen caecum lies upwards and backwards at an angle of 45 degrees from the body of the hyoid bone (Fig. C), and it is along this line that we proceed until the cavity of the mouth is reached. This completes the dissection, and one lifts away *en bloc* the skin about the orifice of the tract; the tract as far as the hyoid bone; the centre of the body of the hyoid bone; a disk of muscle from the mylohyoid muscle; a disk of muscle from the geniohyoid muscle; a tubular portion of the very heart of the geniohyoglossi muscles, and the foramen caecum.

Within this vermiform length of tissues, a seemingly sorry prize for so much labour, lies every secreting cell of the thyroglossal tract. Nothing short of this radical extirpation will guard against recurrence.

RECONSTRUCTION

The geniohyoglossi are brought together with sutures, care being taken that the first stitch completely shuts the opening into the mouth. The tissues about the hyoid bone are approximated, but there is no necessity to attempt to bring the cut edges of the bone together. A

FIG. B.—The dissection nearing completion.

small drainage tube is inserted and the skin closed, mainly with Van Herff's clips, which in my experience is better than any other method of approximating the skin, as it gives the most satisfactory scar.

When the tract is a long one, as when the orifice is situated at some distance below the hyoid bone, the operation is considerably facilitated by making a second transverse skin incision above the first. For instance, in Case III (see below) the fistula was dissected out until the hyoid bone was reached, then a second skin incision was made at this level, and the dissected tract threaded through from the lower to the upper level, after which the dissection was begun anew.

THREE CASES OF THYROGLOSSAL FISTULA TREATED BY SISTRUNK'S OPERATION

CASE I

A married woman of 31 had a thyroglossal cyst excised at the age of 14. Shortly afterwards a fistula appeared in the region of the hyoid bone. Two years later the fistula was excised, but no sooner had the stitches been removed than excretion commenced to issue from the centre of the wound. For fifteen years the fistula continued to discharge, and she wore a little pad to absorb the moisture. Recurrent attacks of inflammation occurred at varying intervals. She presented herself as an out-patient during one of these attacks. After the inflammation had subsided, the indurated scar of the previous operations was excised and Sistrunk's operation performed. The result was highly satisfactory.

CASE II

A schoolgirl, aged 12, came to me with a history that one year previously a thyroglossal cyst had been excised from below the hyoid bone. Almost at once a fistula developed just below the hyoid bone. Repeated attacks of inflammation had occurred during the past few months. Sistrunk's operation was performed. The wound healed well. On inquiry twelve months later it was reported that the wound had healed perfectly.

CASE III

Mrs. S., aged 28. At the age of 7 a thyroglossal cyst had been operated upon and a fistula resulted. Three further operations were performed in her girlhood—apparently local excision of the area or incision during the acute attacks of inflammation. For the previous nine months, she stated, the fistula (Fig. 1, Special Plate) had leaked continuously. Sistrunk's operation was performed, a second transverse skin incision being made at the level of the hyoid bone. Intratracheal ether was given in this case, and I formed the opinion that it is the best anaesthetic for the operation.

It is interesting to note that no disability follows removal of the centre of the hyoid bone. Within forty-eight hours the patient is able to protrude the

FIG. C.—The path of the dissection after passing the hyoid.

tongue normally and to swallow without pain. The body of the hyoid bone apparently regenerates, for in one of my cases one year after Sistrunk's operation had been performed for a thyroglossal cyst a radiogram showed the hyoid bone intact.

REFERENCE

- ¹ Bailey, Hamilton: *Brachial Cysts*, 1929.

PREVENTABLE DENTAL DEFORMITIES*

BY

R. H. McKEAG, M.B., B.Ch., B.DENT.Sc.

(With Special Plate)

The dental surgeon is called upon to correct rather than to prevent malpositions of the teeth. So often does this occur that the accepted classifications of mal-occlusion, or abnormal relationship of the teeth, ignore the cause entirely, and are framed to facilitate treatment. By the time the dentist sees the patient it is usually the case that the mere removal of the cause is insufficient to bring about a cure, and indeed often the original cause has itself disappeared by then.

On investigating the subject I was amazed to find that the vast majority of orthodontic cases in my practice could have been prevented had the defect been noticed at an early stage by someone having the necessary knowledge and authority. I am therefore proposing to put before you the main causes of preventable dental deformity, and to illustrate some of the results following neglect of simple precautions. The models which I ask

* A paper read at a meeting of the Bath and Bristol Branch of the British Medical Association in conjunction with the Bristol and District Section of the British Dental Association.

HAMILTON BAILEY: COMPLETE ERADICATION OF THE THYROGLOSSAL TRACT

FIG. 1.—A long-standing thyroglossal fistula. The semilunar fold is characteristic.

FIG. 2.—Lateral radiogram after an injection of lipiodol into a thyroglossal fistula. The tract passes up to the foramen caecum. (Radiogram by Dr. H. N. Crowe, Worcester.)

R. H. McKEAG: PREVENTABLE DENTAL DEFORMITIES

FIG. 1.—Results of adenoids. Note narrow crowded arches, protruding upper incisors, corners for food to collect in which the tongue cannot clean. This patient has a receding chin and a vacant perpetual grin.

FIG. 2.—Result of thumb-sucking. Note protruding front teeth. The appearance of an otherwise good-looking girl is entirely spoiled by this deformity.