THE USE OF STRIPS OF HERNIAL SAC AS FASCIAL (LIVING) SUTURES.

BY

F. R. BROWN, M.B., F.R.C.S. Ed.,

SURGEON, ROYAL VICTORIA HOSPITAL, DUNDEE; ASSISTANT SURGEON,
ROYAL INFIRMARY, DUNDEE.

SOME years ago (1925), while operating for old-standing oblique inguinal hernia in a man of 57 with weak and lax inguinal musculature, it was my intention to reconstruct

FIG. 1.

the canal by using strips of fascia lata after the method introduced by Gallie and Le Mesurier.¹ Having dissected out and removed the sac, which was unusually long and thick, I was struck by the possibility of using strips cut from the sac itself, in the same manner as strips of fascia lata or of aponeurosis of the external oblique muscle.

In this first case the method adopted was to slit the sac up on either side to within about one inch of the fundus, so that the sac when opened out became a rough rectangular sheet measuring about eleven inches by two and a half inches (Fig. 1). From this sheet

strips a quarter of an inch wide were cut, and the operation was completed by two rows of sutures, following Gallie's technique.

This case proved completely successful, and I have now used the procedure in a large number, with some modifications in the method of cutting the strips, as illustrated in Figs. 2 and 3. Where a strip the length of the opened-out sac—that is, almost twice the length of the intact sac—would be considered too short, strips may be cut as in

FIG. 2.

FIG. 3.

Fig. 2, where the whole sac is cut into one or more long ribbons as required. Fig. 3 shows the sac cut in spiral fashion. In this method, the sac is steadied by the assistant's fingers, or may be held on a glove stretcher or two pairs of long artery forceps. Thus really long strips, suitable for continuous suturing, can be got from sacs of moderate size if desired, although it is generally in the large sac type of hernia that fascial sutures are necessary. The width of the strips can be varied according to the thickness of the sac; in a large thin-walled sac the strips may be as much as half an inch wide in order to have a sufficiently strong suture.

This procedure may be employed (1) as an alternative to the use of fascia lata or of the aponeurosis of the external oblique, or (2) in conjunction with the latter; in a recent case the sac was so small that, when cut in spiral fashion, only one rather weak strip about twelve inches long resulted, and two supplementary strips of external oblique aponeurosis were used.

So far, I have carried out this method only in cases of oblique inguinal hernia. There are obvious limitations to its use, and rarely would a sac of ample size be present in direct and femoral hernias.

REFERENCE.

¹ *Brit. Journ. of Surg.*, 1924-25, xii, 289

UNDULANT FEVER DUE TO BRUCELLA ABORTUS:

ISOLATION OF THE ORGANISM FROM THE URINE.

BY

COLONEL W. P. MACARTHUR, M.D., F.R.C.P.I.,
D.P.H., D.T.M. AND H.,

AND

MAJOR J. B. A. WIGMORE, M.B., M.R.C.S., L.R.C.P.,
D.T.M. AND H., R.A.M.C.

UP to the present seventeen cases of human infection with *Brucella abortus* have been reported in England. In only one of these¹ was the causal organism recovered, in this instance from the blood. The remaining sixteen were diagnosed by agglutination reactions only. It seems appropriate, therefore, to record another case of *Br. abortus* infection, the first in this country in which the organism has been isolated from the urine.

The patient is a lady, aged 53, resident in Manchester. Except for a brief visit to Switzerland ten years ago, she has not left the British Isles during the past twenty years. In April, 1929, she spent ten days holidaying in Ludlow, returning to Manchester on April 22nd. During the second week of May she complained of pain in one ankle, and was feverish and shivery at night, but in the mornings felt better. On the morning of May 13th she was definitely unwell, and her temperature was found to be over 102° F. The febrile attack lasted until May 29th, when the temperature, previously gradually declining, remained normal for twenty-four hours, but on the next day it commenced to rise again. Relapses, of the duration shown, followed the original attack: May 31st to July 16th; July 20th to August 2nd; August 4th to August 7th; September 14th to September 18th; November 2nd to November 27th; December 9th to December 31st. The relapses are counted from the first recorded rise of temperature to the last day that the evening temperature was above normal.

At the beginning of the first relapse the physician in charge, Dr. Florence Robinson, sent specimens of blood, urine, and faeces for examination. Agglutination tests for the typhoid group were negative, and on Professor Shaw Dunn's suggestion the serum was put up against *Br. abortus*, with a positive result. Two blood cultures (June 4th and 10th) were negative. Professor Murray saw the patient in consultation on June 10th, and the case was regarded as probably one of infection by *Br. abortus*, although the causal organism was not isolated.

During the afebrile interval following the fourth relapse the patient came to London, and on November 2nd another relapse commenced. One of us saw her in consultation on November 8th. The general condition was strongly suggestive of undulant fever. She was restless, sleepless, and despondent almost to tearfulness. The temperature fluctuated usually between 101° and 103°, and once reached 104°. Nevertheless there was little prostration, and the efforts of the nurse in charge were necessary to prevent the patient from rising at night and wandering about the house. The pulse rate was from 80 to 90. Tenderness was marked over the splenic area, but there was a difference of opinion as to whether or not the spleen was palpable. The tongue, which had been covered with a silvery fur at the beginning of the relapse, was then clean except for some persisting patches. One ankle was painful and tender. The bowels were loose, and dark offensive motions, free from any trace of mucus or blood, were passed, sometimes as many as six in the day. Obstinate constipation had been a feature of the five previous attacks. On November 10th a red papular rash appeared rather thickly over the body and limbs; rashes of various types are, of course, not uncommon in "Malta" fever. At this stage the tongue had become quite clean, and the appetite was good.

Catheter specimens of urine were taken for culture on November 11th and 12th. The first was sterile. Of the second sample 10 c.cm. were added to each of twelve tubes containing 10 c.cm. of glucose