In 12 cases out of 14 of under six months' duration there was pain in the joints as well as down the course of the nerve.

In only 9 cases out of 24 over six months' duration was there pain in the joint.

REFERENCES.

¹ Gowers: *Diseases of the Nervous System*. ² Oppenheim: *Diseases of the Nervous System*. ³ Barre and Duprez: *Revue de médecine*, 1920. ⁴ Holmes: Osler and McCrae's *System of Medicine*. ⁵ Gibson: *Lancet*, 1893.

PORTAL PYAEMIA SECONDARY TO UMBILICAL INFECTION.

BY

G. BRUCE WHITE, M.B., CH.M.SYD.,

HOUSE-PHYSICIAN, ST. GEORGE'S HOSPITAL, LONDON.

SUPPURATIVE phlebitis, as a result of spreading infection from the umbilicus, is rarely seen nowadays. The case here reported is of interest from several points of view, especially as there was no sign of inflammation of the umbilicus when the infant was admitted to hospital sixteen days after birth.

History.

On November 30th, 1922, when I was house-physician to the Royal Waterloo Hospital, a female infant aged 16 days was admitted at 9 p.m. on account of swelling of the abdomen, first noted twelve hours previously. There had been no action of the bowels for two days, even after a dose of castor oil, and an enema given that morning returned clear. Some blood and mucus had been passed per rectum during the last eighteen hours. The mother stated that the child had had several "convulsions" within a few days of birth, but had appeared healthy until three days before admission, when the skin became yellow. There had been no vomiting.

The infant was fairly well nourished, it was rather deeply jaundiced, and the marked abdominal distension rendered palpation difficult. The umbilicus was perfectly healed, the temperature was subnormal, but the pulse rate was rapid (uncountable). The abdomen was hyper-resonant from distension with gas; the liver was distinctly enlarged, but no other swelling could be detected on palpation. The rectum, the anal canal, and the lower part of the rectum felt swollen and boggy, and the examining finger was streaked with blood.

Mr. Cairns Forsyth saw the child an hour after admission, and the likely causes of the condition were discussed. The onset of the jaundice was too late for physiological jaundice of the newborn, and rather early and abrupt for that due to atresia of the bile ducts. Infection spreading to the liver by the umbilical vein was regarded as improbable, since the umbilicus was soundly healed and the temperature was subnormal. No conclusion as to the origin of the jaundice was arrived at; but since the infant was obviously suffering from intestinal obstruction it was decided that exploratory laparotomy was justifiable.

Operation.

When the child was under anaesthesia the abdomen was again palpated, but no swelling could be detected. The abdomen was opened by a small paracentral incision, the right rectus being displaced outwards. Some clear fluid escaped as soon as the peritoneum was incised and coils of intensely congested small intestine protruded into the wound. No signs of mechanical obstruction of the bowel could be found. The small intestine was everywhere distended, but the large intestine was collapsed. The appendix was normal. The liver was enlarged, of a deep maroon colour, and its lower surface was smooth. The gall bladder appeared normal and bile could be seen in its interior through its translucent wall. As the intestinal obstruction was evidently due to paresis of the bowel, secondary to a grave toxæmia probably of hepatic origin, the abdomen was closed.

Necropsy.

The child died twelve hours later, and at the autopsy the same general condition of the abdomen was observed, but on cutting through the ligamentum teres to remove the liver pus poured from the umbilical vein, which was widely distended (to three-eighths of an inch). On exposing the liver its upper surface showed a number of whitish areas which on section corresponded to small abscesses within the liver substance. The liver was enlarged and between the infected areas was intensely engorged with blood. There were a number of small haemorrhagic spots scattered throughout the lungs, but no evidence of abscess formation in any other viscera or in the brain could be detected. The lower part of the rectum and the anal canal were much congested and swollen and doubtless accounted for the passage of blood and mucus.

It is highly probable that infection took place in a clot in the umbilical vein and spread to the liver. That the navel should have healed soundly when the intra-abdominal part of the umbilical vein was full of pus is somewhat remarkable. The symptoms from which the child suffered are easily explained by the condition of the liver found at autopsy.

I am greatly indebted to Mr. Cairns Forsyth for permission to report the case.

EXCISION OF THE FIBULA IN AMPUTATIONS BELOW THE KNEE-JOINT.

BY

CHARLES NOON, F.R.C.S.,

ASSISTANT SURGEON, NORFOLK AND NORWICH HOSPITAL.

SERVICEABLE stumps after amputations of the lower extremity must possess good movement, be painless, and able to sustain the direct pressure of a part of the body weight upon the artificial limb. Without these attributes it is practically impossible to fit an absolutely satisfactory artificial limb. Much time and money are often wasted in making and attempting to fit artificial limb after artificial limb to unsatisfactory stumps, whereas correct operative treatment before fitting the artificial limb might have saved much time and trouble both to the artificial-limb maker and the patient.

During the last five or six years a great deal has been learned about amputations, amputation stumps, and the fitting of artificial limbs. Some amputations which were before the war considered "first class" have been of late unable to stand the tests to which they have been subjected. There are still probably many amputation cases in the country which require operative treatment before they can be fitted with really satisfactory artificial limbs. Many below-knee amputation stumps which are at present painful and unsound can be transformed into serviceable stumps by excision of the remaining part of the fibula and removal of that part of the external popliteal nerve which lies below the level of the knee-joint.

In a considerable number of cases of amputation below the knee-joint the presence of the fibula appears to cause considerable difficulty in fitting and wearing a satisfactory artificial limb. This difficulty is more especially noticeable in stumps where atrophy of the muscles covering the bones has been excessive. It is well known that structures that do not exert their normal function undergo atrophy, particularly muscles and bones, so that it is easy to explain the excessive atrophy often seen in amputation stumps. It is in these cases especially, where atrophy has been excessive and in which pressure on the prominent head of the fibula causes pain, that removal of the bone improves the condition and produces serviceable stumps.

Although the removal of the fibula has been practised by a few surgeons it does not appear to have received the consideration due to it, taking into account the benefit which is likely to be derived from its practice. The textbooks of surgery scarcely make any mention of the procedure. Muirhead Little in his recent book on *Amputations and Amputation Stumps* (Chapter II, p. 39) refers to the procedure as follows:

"When the amputation is below the knee excision of all that remains of the fibula will add to the amount of skin available, and sometimes enable the surgeon to cover the tibia satisfactorily without shortening it."

In *Orthopaedic Surgery of Injuries*, edited by Sir Robert Jones (vol. 1, p. 478), there is the following reference to excision of the fibula in amputation below the knee:

"In these short stumps there is a tendency for the fibula to tilt away from the tibia and project. In these cases it is a good plan to remove the fibula. In this way a piece of bone is removed which stands pressure badly, and at the same time the bulk of the stump is diminished, so that a flap which would otherwise be too small may suffice."

A man with a serviceable stump, fitted with a satisfactory artificial limb below the knee, is often able to earn his living at a laborious occupation which may necessitate him standing for many hours a day and walking long distances in the performance of his duties. The following cases, taken from a large series of amputations which have come under my care during the last five or six years, will illustrate how excision of the fibula has been the means of obtaining serviceable and satisfactory stumps.

CASE I.—*Gunshot Wound of Right Leg: Amputation below the Knee: Re-amputation below Knee with Excision of the Fibula. Result, Serviceable Stump.*

J. A., a professional golfer, aged 28, was wounded in the right leg on March 16th, 1916, while serving in France; the leg was amputated, and he was transferred to England on April 6th.

Examination (May 1st, 1916).—General condition good. Local condition: The right leg has been amputated below the knee by the guillotine method. The extremity of the stump is covered by a healthy granulating area.