as part of the arched lever of the foot; the two muscles attached to its base—the peroneus longus and tibialis anticus—have become postural in function, assisting in maintaining the arch of the foot. The two terminal phalanges, in place of being the movable parts of a grasping digit, have become the chief part of the fulcrum or base from which the human foot steps off; with each step the weight and force of the body falls on this joint of the great toe. Hence, with the evolution of the human foot the muscles which act on the phalanges, particularly the short muscles acting on the basal phalanx, had to take on a new rôle—they had to adjust and maintain the position of the hallucial phalanges at every phase of the “step off.” Hallux valgus results from a breakdown of the neuro-muscular machinery which guards the condylar joint of the great toe. No doubt ill fitting boots contribute to this breakdown, but when it is remembered that the action of muscles which cross a joint are largely regulated by stimuli which arise within a joint, we see good reason for suspecting that the initial factor in the cause of hallux valgus arises within the joint itself. We do know that people of certain constitutions are particularly prone to disorders in the ball joint of the great toe.

The Normal Position of the Great Toe.

One last point as regards the “normal” position which the human great toe should hold to the rest of the foot. From a study of the footprints of native and unshod races, and also of English infants and children, I realize that a healthy and unblemished great toe may vary in position between two extremes. If on a footprint a line be drawn along the inner border so as to touch the impresses of the margin of the heel and of the ball of the great toe, what may be called an “inner directional” line is obtained. If a similar line be drawn along the outer border of the imprint—from heel to the outer margin of the foot-pad—an “outer directional line” is obtained. In healthy normal children and natives the great toe may be parallel to either outer or inner line, or may lie at any angle between these lines. The inner direction is the more simian, but which is the more liable to become the site of hallux valgus I have no evidence.

A CASE OF HYPERPIESIA.

REPORTED BY

D. C. L. VEY, M.C., B.A., M.B. CANTAB., M.R.C.S., L.R.C.P.,
HOUSE-PHYSICIAN, UNIVERSITY COLLEGE HOSPITAL.

With Commentary by

SIR CLIFFORD ALLBUTT.

FREQUENT as this malady is amongst the well-to-do classes of life, it is—so far as my experience goes (C. A.)—scarce in hospital practice. Moreover, by the duration of the malady, and its consistency for some long period with good health, the story is often broken or incomplete. When the patient has to seek advice, he may pass from one physician to another; in private practice, for various reasons, necropsies are rarely obtained, and when obtained may be very perfunctory. Consequently, until Dr. Batty Shaw made, in his recent book, a solid and valuable contribution to its pathology, our stock of fully observed instances of hyperpiesia was very scanty.

A patient, a man aged about 60, died during last Lent term in the Addenbrooke's Hospital with arterio-sclerosis, a very large heart, and a systolic pressure of 210 mm. But little was known of his history, and before death there had not been much time for an adequate series of biochemical and other observations. The necropsy was made by Dr. J. F. Gaskell, who reported that the kidneys (of which I made a naked eye inspection) were quite normal. Death was by cardiac defeat and its effects. The case was clearly proved to be one of hyperpiesia.

In this default of instances it seems well therefore to add to our store of evidence by publishing the following notes by Dr. Vey of a case of this kind at University College Hospital. Dr. Vey, then house-physician, watched the patient very closely and made ample clinical notes during the last fifteen months of his life, including the laboratory reports and the *post-mortem* findings. Dr. Vey described the case in his thesis for M.B. at Cambridge; and, with his permission, I have made (I hope not too

drastic) an abridgement of his notes. I think that I have not omitted any relevant abnormal feature or reaction.

Report of Case by Dr. D. C. L. VEY.

A male patient, aged 61, a clerk, was admitted into University College Hospital, under Sir John Rose Bradford (and later under Dr. Sidney Martin), five times in fifteen months. The first occasion was on September 3rd, 1921. He had complained of dyspnoea on ascents for ten months, and of swelling of feet and abdomen for nine months. Previous personal history good; no bad habits.

On admission, respiratory rate 32, much general oedema; liver full; heart large and heaving; pulse regular, large in volume, rate 50 to 72. Systolic blood pressure 200. Urine very scanty; specific gravity high; albumin abundant; a few epithelial cells and leucocytes; no casts. Under treatment, which included digitalis, he improved greatly.

Second admission, December 28th, 1921; symptoms as before, but drowsiness also. Bled to 24 oz. Heart still larger, beat in seventh left space in mid-axillary line. Second aortic sound much accentuated. Blood urea, on two occasions, 20 and 37. Blood pressure (systolic) 210. Improvement as before. Blood pressure on discharge 180.

On the third and fourth admissions similar reports and results. Fifth admission, November 8th, 1922: oedema and dyspnoea worse than ever; but pulse regular, full and hard, rate 80. Radials thick; blood pressure 190. Heart as before. No murmurs in any area. Urine in first twenty-four hours 34 oz., and about the same for next seven days; specific gravity 1018; some albumin. Headache and drowsiness. Respiration at times tidal. No twitchings. Multiple incisions of legs answered well, relieving symptoms; and blood pressure fell a little to 180–200. Urine now generally reached to 70 oz., and on each of two days to 120. Blood urea on admission was 48 mg. per 100 c.cm.; a fortnight later it had fallen to 24. During a period of three weeks the daily percentage and the totals of urea in the urine ranged between 1.5 and 2.35; the total amount averaged about 40 grams. Chlorides averaged 3 grams daily. Two urea concentration tests were made on November 15th and December 5th. On the former date the concentration was 1.6 per cent. at the end of the first hour (vol. 160 c.cm.), and 1.7 per cent. (vol. 140 c.cm.) at the end of the second hour. On the second date 2.05 per cent. (vol. 224 c.cm.) was noted at the end of the first hour, and 2 per cent. at the end of the second hour.* There never had been any pyrexia, and the patient, as before, was doing well, when he had a sudden rigor; he was distressed and speechless; the temperature rose to 104.8°, he became unconscious, and died. The cause of this event was not apparent at the necropsy. A few days before death recent small haemorrhages had been reported in the right fundus oculi, and a single larger one in the left.

The diagnosis during life was very difficult. “Chronic parenchymatous nephritis” was suspected, though the extreme cardiovascular symptoms seemed hardly to confirm the suspicion. Dr. Goodhart in the *post-mortem* room, before the necropsy, expressed the opinion that “the kidneys would be found normal, or nearly so.”

The relevant features of the necropsy were briefly as follows: Heart, 39 oz.; left ventricle much hypertrophied and dilated, and right ventricle also. Aorta, traces of old syphilis; many patches of atheroma, calcareous and ulcerative. Pulmonary artery and branches dilated, stiff on section, and showed atheromatous patches. Spleen large and fibrotic, with internal thickening of the arterioles, and fat droplets. Liver nutmeg, and some arteriolar thickening. Kidneys each 9 oz.; except that the renal artery was stiff, no abnormal appearance to the eye. Brain normal to the eye. Microscopically the myocardium revealed well marked interstitial fibrosis; no fatty degeneration.

From the kidneys frozen sections were taken and stained with haematoxylin and Scharlach R.; and paraffin sections were stained with haematoxylin and eosin. The larger arteries showed a moderate degree of internal thickening; this was more pronounced in the smaller, especially the interlobular and afferent; but not to obliteration. Fat droplets in the smaller vessels. Glomeruli “surprisingly little affected; a very few showed some atrophic change with fat droplets, but nearly all were remarkably healthy.” “The tubules were practically free from change.” Interstitial fibrosis was minimal.

Such in epitome is Dr. Vey's record of a case of hyperpiesia, in its later phases thoroughly investigated. That the patient, as was the case with the one mentioned in Addenbrooke's Hospital, died with more or less of uraemic symptoms is true; but it is now fully recognized that such symptoms may, and often do, occur in the “cardiac kidney,” but in case of the patient's amendment pass off entirely. That the syndrome is essentially incidental in these cases and, in periods of amendment, transitory, has been very fully illustrated by certain French physicians; but as I am writing away from home I cannot give references. Dr. Vey reported drowsiness on the second, third, and fourth admissions; but during the temporary recoveries the uraemia passed off.

Little by little, then, our knowledge of hyperpiesia in its later phases is increasing, yet its causes still lie in obscurity, and, except in early phases, we are gaining but little power of relief. To some amendment we can attain, but in the later issues, and unfortunately these are too often the first symptoms

* The amounts of protein and salt taken in the diet were not calculated, but the diet was a very light one.

to drive the patient to seek medical advice, cure is out of the question. Venesections do some temporary good—and are useful at critical moments—but frequent repetitions hardly prove acceptable or satisfactory. Calomel still holds its ground as perhaps the best single drug. The high-frequency current (diathermy) is the best of the "symptomatic" remedies, but too often I fail to persuade the x -ray specialists to take much interest in invisible results. In the long run intermittent courses of calomel, diet, regulated life, diversions of mind, and an annual visit to one or other of certain spas seem at present to be all we can do. C. A.

THE CUTANEOUS TUBERCULIN TEST (PIRQUET):

WITH REFERENCE TO ITS FAILURE IN ADVANCED
TUBERCULOUS DISEASE IN CHILDHOOD.

BY

CHARLES McNEIL, M.A., M.D., F.R.C.P. EDIN.,

PHYSICIAN, ROYAL EDINBURGH HOSPITAL FOR SICK CHILDREN; LECTURER
ON DISEASES OF CHILDREN, UNIVERSITY OF EDINBURGH.

THE cutaneous tuberculin test (Pirquet) is now accepted as a delicate indicator of tuberculous infection in childhood—so delicate, indeed, as often to reveal a latent focus, buried out of reach of clinical methods of detection, and producing no effect upon sound health. On the other hand, it is common doctrine that the test fails in a considerable number of cases of advanced and widespread tuberculous disease in childhood. It is enough to say that these failures of the test are admitted by Pirquet himself,¹ by Calmette,² and are confirmed by many published investigations, of which those of McLean and Jeidell,³ and Dunn and Cohen,⁴ may be quoted as representative and recent.

It would seem rash to question such weighty scientific teaching and general clinical experience, and in a previous communication to the *BRITISH MEDICAL JOURNAL* (1909)⁵ I have myself subscribed to the frequent failure of the test in advanced tuberculous disease. But for some time I have used a more careful technique in the performance and reading of the Pirquet test, and with this I have not failed to obtain a positive reaction in any case of severe tuberculous disease proved to be such by the discovery of the bacillus or by *post-mortem* examination. It seems worth while, therefore, to describe this technique and to give some examples of its results.

TECHNIQUE OF THE PIRQUET TEST.

There is great variation in the directions given for the performance and reading of the test. Undiluted tuberculin is now generally advised, but a dilution is permitted by Calmette and others. For the preparation of the skin, multiple scratches or a single scratch, or the use of a dental burr or small chisel (Pirquet), are variously recommended. The test is advised to be read as positive or negative in twenty-four hours, or at latest forty-eight hours, though it is admitted that late reactions may occur after this time. The control test is made by a similar lesion on the dry skin and the application of a drop of normal saline. The technique which I now use, and which is, I think, more thorough and less free from error, is as follows:

1. *Undiluted Old Tuberculin (O.T. Koch) must be used.*—In my first series of cases, published in the *JOURNAL* in 1909, a 25 per cent. dilution of old tuberculin was used, and it is very probable that some of the failures there recorded were due to this. Calmette himself still advises such a dilution. In another paper⁶ I have published photographs showing the simultaneous use of undiluted tuberculin and 25 per cent. dilution in two cases, and the diminution of the test in the one case and its failure in the other with the diluted tuberculin. It is necessary, therefore, to emphasize the importance of using undiluted old tuberculin for the cutaneous test.

2. *Preparation of an Area of Skin to Ensure Contact of the Tuberculin with Vascular Tissue.*—The method of the single scratch, or of a number of scratches, is too haphazard. Pirquet's method of placing a drop of tuberculin on the skin and of making a lesion by rotating through it a small drill or borer, may also fail to reach and expose the vascular cutis. I always use the following method: A small circular area of skin on the front of the forearm is chafed with the sharp point of a darning-needle until the epidermis is removed and the pink cutis vera is exposed. Care is taken to avoid bleeding. The eye of the needle is now charged with un-

diluted old tuberculin, and pressed into the prepared vascular surface with a rotary motion. A small bruised pit is thus formed from which the tuberculin is rapidly absorbed. This procedure is similar to that advised by Pirquet; but it differs in the careful preliminary excoriation of the skin, and thus renders more certain the contact of tuberculin with the skin capillaries; this difference is enough, I think, to secure success in cases where the tuberculin reaction is feeble. In seven cases in which both methods were tried side by side the reaction was positive in five with the method just recommended, and in only two with Pirquet's method. The reaction is a vascular reaction, and it is only after a careful removal of the tough epidermis that contact with the blood vessels of the skin can be ensured.

3. *The Time-Reading of the Test.*—The routine in clinical work is to read the test as positive or negative at the end of twenty-four hours, and later observations are seldom made. This is another source of fallacy. It may be granted that in a large majority of cases a positive reaction is visible after one day; but it is quite common for the reaction to be delayed until the end of two or three, or rarely four days. Therefore, though a test may be pronounced positive at the end of one day, it is never safe to record it negative until four days have elapsed. I shall give presently an example (Case 1) of how easy it is to make mistakes when, in the routine of clinical work, the arm is examined after twenty-four hours, is recorded as negative, and is never looked at again.

In a previous paper⁵ I suggested the definition of a positive reaction as a "red papule that persisted for seven days." Since that was written in 1909 my subsequent experience has confirmed me in this definition. Calmette² states that the reaction generally disappears in "a few days." I have watched the behaviour of many positive reactions for weeks. Even in feeble reactions I have never failed to see a dark discoloured papule at the end of two weeks; and it is common to see clear evidence of a positive reaction at the end of a month. The tuberculin cutaneous reaction is a chronic enduring reaction: it may be slow in appearing, but it is always long lasting; and in that respect it shows its kinship to the slow inflammation of a true tuberculous process. There need be no fear that if a test has not been read until after a week a positive reaction may have come and gone in that time. It is worth repeating that while a positive reaction may be recorded after a day, a negative reaction (and this is the more valuable clinical aid) may not be accepted until at least four days have elapsed. It would be a safer, though perhaps less convenient, way to read the test at the end of a week rather than at the end of a day. In all important and doubtful cases the practice of observing the arm daily for a week is to be recommended.

The size of the papule is of little importance. It generally exceeds 5 mm. with the above technique; but in a few cases is less than that, and yet is to be regarded as positive if it persists for a week.

4. *The Use of a Control.*—Up to a few months ago the control I used was a 50 per cent. solution of glycerin and water. Since then I have used normal horse serum, which has proved a more severe control, as it gives not seldom a small inflammatory papule. This, however, has always disappeared in four days. But its appearance makes it necessary to delay the recording of a small or doubtful positive reaction until this time has elapsed; if this is not done, pseudo-reactions, sometimes called traumatic reactions, may be accepted as true positives. It will thus be seen that the practice of reading the cutaneous reaction at the end of one day may introduce two kinds of fallacy: it may omit late and feeble positive reactions which may develop after a day; and it may include false positive reactions which will have disappeared in a few days. But the use of a normal serum control ought to guard against the latter fallacy.

There are obstinate critics who argue that any albuminous substance rubbed into the skin may produce an inflammatory reaction, and who put no faith in the specific character of the tuberculin test. Such an argument can neither be proved nor disproved. But *post-mortem* evidence, which has now been produced in great amount, makes it reasonably certain that a positive cutaneous reaction indicates a tuberculous focus in the body.

Alleged Failure of Test in Advanced Tuberculous Disease.

I shall now cite a few examples from my own experience to show that, with the technique just described, positive reactions can be obtained in cases of massive and advanced