the phlegm, and I had not a single death from tuberculosis to record down to the year 1920. Moreover, I had one striking control case.

A young boy was brought to us. He had positive physical signs at one apex, and at the Lister Institute a few tubercle bacilli were found in his phlegm. We treated him for two months and he gained 2 lb. and improved very much. Unfortunately we notified the case. Thereupon a lay visitor went to the house and urged the mother to take the boy to Brompton Hospital. I lost sight of the boy for more than a year, when the mother told me the tragic story. At the instance of the lay visitor the boy went to Brompton and was six weeks in the hospital for observation. The mother was then told that the boy had not tuberculosis. (Tubercle bacilli had been found by us.) When I saw the boy again he was in a hopeless third stage, and he died in a few months. Our diagnosis was tragically confirmed.

Thus the only early case that we were not allowed to treat died within two years; surely a striking control. These results obtained by me in England almost exactly tally with the results I have published in my book.³ My records in Australia and London are quite good enough to give me unswerving faith in large doses of tuberculin given subcutaneously as easily the best remedy for tuberculosis in all its forms.

REFERENCES.

¹ BRITISH MEDICAL JOURNAL, March 24th, 1923, p. 493. ² Ibid., March 31st, 1923, p. 556. ³ *Tuberculin in the Diagnosis and Treatment of Tuberculosis.*

THE TREATMENT OF PNEUMONIA.*

BY

A. G. NEWELL, M.D., D.P.H.,

HARRINGAY.

PNEUMONIA causes more deaths than tuberculosis, and more people die from it between the ages of 35 and 55 than at any other period. It carries away insidiously the aged without an acute attack. The death rate is put at 20 per cent, and with influenza at 30 per cent. This means one death out of every five or six patients; annually over 40,000 persons die from it.

Pneumonia at its outset is a septicaemia caused by a diplococcus. All its symptoms are the result of the influence of the bacillary toxin on the nervous system. Its entrance is always at some local focus of the lung evidenced early by dullness and diminished air entry, but its spread is often very rapid. Hence the importance of the early localization and diagnosis and active measures.

The early attempt at defence is consolidation, to limit the focus of infection; the organisms are surrounded, and there is a leucocytosis whereby the phagocytic action of certain blood-cells destroys the organisms. It must be remembered that the lungs contain a larger number of leucocytes than any other organ, except the marrow (their birthplace) and the spleen (their grave). The clotting of the fibrinous exudate in the alveoli which gives rise to consolidation, and the attack by emigration into this of the leucocytes shows an attempt on the part of Nature to localize the infection and prevent its spread. But leakage occurs, and before consolidation takes place certain organisms and toxins have already spread to the blood, causing the acute septicaemic symptoms. Nature's third attempt at defence is the formation of antibodies. Let me summarize the main principles, as it appears to me, on which we should go in attempting to counteract the septicaemia.

There is acute nerve exhaustion from intoxication. This demands absolute rest. The least movement of the patient is exhausting, especially any efforts by himself. Once the diagnosis is made he should not be moved for examination. He should not be allowed to do anything, and bed pans and urine glasses are essential. With this acute nerve exhaustion there is nerve exhaustion to the heart from the poisoning of the vasomotor centre and of the heart muscle. Therefore the importance of rest is especially demanded by an organ called upon to do extra work by the resistance to the circulation in the lung alveoli. A patient may fall back dead on attempting to sit up. The consolidation of the lung alveoli lessens the oxygenation of the blood, and has therefore thrown out of action a defensive auxiliary of the blood. This demands free ventilation around the patient.

The high temperature, the rapid breathing, rapid pulse, the

pain, all add to the patient's discomfort, and therefore it is essential that his comfort and his environment should be attended to by a skilled nurse, or one to whom all the particulars of the necessary details are made plain.

Various lines of treatment have been adopted in the past:

1. Active bleeding and purgation; its ancient history and objects are well known, and I need not dwell on this.

2. Poulitices and expectorants may be said to have died out.

3. "Expectant" treatment may be summed up as doing what each individual fancies best. Where groups of cases have been treated by certain definite drugs it has not infrequently been claimed that some of them were "specific"; types of cases may vary greatly in severity, according to the virulence of the organism, or the influence of the locality, or of the acquired natural resistance, so that claims founded on statistics are of little value. I believe at present there is no specific drug for this disease. Any drug to be specific, or any combination of two drugs to deserve such a claim, must show equal results in the hands of any practitioner anywhere. By the expectant treatment thousands of lives are lost annually.

4. "Chloroform" treatment has been used in America. Its object was to give rest to the inflamed lung. It has not had, so far as I know, a long run. It aids, in my opinion, the basic factors causing the distress, and can have no influence on the intoxication.

5. The application of ice has been advocated by many, the late Dr. David Lees among others, and has been tried in Dublin hospitals. It demands the earliest recognition of the disease—the localization at its earliest site. Two ice-bags are placed so as to include between them the inflamed lung. It has a scientific and rational basis, and good results have followed its use. The pneumococcus is very sensitive to changes of temperature, and will not live below 28° C., nor above 42° C.—the optimum being 37.5° C. The greater the virulence of the organism the more marked its sensitiveness to temperature. The ice inhibits its multiplication, lowers the patient's temperature, adds to his comfort, and is helpful to sleep. I have often used ice in high temperatures with benefit. In general practice no doubt public opinion would be shocked, but it is of great use to give ice to suck. Chills do not occur. It takes time to reduce high temperatures by rubbing with ice, but the sensation is comforting. I have often seen and used ice for the hyperpyrexia of sunstroke, the whole naked body being on blocks of ice and rubbed with ice.

6. The object of lung puncture is to cause a local accumulation of polymorphonuclear leucocytes by setting up a focus of irritation in the pneumonic areas and so stimulating phagocytosis. It is applying to the lung Kraske's principle in the local treatment of erysipelas, his treatment for the latter disease being to make superficial incisions about an inch or two from the spreading margin all round the affected part. It is considered analogous to laparotomy in the ascitic variety of tuberculous peritonitis. This line of treatment demands accurate determinations of situation and area of consolidated patches; a hollow needle or trocar about two inches long is put into the consolidated patch and left there for thirty seconds. If more than one patch is present, and frequently general practitioners get these cases with many, the largest is selected, and if there are several large patches other punctures are made. Strict asepsis is essential, and the best sites are said to be the axillary lines. Striking results have been claimed by its advocates, but as a method for general practice it does not seem desirable with other measures yet available.

7. Studies at the Rockefeller Institute of Medical Research (referred to in the memorandum on pneumonia issued by the Local Government Board in 1919) led to the recognition of four types of pneumococcus; so far antipneumonia serum has been found effective only in cases due to Type 1. In so acute a disease we cannot afford to wait for the determination of the type, and as the serum has to be injected every eight hours it is out of court in general practice.

A bright hope lies in the success attending vaccine treatment. Recovery depends on the patient providing sufficient antibodies against the bacteria and their toxins in the blood; this and this only is the specific treatment available. Early diagnosis is essential: given during the first twenty-four hours, brilliant results have been obtained. A vaccine must be used early, and it is not necessary to wait for a vaccine to be prepared. By the use of vaccine we are helping Nature, and by accumulating antibodies in the blood we have them available for the lung when the fibrin has been liquefied. In other septicaemias (streptococcus, staphylococcus, gonococcus, *B. coli*, etc.) vaccines have been successful. Should we not

* A paper read before the North Middlesex Division of the British Medical Association.

all adopt this method and see the result on all hands? Is the gravity of the disease and the saving of life not sufficient to warrant its trial? I think it is. Dr. Wynn of Birmingham says:

"I know nothing in the whole range of medical treatment so dramatic as the rapid defervescence which follows the injection of a suitable dose of vaccine to a patient suffering from pneumonia or influenzal pneumonia within twenty-four hours of its onset. The vaccine I have used is made from primary cultures or at most first subcultures. It should not be detoxicated or sensitized. For acute pneumonia it contains several strains of pneumococci. . . . The important points are that it should be given early and in sufficient amount."

He gives 100 million pneumococci to an adult; to a child aged 12 to 14 he gives 40 to 50 million, and to a child aged 2 to 3 years 10 to 20 million. These doses have been arrived at after many years of experience. When such a dose is used during the first day of the disease, in the majority of cases the temperature falls to normal on the day following and no further treatment is required. By each day's delay defervescence is less likely, and the dose has to be repeated every twenty-four hours until the temperature is normal. His results² show:

Day Treated.	No. of Cases.	Recoveries.	Deaths.	Temp. Normal in —	
				24 hours.	48 hours.
				Per cent.	Per cent.
First	28	28	Nil	71.4	85.7
Second	23	22	1	47.85	56.5
Third	22	20	2	50.0	50.0
Fourth	20	15	5	30.0	40.0
Fifth	14	12	2	35.7	63.5
	107	97	10	50.0	65.0

In pneumonia there is no evidence of a negative phase, and so the early large dose is given safely.

Stock vaccines should be used from first cultures and from a highly virulent strain. Even if a case is injected from a stock vaccine and the case does not afterwards develop consolidation, no harm is done. No harm is done by using antitoxin for a suspected case of diphtheria. Sir Almroth Wright, in his work on the use of vaccines in pneumonia, states that inoculation in the form of a single large dose, administered in the incubation period, often arrested the disease and prevented death. I put this method forward with a plea for trial, as it seems to me our best line of action for the future. My results from stock vaccines are striking.

I cannot lay too much stress on the hygiene of the room—free ventilation is essential. Cases have been treated in America in all weathers on the roofs of hospitals, and in the Philadelphia General Hospital, where the cases were of unusual severity, the death rate of 53 per cent. before the adoption of the open-air treatment was reduced after its adoption to 38 per cent. In the Prince Edward Hospital, Sydney, the patients have since 1908 been treated in the open day and night, and have recovered more rapidly and with less complications and less necessity for drugs. I always advise the windows to be well opened, and so long as there is no discomfort from draughts and there is sufficient light covering, no harm will accrue. Overheating of the room should be avoided—the patient is already hot enough; fresh air is the best germicide, and the patient is gasping for a cooler air.

The actual degree of temperature is of little scientific interest, as one can usually tell from the skin and the pulse the existence of a feverish state. Its record is of value to the nurse to avoid too heavy clothing and at a crisis for more. There is a tendency to put on too much covering both by nurses and relatives. It adds to the discomfort of a feverish patient. If the patient is hot it is well he should lose heat. Pneumonia jackets are useful provided they are thin and not too heavy.

Pulse and breathing rates should always be recorded. The respiration-pulse ratio normally is 1:4; but in pneumonia it may be 1:3, 1:2, 1:2.5, 1:1, or even 2:1. It is the index factor of how the patient is going, and helps in prognosis under ordinary lines of treatment. A case is not necessarily hopeless with 1:1. The heart condition decides.

Potassium iodide and creosote have been advocated. With a tuberculous focus it is possible to loosen some fibrous tissue of a semi-healed area. Inunction of 10 minims of pure creosote in half an ounce of water, with or without one drop of oil of peppermint, has been used with success; it causes sweating and, it is claimed, aborts the disease. One of the oldest and best drugs is calomel. Digitalis is recommended from the outset by some, strophanthus by others. Neither of them raises blood pressure, and the cardiac muscle is enabled to act with regular impulse. They are the only cardiac stimulants necessary, and if employed should be used from the outset. Strychnine and alcohol are probably of little avail. Tincture of nuxvomica with digitalis is better than injections of strychnine for a flagging heart. To produce sleep 1/6 grain heroin may be given: 5 to 10 grains of Dover's powder in a saline has been recommended, but I never use it. Paraldehyde in doses of 1 to 2 drachms in very sweet tea is safe. Soap liniment will relieve localized pleuritic pain. For heart failure injection of 10 per cent. glucose is said to be the most valuable measure.

The cyanosis depends on anoxaemia of the arterial blood, and if oxygen is administered properly and by a suitable apparatus, such as Haldane's, it has been proved that this anoxaemia will be diminished and so the cyanosis relieved. Otherwise it is probably useless. Too often it is used as a last resource when nothing could be achieved. It is more important to let the fresh air into the room from the very outset.

As regards diet, milk (citrated if there is vomiting), with soda-water or barley water, or a little flavour of tea, or malted milk may be given. All sugars are good for the heart, so from the outset sugar or honey should be given. Water is essential in small quantities, as also is ice. Two pints of water should be given in the day—albumin water or water flavoured with beef-tea; beef-tea alone is not advisable. The mouth should be cleaned every two hours.

PERNICIOUS ANAEMIA WITH FOCI OF INFECTION IN THE ALIMENTARY TRACT.

BY

VINCENT COATES, M.C., M.D., M.A.CANTAB.,

PHYSICIAN, ROYAL MINERAL WATER HOSPITAL, BATH; ASSISTANT PHYSICIAN,
ROYAL UNITED HOSPITAL, BATH.

It has long been held that foci in the alimentary tract may play an important part in the production of pernicious anaemia; it has not been clearly established exactly how this obtains, and the theory does not reconcile the facts that, whereas pernicious anaemia is comparatively uncommon, septic foci are relatively of frequent occurrence.

Hurst¹ believes that achlorhydria is almost invariably the predisposing cause. He collected 12 cases in which this was known to have preceded the onset of anaemia by periods of from one to twelve years. He considers the absence of HCl, probably in the majority of cases of constitutional origin, to be the important factor, and not the cause of the absence. According to him the achlorhydria not only favours infection but results in the arrival in the intestine of proteins in an unaltered condition. He classifies the possible results which may accrue from achlorhydria thus:

1. Delayed protein digestion, with an undue proportion remaining in the lower part of the ileum, in a form readily prone to bacterial decomposition.
2. Provision of material likely to form toxins, which may be (a) haemolytic, the cause of the anaemia; (b) neuro-toxic, causing subacute combined degeneration of the spinal cord; (c) a toxin stimulating bone marrow to form abnormal types of red blood cells.

The fact that subacute combined degeneration of the cord may appear with little or no evidence of blood changes in no particular detracts from the likelihood of the above theory being practically possible, and this is enhanced in value by the good results Böttner and Werner² obtained by duodenal lavage. Nor does the order in which changes may appear in the blood or in the cord. On the other hand, Biffis³ states that achylia gastrica may be absent, and that it occurs later than the blood changes. Maynard and Sturton⁴ record two cases of pernicious anaemia with foci in the alimentary tract, one with dental sepsis and the other with colitis. The autopsy of the latter showed ulcers in the ileum. They

* *Lancet*, September 2nd, 1922, p. 493.