THE MODERN THEORY OF MAN'S POSTURE.

Thus it will be seen that modern anatomists no longer regard the postural adaptations of the human body as the result of a transformation peculiar to man, as did Lamarck and Darwin. They regard man's gait and posture as the culmination of a series of evolutionary phases which are to be traced in the bodies of orthograde primates throughout the greater part of the Tertiary duration of the earth's history.

THE COMPLEXITY OF THE PROBLEM.

The great extension of our knowledge of the constitution and mechanism of the central nerve system, largely the outcome of the discoveries made by Sir Charles Sherrington during the past thirty years, has shown us that the evolution of the orthograde posture was a far more complex and difficult problem than the great pioneers had supposed. We know how complex and delicately adjusted are the nerve centres which carry out the spinal reflexes of even pronograde apes; with the elaboration of the orthograde posture all of these spinal centres must have undergone a structural elaboration as well as a functional adjustment. Further, the orthograde posture, even in its more primitive form, demands a higher degree of co-ordination of the reflex centres in the spinal cord—centres or clusters which extend from one end of the cord to the other; consequently the higher centres—the red nucleus and other cell masses of the mid-brain, which regulate the postural tone—and the cerebellum, which has so much to do with the "timing" of the spinal centres, must have undergone a high degree of elaboration and expansion. The elaboration of the centres for automatic control of the orthograde muscular system was accompanied by an expansion of the basal ganglia and cortical centres of the brain. During the evolution of the orthograde posture brain and body had to become modified in unison.

The researches of Dr. Leonard Hill have made us aware of the elaborate mechanism needed to regulate the arterial blood pressure when the posture of the body is changed. In the early pronograde primates the vasomotor postural mechanisms must have been already highly evolved, but it is clear that, with each step towards the plantigrade posture of man, there must have been further specializations in the reflex centres which control the distribution of blood and in the structures which support the weight of the blood mass. Further, we shall see that all the structures concerned in the mechanism of respiration underwent a change in form, function, and control. The evolution of the orthograde posture entailed a structural revolution in all parts and systems of the body, from the crown of the head to the soles of the feet.

THE MACHINERY OF STRUCTURAL EVOLUTION.

When we see that a change from the pronograde to the orthograde posture entails a modification of nearly every muscle of the body as well as of corresponding changes in the nerve centres which control them we have to seek an answer to this problem: By what evolutionary machinery does Nature bring about simultaneous and harmonious modifications in a thousand apparently independent units of structure? The explanation offered by Lamarck, and also to a certain extent by Darwin, was that in the animal body there exists a mechanism by which structural modification acquired by one generation as the result of use and habit can be passed on, in ever-increasing degree, from generation to generation. The existence of such a machinery for transmission is not only unproved, but for several reasons is highly improbable. Darwin observed that there was evidence of some controlling factor which correlated the growth of associated structures. The nature of the mechanism which regulates the development and growth of parts and so fits them for performing their function in a new way is becoming apparent bit by bit as we get to know more of the behaviour of animal tissues. Phagocytes are attracted to sources of infection by a chemotactic mechanism. Kappers of Amsterdam has demonstrated that groups of developing nerve cells migrate in swarms and arrange themselves at sites favourable for the reception of the stimuli with which they have to deal; to the mechanism involved he gives the name "neuro-biotaxis." Cells removed from the body of an embryo and grown in artificial media exert a regulating influence on each other; in a sutured wound of the intestine epithelium seeks out epithelium, and muscle cells seek out muscle cells in the process of union. The testicle throws into the blood a hormone which stimulates and regulates the growth of a crop of structures. In acromegaly we see a structural transformation effected in all systems of the body. It is along such

lines as these that we are to be guided to the adaptational machinery which Nature has employed in bringing about evolutionary changes in posture and in the creation of new types of animals. We are only beginning to realize the elaborate growth machinery which is at work in shaping the body of the embryo; the one thing we can be certain of is that this machinery is constituted, not on an anatomical, but on a physiological basis. Further, it is not a supernatural machinery but one which, given means, opportunity, and industry, the mind of man can study and determine.

The Goulstonian Lectures

ON

THE NATURE OF ARTERIO-SCLEROSIS.

DELIVERED BEFORE THE ROYAL COLLEGE OF PHYSICIANS
OF LONDON

BY

GEOFFREY EVANS, M.D.CANTAB., F.R.C.P.LOND.,

ASSISTANT DIRECTOR, MEDICAL UNIT, AND ASSISTANT PHYSICIAN,
ST. BARTHOLOMEW'S HOSPITAL.

[With Special Plate.]

LECTURE I.*

In leaving £200 for the maintenance of these annual lectures, Dr. Theodore Goulston directed that a dead body should, if possible, be procured and two or more diseases treated of. Arterio-sclerosis, as it happens, fulfils both conditions, since it is largely based on *post-mortem* study and includes two or more diseases. It is too often depicted as a permanent pathological change, the final result of a process of arterial decay. There is a mark of finality about these structural changes in vessel walls due to the accidental circumstance of their being chiefly studied after death, due also in part to the degenerative changes being more obvious than the proliferative, and in part to arterio-sclerosis being more common in the old than the young; hence the discovery of these changes during life tends to be accepted as evidence of wear and tear significant of an irrevocable step towards the grave. It is my object to bring forward some facts that will give this slow process of arterial "decay" some vitality and life, for I think it can be shown to be a process of disease with an onset that as yet escapes clinical recognition, and a course in which arrest or recovery are commoner than death.

DEFINITION OF ARTERIO-SCLEROSIS.

The problem of arterio-sclerosis is faced at once by the difficulty of its definition. Osler¹ defined it as "a condition of thickening of the arterial coats, with degeneration, diffuse or circumscribed." This definition states the two essential criteria of arterio-sclerosis—namely, the new formation of tissue and the presence of degeneration; the word "thickening" is better than the word "hardening," which is often used, because it marks the new formation of tissue. Forms of both acute and chronic arterial disease with thickening of the arterial coats but without degenerative lesions are excluded by definition from arterio-sclerosis. Thus periarteritis nodosa, thrombo-angiitis obliterans of Leo Buerger,¹¹ intimal thickening in ligatured vessels, and the endarteritis of Friedländer,¹² in all of which thickening of the walls occurs characteristically without degenerative change, are excluded from arterio-sclerosis.

The definition, however, makes no reference to the nature of the underlying process nor to the problem of causation, and it is therefore too inclusive. In their definition of arterio-sclerosis Thorel³ and Marchand⁴ take into account the underlying process, and in addition to emphasizing the importance of intimal thickening refer to "degenerative changes, inflammatory and productive processes" (Marchand), while Thorel speaks of "hyperplastic changes, and perhaps also inflammatory proliferation." In the second place, arterial lesions whose causation is known, even if they show both thickening and degeneration, do not belong to arterio-sclerosis, but are placed under their specific title. Thus syphilitic mesaortitis, and syphilitic and tuberculous endarteritis do not belong to the group of arterio-sclerosis, though syphilis may have to be taken into account as a factor in the causation of arterio-sclerosis when the anatomical lesion is that of arterio-sclerosis and the possibility of its being syphilitic is suspected but not assured.

* Abstract of lecture delivered on March 6th.

THE PROBLEM OF ARTERIO-SCLEROSIS.

As Sir Clifford Allbutt⁴ has often insisted, arterio-sclerosis is not a disease. The word should only be used in a pathological sense to define a lesion which is common to many diseases. There are three distinct aspects of the problem—namely, the description of the vascular lesion, the relative importance in the production of this lesion of various pathological and physiological processes, and lastly its causation. There is fair measure of agreement as to the description of the lesion; there is, however, no accepted opinion as to the nature of the processes concerned in its production, and its causation is largely hypothetical.

CLASSIFICATION OF ARTERIO-SCLEROSIS.

Arterio-sclerosis is too large a subject to be considered as a whole in detail. It is conveniently divided into four main types based on anatomical changes—namely, nodular, diffuse hyperplastic, senile, and Mönckeberg's sclerosis.⁵ This is Councilman's classification⁶ with the addition of Mönckeberg's sclerosis. The classification of arterio-sclerosis is fully discussed by Sir Frederick Andrewes,⁷ but I have not followed his classification, since several forms of arterial disease, such as syphilitic mesoarteritis and arteries undergoing involution, which he includes under the term "arterio-sclerosis," are excluded by definition from this review.

At the same time the four main forms of arterio-sclerosis mentioned do not include all varieties of arterio-sclerosis commonly met with. There are among others more or less mixed forms that must be placed in the class to which the more prominent lesions belong, and there are forms of diffuse intimal thickening, particularly in the aorta and larger vessels, which are fibrotic rather than hyperplastic, and which seem to undergo fatty degeneration at an early stage in their evolution. For the present I am chiefly concerned with the type of arterio-sclerosis termed "diffuse hyperplastic sclerosis,"^{8,16,22} and its relation to the other three types of arterio-sclerosis will be considered later.

DIFFUSE HYPERPLASTIC SCLEROSIS.

This type of arterial disease is the same as that termed "diffuse arterio-sclerosis" by Osler, "secondary arterio-sclerosis" by Thoma,⁹ and "arterio-capillary fibrosis" by Gull and Sutton.¹⁰ It is found in children¹¹ with renal disease and in adults with chronic nephritis (chronic parenchymatous nephritis) and true chronic interstitial nephritis,⁸ also (as will be discussed later) in association with left ventricle hypertrophy and raised blood pressure, and lastly, in certain cases as an incident when death has resulted from some other not related cause. My observations on the histology of the lesion are based on the opinion that the nature of the arterial lesion and its evolution is the same in these different conditions and diseases, excepting that in the case of chronic nephritis it may be limited in its distribution and may indeed exceptionally be limited to the kidneys. The histology of diffuse hyperplastic sclerosis was the subject of controversy between Johnson and Gull, but Jores has made the greatest contribution to our knowledge of the lesion.^{12,14} Aschoff and Gaskell¹³ have followed up his work, and I have used the latter's method of gelatin imbedding in all my material for microscopic examination.¹⁵ A short résumé of the main features of the lesion as previously described in detail⁸ will now be given, with the addition of the description of the earliest stage of the lesion which I have not previously published.

The Histology of Diffuse Hyperplastic Sclerosis.

The essential lesion is a thickening of the intima in the smaller and smallest arteries. This observation is in complete agreement with Jores, and there can no longer be doubt on the matter. Reference will be made later to some observations on the histology and physiology of capillaries, and it will perhaps be shown in the future with more perfect histological technique that the earliest and most important changes are in the capillary endothelium. The intimal thickening is very uniform in the circumference of the vessel, and it may be so great as completely to occlude the lumen in the terminal arterioles, while the lumen of the parent arteries is often greatly reduced. In serial sections of kidney tissue the intimal change is fairly uniform in the length of the vessel both as regards the terminal arterioles and their parent vessels. There is no fatty degeneration in the small arteries, whereas in the arterioles there is marked fatty degeneration of the swollen intima. In a cross-section of the arteriole stained with Sudan 3 the lumen may appear completely

filled with a plug of fat, or lying a little eccentrically may be seen a rosette of nuclei, the remnant of degenerated endothelial cells.

All vessels in a single organ are by no means equally affected. In slight or early cases the affection is limited to occasional vessels. When only occasional arterioles are affected, even when the lesion is complete and the arteriole lumen appears to be completely occluded with a plug of fat, the term "diffuse hyperplastic sclerosis" is not of course applied to the condition. For example, in the case of an apparently healthy young man, aged 19, who was killed by accident, I found two arterioles in the spleen in this condition; the rest of his cardio-vasculo-renal system was perfectly normal. The lesion is regarded as one of diffuse hyperplastic sclerosis when both arterioles and their parent vessels are affected, and as a rule it is found that the lesion in the parent vessels is more uniform, though it varies in degree, than in the terminal vessels. The changes found in capillaries will be considered later in discussing nephritis; their walls undergo hyaline swelling and focal fatty degeneration, the lumen becomes occluded, and they become empty of blood.

The intimal lesion in diffuse hyperplastic sclerosis is the structural change by which this form of arterio-sclerosis is recognized. Its chief incidence is on the finer ramifications of the vascular system, the terminal arteries and their parent vessels being chiefly or only affected; when the arteries from which the so-called parent arteries are given off are affected, they show similar changes to those seen in the parent arteries, while the intima of vessels of the size of the renal and radial artery is fairly normal in appearance. I have not identified a characteristic change in the larger vessels such as the aorta and its branches apart from the medial hypertrophy which is apparently present in marked cases of diffuse hyperplastic sclerosis associated with a raised blood pressure and left ventricle hypertrophy. In some cases the larger arteries show structural changes characteristic of nodular or senile arterio-sclerosis.

The distribution of the lesion in the several organs of the body shows considerable uniformity. The kidney and spleen are most commonly affected; so far as I have observed the spleen is not affected independently of the kidney, but since there are occasional cases in which the kidney is affected without the spleen being involved, it follows that of the two organs the kidney is rather more often affected than the spleen. The incidence of the lesion in the splenic vessels is a criterion of its general incidence in other organs, the kidney affection being a less reliable criterion because of the association of the vascular lesion with chronic nephritis. The brain appears, from my observations, to be the next organ most liable to this vascular lesion; and after this come the pancreas, liver, suprarenal glands, and, less often, the vessels of the stomach and intestines. I have seen no specimen in which the lesion was present in the heart; 70 cases have been examined, including many in which the lesion was well developed in other organs, hence it follows that its presence in the heart vasculature must be uncommon. The lesion has not been found in skeletal muscle. Other organs have been examined, and the lesion has been found in lymphatic glands and in one case in the ovaries; no example of the lesion has been found in skin, pituitary, thymus, urinary bladder, prostate, or uterus, but too few specimens have been examined to lend weight to the absence of the lesion in these organs.

Nature of the Intimal Thickening.

The sequence of changes that take place in the thickening of the intima cannot be determined by the investigation of the condition of the vessels in a single subject, and it is of course an assumption to describe as occurring in a sequence the various changes observed in different vessels in organs taken from different subjects; at the same time I believe that the changes in the arterial wall occur in the order that I now intend to describe. The first change to occur in the arterioles is a swelling and proliferation of the endothelial cells (Fig. 1). The new endothelial cells become flattened and undergo hyaline swelling, and the next stage in the process is the appearance of fatty granules in the cells (Fig. 2). The stainable fat increases, the granules coalesce, the cell outline is lost, and finally the cell protoplasm is replaced by fat. It is at this stage that the arteriole is frequently seen in cross-section with its lumen plugged with fat and the nuclei forming a tiny rosette in the centre. Ultimately the nuclei disintegrate and disappear, and the last cells to survive are those

nearest to the lumen, perhaps because they are the youngest, most recently formed cells, and because of their situation have the best supply of oxygen. In serial sections of arterioles parts that have undergone hyaline swelling merge insensibly into areas, or rather lengths of fatty degeneration; there is also a variation in the amount of Sudan 3 taken up, and its colour varies between yellow and vermillion red. When the arteriole is traced into the vessel from which it springs the fatty degeneration is seen to stop short at the wall of the parent vessel. This vessel characteristically contains no fat within its walls, and is in this striking contrast to the fatty degeneration in the terminal arteriole.

Changes in the Parent Vessels.

The thickened intima of the parent vessels—a thickening which may be so marked as nearly to obliterate the lumen—consists of cells, fibrils of connective tissue and elastic tissue, and hyaline intercellular substance. The prominent feature of the thickened intima is the amount of hyaline substance present, the new formation of connective tissue, and the fine fibrils of elastic tissue that permeate the whole coat. In active and early lesions there is a considerable increase in the number of cells present, but more often there is a relative paucity of cells, though allowing for the great increase in the thickness of the intima there is even in these cases an increase in the number of cells over those present in the normal intima.

1. *Cellular Proliferation.*—Endothelial proliferation is more difficult to demonstrate in the parent vessels than in the arterioles, but there is no doubt of its occurrence. Fig. 3 shows an active stage of endothelial proliferation in a vessel taken from a section of a kidney with chronic nephritis. Fig. 5 shows a later stage, an appearance more frequently seen; the inner layers of cells are flattened endothelial cells, and they are seen lining the lumen in a close-up view of the same artery shown in Fig. 6. It is probable that some of the cells in the thickened intima arise by proliferation of the subendothelial cells.

2. *Origin of Elastic Fibres.*—Elastic fibrils are laid down by cells morphologically identical with connective tissue cells,²² and in addition to this method of formation it has been suggested by Jores that the elastic tissue of the hyperplastic layer arises by a process of growth and splitting of the internal elastic lamina. The appearance of some sections of diseased arteries suggests the possibility of elastic fibrils being laid down directly by endothelial cells. In a case of endarteritis obliterans (type of Winiwarter) affecting the anterior tibial artery there was great thickening of the intima arising suddenly, almost like a shelf, in the lumen of the vessel. The thickened intima was enormously cellular and the cells had the appearance of flattened endothelial cells, in between which were found fibrils of elastic tissue almost enfolding the cells in their meshes. At one place the endothelial cells were arranged in a number of concentric layers enclosing a lumen, and near the lumen of this vessel, which seemed to be in the process of formation *in situ*, was a thick folded layer of elastic tissue, and lying flat on it, almost like slugs on celery, were endothelial cells.

3. *Origin of Hyaline Substance.*—The formation of hyaline is not without doubt a degenerative process. Adami¹⁹ uses the word "inflammation" as reaction following injury, and the response to it. For the sake of clearness for my present purpose I would add the words "and to the effects of it." In distinction from a process of inflammation a process of degeneration is one in which there is no reaction on the part of the affected tissues to the change in their environment, the changes observed being a passive result of the alteration in environment, and the later phases of these changes are loss of function and cell death. I apply the term "degeneration" when fat appears in cells that do not normally contain it. Thus if an intimal cell contains fat it is regarded as undergoing the process of degeneration, whereas, of course, the appearance of fat in a liver cell has no such significance. It is agreed that the appearance of amyloid and calcification is evidence of degeneration. It is with the interpretation of the earlier phases of the process of degeneration that the difficulties occur. Thus it seems that cloudy swelling and a hyaline change may be either inflammatory in the sense that I have defined the word (that is, in its widest sense) or degenerative. Hyaline change is indeed commonly classed among the processes of degeneration in textbooks of pathology, and is in fact referred to as hyaline degeneration. In many instances the origin of the hyaline

material is difficult to determine, in other cases there is little doubt as to its origin. Thus masses of fibrinous deposit may become hyaline, as may blood coagula in the thrombi of aneurysms, and similarly the thrombi of any vessel may undergo hyaline change. The hyaline change occurring under these circumstances is regarded as a degenerative change, and the clearest example is the change of dead protoplasm washed with tissue fluids to hyaline material.²³ Hyaline material appears also to be derived directly from cells, either as a secretion by glandular epithelium or as an accumulation of hyaline substances in the protoplasm of the cells. It is probable that the hyaline in the intima of the thickened arteries now under discussion arises in this way by a process of secretion by the endothelial cells or by the subendothelial cells. The coincident appearance of hyaline substance and cellular proliferation, the fact that the change does not progress to fatty degeneration, and lastly, the clear outline of the cells in the thickened intima and the good staining of their nuclei, are reasons for refusing to accept the presence of hyaline substance as evidence of a degenerative change. This point of view is clearly stated by J. Lorrain Smith, who writes:

"Hyaline material is deposited in the walls of capillaries, causing thickening and gradually leading to closure of the lumen with atrophy and disappearance of the endothelial cells. This change in the capillaries is seen in senile conditions, and also in chronic inflammation, where it extends beyond the vessel to newly formed connective-tissue fibrils. In these examples it originates chiefly in the endothelium and may be regarded as secretory in form."

4. *Origin of Connective Tissues.*—The question of a degenerative process being involved in the formation of the fine connective tissue fibrils that permeate the thickened intima in diffuse hyperplastic sclerosis does not arise. There is no doubt that the process involved is one of inflammation in the sense in which that word has already been defined; in fact, the definition need not be so wide, and the word in this connexion can be used in the limited sense of "reaction following injury and the response to it." As has already been seen, there is no necessity for regarding the hyaline substance as a degenerative change, and therefore there is no need to explain the formation of connective tissue as an effect of the presence of hyaline substance. It is generally held that the connective tissue which often forms an appreciable part of the intima thickening is derived from the subendothelial cells which by their proliferation contribute to the cell increase in the thickened intima. In addition to its origin from these cells it is possible that some of the connective tissue is derived from endothelium. Thus Baumgarten has made observations that indicate the origin of fibroblasts and connective tissue from vascular endothelium under mild grades of irritation. Sir Bernard Spilsbury allows me to quote his observation of the appearance of connective tissue fibrils immediately under the growing layer of endothelial cells as they spread over a newly formed thrombus on a heart valve; in this latter case there were no cells in the neighbourhood from which the connective tissue could have been derived other than the endothelium.

Summary and Discussion of Changes in Intima, Media, and Adventitia.

Joseph Coats²⁴ maintained that the essential constituent of blood vessels is the intima. The endothelium is the universal lining of the vascular system; every vessel, in fact, has an intima, but only the arteries and veins have a media and adventitia; also in the formation of new vessels the primary event is the formation of a tube composed of endothelium by a process of budding from existing vessels. The paramount importance of the intima in health is maintained in disease, and I believe that the endothelial changes are the key to the interpretation of the lesion in diffuse hyperplastic sclerosis. It may be possible in the future to be more explicit in this statement and to bring forward convincing evidence that the endothelium plays the greater part in the intimal change. Thus specimens have been shown in which there is active proliferation of the endothelium; evidence has been brought forward that raises the question of elastic tissue formation by endothelium; the formation of hyaline material by a process of secretion from endothelium has been discussed; and finally, the formation of connective tissue by endothelial cells may be accepted as a fact. In short, the various constituents of the thickened intima in diffuse hyperplastic sclerosis, the proliferation of cells, the increase of elastic tissue, the formation of connective tissue, and the production of hyaline substance,

FIG. 1.—A terminal arteriole showing proliferation and swelling of endothelial cells; the lumen is almost obliterated and is occupied by a mononuclear cell. The earliest stage of diffuse hyperplastic sclerosis.

FIG. 4.—Same vessel as Fig. 3, stained with Weigert's elastic stain to show new formation of elastic tissue.

FIG. 2.—A later stage of the lesion, showing an arteriole with proliferated endothelium, the outer cells having undergone hyaline swelling and fatty degeneration (shown by black dots, section stained with Sudan 3), with loss of cell outline. The lower arteriole shows a still later stage, in which the lumen is completely obliterated and the cells further degenerated. The media is intact in each arteriole.

FIG. 5.—A later stage of the same process. The endothelial cells are now flattened; the new-formed elastic tissue is more regularly disposed.

FIG. 3.—Endothelial proliferation in a parent vessel with simultaneous formation of elastic tissue. No fatty degeneration present. Early stage of lesion.

FIG. 6.—Close-up view of part of wall of same vessel to show the flattened endothelial cells, the new-formed elastic layers, the internal elastic lamina, the media, and surrounding kidney tissue in this order from above downwards.

may all result from the increased activity of the endothelium. It will be remembered that in pure forms of diffuse hyperplastic sclerosis—and in speaking of "pure" forms of diffuse hyperplastic sclerosis I mean to exclude those cases in which senile arterio-sclerosis complicates the picture—fatty degeneration is confined to the terminal arterioles of the kidney and the smaller vessels in other organs. The presence of fatty change in these vessels will be referred to later, but the point that I would make now is the absence of any change that is beyond doubt degenerative in the vessels of second size—that is, the parent vessels of those affected by fatty change.

In addition to the thickening of the intima there is sometimes hypertrophy of the media in diffuse hyperplastic sclerosis. This is the only change in the middle coat, pathological changes being characteristically absent. The medial hypertrophy cannot otherwise be regarded than as a physiological response to the increased blood pressure that is present in the majority of cases of diffuse hyperplastic sclerosis. I have previously called attention to the fact that diffuse hyperplastic sclerosis of limited extent may be found after death in bodies in which the heart is not increased above 15 oz. in weight, and in which the blood pressure during life was below 180 mm. Hg. Hypertrophy of the media is not evident in these cases. There are also cases of diffuse hyperplastic sclerosis in older cases in which fatty degeneration of the media occurs and the media appears stretched in section and apparently thinner than expected in a normal artery of the same size. I have not made exact comparison in these cases, and can only express the opinion from my observations that medial hypertrophy is a common accompaniment of marked cases of diffuse hyperplastic sclerosis, and that it is present in cases in which there was a definitely raised blood pressure during life or marked hypertrophy of the left ventricle after death, though there are slighter cases of diffuse hyperplastic sclerosis in which no medial hypertrophy occurs. This medial hypertrophy is neither the essential element nor a constant feature of diffuse hyperplastic sclerosis. It is possible that the raised blood pressure precedes the vascular lesion in some cases, and it may be, therefore, that medial hypertrophy precedes intimal thickening under these conditions; in any case the medial hypertrophy is frequently so pronounced that it is regarded by some observers as the essential element in this form of arterio-sclerosis.¹²

Gull and Sutton emphasized the thickening of the adventitia in arterio-capillary fibrosis. This is present in some cases, and I have noticed it in the renal vessels and perhaps also in the spleen. It is difficult to distinguish between an increase in the adventitia and an increase in the perivascular connective tissue which is present in considerable quantity in both these organs. It has escaped my observation in other organs of the body, and needs closer study than I have given it.

Conclusion.

The main feature of diffuse hyperplastic sclerosis is a thickening of the intima, accompanied sometimes by hypertrophy of the media. The intimal changes are the essential lesion. Activity of the endothelial cells is largely responsible for the intimal thickening. In this thickening the process of degeneration plays a secondary part; it is only certainly present in the terminal arterioles of the kidney and the smallest vessels of the other organs. The characteristic feature of the lesion is an intense tissue activity, which is the very antithesis of a process of decay. The nature of the process concerned in the production of this lesion will be considered in the next lecture.

REFERENCES.

- ¹ Osler, Sir William: *Principles and Practice of Medicine*, eighth edition, 1918, 841. ² Thorel, Ch.: *Labarrach-Ostertag*, 1915, 1, 171. ³ Marchand: *Verh. d. d. Cong. f. inn. Med.*, 1904, 21, 58. ⁴ Allbutt, Sir Clifford: *Diseases of the Arteries and Angina Pectoris*, 1915. ⁵ Mönckeberg, J. G.: *Virchow's Arch.*, 1903, 171, 141. ⁶ Councilman, Wm. T.: *Trans. Assoc. Amer. Phys.*, 1891, 6, 179. ⁷ Andrews, Sir Frederick: *Local Govt. Board Reports*, 1911-12, Suppl. App. B. ⁸ Evans, Geoffrey: *Quart. Journ. Med.*, Oxford, 1921, 14, 215. ⁹ Thomas, R.: *Textbook of General Pathology*, translated by Bruce, 1896; *Virchow's Arch.*, 1886, 104, 203; *Ziegler's Beitr.*, 1891, 10, 433; further reference given by Marchand, loc. cit. ¹⁰ Gull, Sir William, *Writings of New Syd. Soc.*, 1894, 325. ¹¹ Evans, Geoffrey: *Quart. Journ. Med.*, Oxford, 1922, 16, 33. ¹² Turnbull, Hubert M.: *Ibid.*, Oxford, 1915, 8, 201. ¹³ Jores, L.: *Wesen u. Entwicklung d. Arterio-sklerose*, Bonn, 1903. ¹⁴ Jores, L.: *Virchow's Arch.*, 1904, 178, 367. ¹⁵ Gaskell, J. F.: *Journ. Path. and Bact.*, 1912, July, p. 18. ¹⁶ Gaskell, J. F.: *Ibid.*, 1912, 16, 287. ¹⁷ Buerger, Leo: *Amer. Journ. Med. Sci.*, 1908, 136, 567; and 1910, 139, 107. ¹⁸ Friedländer, Carl: *Centralbl. f. med. Wissen.*, 1876, No. 4. ¹⁹ Adams, J. G.: *A.butt and Rolleston's System of Medicine*, 1911, 1, 786. ²⁰ Smith, Lorrain: *General Pathology*, edited by Pembrey and Ritchie, 1913, 207. ²¹ Coats, Joseph: *Manual of Pathology*, third edition, 1895, 466. ²² Jores, L.: *Ziegler's Beitr.*, 1893, 24, 4, 8.

An Address

ON

TREPHINATION OF THE LIVING HUMAN SKULL IN PREHISTORIC TIMES.*

BY

T. WILSON PARRY, M.A., M.D. CANTAB., F.S.A.

[With Special Plate.]

SIXTY years ago not even the most eminent scientist in Europe was aware of the fact that, in prehistoric times, our primitive ancestors used to trephine the living skulls of their fellow tribesmen with implements made of stone. The knowledge of this was rendered possible by a discovery made by a general medical practitioner. And here I should like to say that such of you as are about to enter general practice will find, I think, this branch of our profession of a peculiarly satisfying nature, in that not only does it have for its aim the carrying out of ethical ideals by sound practical methods, but there is scope, at leisure moments, for the study of those wider interests that do not actually belong to our profession, but which are stimulated by the atmosphere created by our education, trend of thought, and environment.

Prunières, a general practitioner of Marvejols in the department of La Lozère, France, was a man who spent much of his spare time in antiquarian research. While examining a large dolmen, near Aiguères, in 1865, he discovered a skull with a curious hole in it. The hole had smooth, shelving edges which looked as if they had been polished. Prunières was puzzled as to the nature of this hole and came to the conclusion that, finding the skull in a dolmen, it had belonged to a Neolithic savage and had been converted into a drinking cup, a custom which not infrequently exists among savage tribes, and that this smooth-faced hole had been specially designed for the application of the lips. Now these dolmens were constructed during the third or Carnac stage of the Neolithic period, which ended in north-western Europe about two thousand years before the Christian era. The whole of the Neolithic period is computed to have lasted some twenty thousand years, the third and last division probably occupying from four to six thousand years, so that the age of these dolmens may be estimated as being not less than four and even as much as eight to ten thousand years old—the last word in these megalithic buildings being that magnificent pile known as Stonehenge, which was erected at the beginning of the Bronze Age.

Prunières was not correct in his surmise about the drinking cup. To Broca belongs the credit of explaining that the smooth, polished-looking surface of the sides of the hole in this skull was due to a healing process that had taken place in a wound of the bone during life. After this first discovery many other specimens were unearthed from the dolmens existing in those regions, so many, indeed, as to compel one to believe that, during the latter part of the Neolithic period in France, a fetish of a most extraordinary kind was prevalent among the inhabitants, and this fetish consisted in trephining the healthy human skull for some purpose of a superstitious nature, for none of the skulls showed any sign of either fracture or disease.

And now comes the point when it might very reasonably be asked, How can you definitely prove that these holes, which we now know were made during life, were the result of an operation, deliberately planned and carried out by Neolithic man? Is not this suggestion merely guesswork? Broca, having found out all he could from the nature of the specimen itself, and from the place where it was found, and the associations, turned his eyes in another direction.

Here it may be explained that all primitive tribes, on their way upward from savagery to civilization, pass through the same phases. Every nation starts with its so-called "Stone Age," when the chief material used for the making of their implements is stone. Wood, slate, shell, bone, and teeth are also used during this period. When culture has proceeded farther, metals are discovered, the first being the more malleable ones—copper and tin. By accident or experiment they find that a harder and much more serviceable metal is obtained by blending these two metals by heat, bronze being the outcome—thus the Bronze Age. When iron is discovered a stupendous advance has been made, and the early Iron Age

* Given before the Listerian Society of King's College Hospital on January 17th, 1923.