is that there is a blood infection. This can only be shown by blood culture, frequently repeated if necessary. Dr. W. S. Cosbie, a member of my staff in Toronto General Hospital, took cultures from all puerperal patients running a temperature. His investigations are not completed, but we were surprised to find that so many of them had positive cultures at one time or other. Most of these made a complete recovery without special treatment.

In the treatment of true cases of bacteraemia we have used various kinds of intravenous medication such as magnesium sulphate, eusol, and salvarsan. We have also used serums. With all of them we have had recoveries, and at first attributed these to the drug used, until we found that, with no intravenous medication and with no serum, just as many patients got well. The great difficulty in assessing the value of these forms of treatment is that spontaneous recovery so often does take place.

What I have said regarding the treatment of puerperal infections occurring after a full-time delivery applies equally to those arising after abortion, either complete or incomplete. I make it a rule never to curette the uterus or carry out any intrauterine manipulation in a case of abortion if the patient has a rise of temperature, unless there is severe haemorrhage. If the abortion is incomplete and a mass is felt projecting through and blocking the cervical canal, this may be very gently removed, but more interference than that I think unwise. After the temperature has been normal for some days, if there is still evidence that the abortion is incomplete by the persistence of bleeding and the patulous condition of the cervix, gentle curetting may be carried out. Even after waiting this time it is almost invariably found that the patient's temperature rises after the operation, and she not infrequently has a rigor, showing almost certainly that there has been a blood invasion. Fortunately, in most cases, this is only temporary.

I have only touched very briefly on the whole subject of puerperal sepsis, but my purpose is served if I initiate some discussion on the inadvisability or otherwise of active interference with the interior of the uterus in the presence of fever. From all the evidence of the literature and from my own experience I hold that it is inadvisable.

THE USE OF QUINIDINE IN AURICULAR FIBRILLATION.*

BY

FRANCIS R. FRASER, M.D., F.R.C.P.Ed.,

DIRECTOR, MEDICAL PROFESSIONAL CLINIC, ST. BARTHOLOMEW'S HOSPITAL, LONDON.

THERE have now been recorded several hundreds of cases of auricular fibrillation in which normal rhythm has been restored, and it can be accepted as demonstrated that quinidine can bring about this change of mechanism in a large proportion of all cases.

Like all powerful drugs, quinidine can produce effects that are unpleasant, or even dangerous, and so it is necessary to learn in what cases the conversion can be effected without producing such results. The advantages to be gained might conceivably be so great that the risk of unpleasant symptoms would not be a serious deterrent to giving the drug, but in digitalis we have a method of treating the effects of auricular fibrillation that is so successful that it must be seriously considered in what cases a successful quinidine result surpasses in therapeutic value the effect of digitalis treatment.

Answers to the two following questions must be obtained before quinidine is to find a place in practical therapeutics:

- (a) In what cases can the conversion be brought about without risk of producing dangerous symptoms?
- (b) In what cases does the therapeutic effect of this conversion surpass the effect of digitalis treatment?

Discussion may well help to answer two further questions:

- (c) What is the best method of administration in regard to dosage?
- (d) For how long after the conversion should the quinidine treatment be continued?

Many other questions must arise, but preliminary answers to these four are desirable before the necessary amount of

evidence will be collected on which sound therapeutics can be based.

Among the earlier cases in which I tried quinidine was a woman, aged 54, with a greatly enlarged heart, severe mitral regurgitation, and stenosis of rheumatic origin, and auricular fibrillation. Quinidine sulphate (5 grains night and morning) caused auricular flutter to develop, and after eight days, during which the auricular rate was about 200 and the ventricular rate 80 to 90, the pulse rate rose suddenly to 200, probably as the result of a 1:1 response. On stopping the quinidine and commencing digitalis fibrillation returned.

A woman, aged 31, with mitral stenosis of rheumatic origin and severe heart failure, at rest in bed had a ventricular rate of only 50 to 65. After five days of quinidine sulphate (5 grains three times a day) auricular flutter resulted, and after six days more on 7½ grains three times a day she had a Stokes-Adams attack with ventricular standstill.

A woman, aged 54, with a greatly enlarged heart, mitral regurgitation, and auricular fibrillation, evidence of arterial and renal disease, and a ventricular rate of 180, responded moderately to digitalis. At a ventricular rate of 90 numerous ventricular extra-systoles occurred. With quinidine sulphate (15 grains at intervals of six hours) normal rhythm resulted after four doses, but was accompanied by precordial distress and nausea. On stopping the quinidine auricular fibrillation returned a day later with relief from the subjective distress.

Another limitation to the administration was seen in a woman, aged 48, with mitral stenosis of unknown origin and auricular fibrillation of probably only a few weeks' duration. The response to digitalis was good, but after nine days of quinidine sulphate, during which she received 175 grains in all, a severe toxic erythema developed with headache, nausea, and fever.

After the experience with these cases careful selection was made, avoiding all cases with evidence of much damage to the cardiac structures such as great enlargement, heart-block, or severe valvular disease.

A man, aged 39, with syphilitic aortitis and severe heart failure but no evidence of valvular disease, responded well to digitalis and received antisyphilitic treatment. Three months later normal rhythm was restored after ten days, during which 180 grains of quinidine sulphate was given. The treatment was continued for three months on diminishing doses, and nearly two years later the rhythm is still normal except for ventricular extra-systoles, and he feels fit to return to his work as a stevedore.

In the case of a man, aged 26, with no clear evidence of cardiac enlargement or of valvular disease and no evidence of infection other than malaria, a successful quinidine effect has continued for twenty months, and he is now working as a labourer. The auricular fibrillation had been present for six weeks before admission, and the digitalis response was good.

A similar successful result was seen in a man aged 40 with exophthalmic goitre, without evidence of valvular disease, in whom auricular fibrillation had been present probably for five years, and was still present three weeks after partial thyroidectomy by Mr. Dunhill. Normal rhythm resulted after four doses of quinidine sulphate (5 grains three times a day), and has persisted with gradually increasing efficiency for five months.

A woman, aged 45, with exophthalmic goitre of a mild degree had probably had auricular fibrillation for some years. Severe heart failure responded well to digitalis, and normal rhythm resulted after six doses of quinidine sulphate (7½ grains three times a day). The dosage was gradually reduced and treatment discontinued four weeks later. An increase in the thyrotoxic symptoms was accompanied by a return of fibrillation. Partial thyroidectomy was performed by Mr. Dunhill, and as fibrillation was still present a week later quinidine was given, and normal rhythm returned after four doses of 5 grains three times a day.

Another case of this nature shows similar features.

A girl, aged 26, with mitral stenosis of rheumatic origin, responded well to digitalis objectively but continued to have severe dyspnoea and palpitations. Quinidine sulphate (5 grains three times a day) restored normal rhythm after seven doses, with relief from the palpitations. This dosage has been continued for nearly a year now, and a severe relapse occurred during a return of the rheumatic infection. Relapses of a few days' duration have occurred on three other occasions; on one she had a bad cold, on another much worry in connexion with the death of her mother, and on the third she tried to do some heavier work than usual.

There are cases of thyrotoxic auricular fibrillation in which normal rhythm was restored a few days after partial thyroidectomy without other interference, and cases in which auricular fibrillation occurred only for a few hours after operation during the stage of increased excitement. They suggest that the auricular fibrillation is due to an active toxic action, and that normal rhythm may result simply from the cessation of this action. The difficulty in obtaining a stable normal rhythm with quinidine in active toxic cases, and the stability obtained after thyroidectomy, indicate that in this type of case to obtain a stable normal rhythm the thyrotoxic factor must be treated.

In the other types of cases the indications of a similar active agent are not so complete, but the accumulation of more evidence may decide in favour of a similar state of

* Abstract of remarks made in opening a discussion at a meeting of the Therapeutic Section of the Royal Society of Medicine, March 13th, 1923.

affairs in them also. The stability of the syphilitic case in which antisyphilitic treatment was given supports this view. The stability in the case in which no definite cardiac lesion could be found and in which the only known infection was malaria may well be due to the cessation of the causal factor, whatever it was, before the patient came under observation. The instability of the rheumatic case, even when quinidine treatment has been continued for nine months, is suggestive of a constantly or intermittently active factor tending to produce fibrillation. This idea is fully in keeping with the theory derived from the clinical study of such cases—that the rheumatic infection may continue for years, causing gradually increasing stenosis of the mitral valve, often without well defined attacks of acute rheumatic infection. In such a case it is of interest to note the additional factors present during the relapses into auricular fibrillation. Unfortunately there is no successful method of treating the rheumatic infection of the heart.

In addition to active agents it is probable that auricular fibrillation can result from permanent structural changes in the heart, in which case a stable normal rhythm after cessation of the quinidine administration would be unlikely.

CONCLUSIONS.

1. Quinidine treatment will result in a stable normal rhythm if the cause of the auricular fibrillation can be successfully treated or has ceased to be active.

2. Cases in which there are extensive structural changes in the heart are not suitable for quinidine treatment, especially if the response to digitalis is not good, as dangerous symptoms may result.

3. In some cases at least the comfort and efficiency of the patient is greater when a normal rhythm results from quinidine treatment than when digitalis treatment alone is used.

4. In the cases in which the results are satisfactory larger doses than 10 grains of quinidine sulphate at intervals of six hours were not necessary, nor smaller initial doses than 5 grains three times a day.

5. If necessary to maintain normal rhythm, quinidine may be continued to be administered indefinitely in doses of at least 5 grains three times a day, and it is probably advisable in all cases to continue it for some weeks after normal rhythm has resulted, though in gradually decreasing doses.

Memoranda:

MEDICAL, SURGICAL, OBSTETRICAL.

FIXATION OF CAECUM IN CHRONIC INTUSSUSCEPTION.

The following case of chronic intussusception which took an acute form came under my care in November of last year, and seems of sufficient interest to warrant publication.

N. C., a boy aged 4 years, was admitted to hospital on November 16th, 1922. The mother gave the following history:

On November 4th the child complained of severe pain in the region of the stomach. The pain was attributed to worms, and no significance was attached to it. The pain, however, recurred at varying intervals for the next few days. On November 7th the child vomited. The bowels had been quite regular, though once the mother noticed some "slime" in the stools. On November 11th the child was given two "yellow worm powders" and a large dose of castor oil. Two days later two round worms were passed. He still complained of pains, and cried almost incessantly until November 16th, when he was brought to hospital.

On admission his general condition was good. An indefinite tumour could be felt below the right costal margin, and there was loss of resistance in the right iliac fossa. He did not complain of pain. On November 17th and 18th the child had normal motions, but slight abdominal pain. On the 19th his stools were blood-streaked, the pain was worse, and a good deal of mucus was present in the motion. A typical sausage-shaped tumour could be felt in the region of the descending colon.

Operation.

The apex of the intussusception was in the left iliac fossa, about the beginning of the sigmoid, and was reduced with some difficulty. There was some extravasation of blood in the mesenteries of the small intestine and ascending colon, and the vessels were very much engorged.

An area about the size of half a crown at the apex of the caecum was grey and sodden. The appendix was healthy. The caecum and ascending colon were very mobile and the mesentery of the small intestine long. Owing to the congested state of the mesentery and ascending mesocolon I did not like to put stitches

in them, for fixation purposes, so I decided to use the appendix—(1) for fixation of the caecum, (2) by traction on it to turn the doubtful apex of the caecum up against the anterior abdominal wall, in case it should prove non-viable. I ligatured and divided the meso-appendix, and brought the appendix out through a stab wound just below McBurney's point, fixed it with a couple of sutures, and put a gauze wick down to the apex of the caecum, which was directly under the stab wound.

In forty-eight hours I removed the appendix level with the skin and withdrew the gauze drain; I also destroyed the mucous membrane with a probe dipped in pure carbolic.

The boy made a good recovery. The appendix required a second application of carbolic, as the mucous membrane was rather redundant, and secreted a little moisture.

A seven-hour picture after an umbrose meal showed the caecum in the position in which it was placed just below McBurney's point. It would be interesting to know—

1. In what percentage of cases there is recurrence of intussusception after operation.

2. If the risk of recurrence is enough to make an attempt at fixation worth while.

3. If fixation of the caecum by the appendix is suitable for prolapsed or mobile caecum, etc.

Londonderry.

J. B. ALEXANDER, M.B., B.Ch.

SUBOCCIPITAL MENINGOCELE SUCCESSFULLY REMOVED.

The following case, in which a large meningocele which was present at birth was successfully removed by operation, is, we think, of sufficient interest to warrant publication.

Marie P. was born on May 19th, 1921, with a large meningocele, the birth being uneventful and taking place before the arrival of a doctor. The baby was otherwise healthy. It was found that the growth steadily increased after birth, and, the covering becoming thinner, it was obvious that rupture was imminent. It was decided to operate, and the baby was removed to North Lonsdale Hospital. Before operation the dimensions were: horizontal 6½ in., vertical 5 in., circumference at junction with skull 7 in.

Operation.

On August 1st, 1921, assisted by Dr. T. E. Roberts, I dissected back the skin and exposed the underlying membrane. This was very difficult near the neck of the sac, as there was practically nothing between skin and membrane. With a large opening

Photographs showing the child before and after removal of the meningocele.

between sac and the skull there was danger of collapse from sudden loss of cerebro-spinal fluid; to obviate this an intestinal clamp was loosely applied to the neck of the sac. The sac was then opened at a distance to verify the absence of nerve tissue. The clamp was quickly closed, thus preventing leakage of fluid. The sac was cut away and a continuous suture applied round stump and clamp, as in Barling's operation for piles. The clamp was withdrawn and suture quickly tightened. A few interrupted sutures were applied where fluid was oozing. The deeper structures were drawn over the stump and the wound closed. An anchor dressing was applied.

For a few days the temperature oscillated between normal and 102°, with muscular twitchings suggestive of impending convulsions. Bromides were given, and the temperature came down to normal and all twitchings ceased. Owing to fear of yielding of scar the anchor dressing was kept on for two weeks and cut its way out, leaving a granulating surface which did not involve deeper structures. Secondary suturing closed this, and complete healing quickly followed. The baby was sent home well five weeks later, and has remained well and normal since. She has just recovered from whooping-cough and is beginning to walk.

I am indebted to Dr. Kathleen Rutherford for the photographs, which show the baby before and after operation, also for her very helpful interest in the case. The ultimate recovery of the baby was largely due to the skilful nursing of Sister Levison.

JOHN LIVINGSTON, M.D.Durh.,
F.R.C.S. Edin.,
Honorary Surgeon, North Lonsdale
Hospital.

Barrow.