to the conclusion that a difference of 30 mm. or upwards in the sphygmometer readings obtained from the two upper limbs speaks strongly in favour of aneurysm; that a difference of 20 mm. may mean either aneurysm or dilatation of the aorta; and that with a still slighter difference may be found either of the conditions just mentioned, or arteriosclerosis, or mediastinal tumour. Broadly, the position is that to be of diagnostic value the inequality between the measurements must be a pronounced one, and even then, as

FIG. 5.—Sphygmometer: Right, 40; Left, 160.

shown by the case from which Fig. 5 was taken, the possibility of a unilateral interference with the arterial circulation other than aneurysm must be remembered. Further, such differences as are illustrated by the readings recorded in Figs. 1, 3, and 4 are unusual even in cases of aneurysm, and frequently—probably in the majority of cases—the readings are identical or approximately identical. A parallel remark applies to the other pulse differences noted in this paper. Neither unilateral delay in the pulse wave, nor modification in the height or in the form of the wave, is to be observed as a frequent clinical experience in thoracic aneurysm. The absence of such conditions from an individual case is therefore of little diagnostic moment, and while their positive values, when present, may be considerable, even these are subject to such qualifications as are here described and illustrated.

REFERENCES.

- ¹ *Practitioner*, December, 1909. ² *Lancet*, 1907, vol. ii, p. 1516.

A NOTE ON THE NATURE OF THE BLOOD SUGAR.

BY

L. B. WINTER AND W. SMITH.

(From the Biochemical Laboratory, Cambridge.)

IN a recent paper¹ we gave the results of an investigation into the nature of the blood sugar by a new method. Hewitt² has lately criticized both the method employed and also the interpretation placed thereon. He states that we "reached the conclusion that the normal blood sugar is γ glucose." In discussing the experimental evidence which we brought forward in our original paper we stated: "The nature of the normal blood sugar we leave for the present undecided." Later, in view of the fact that Hewitt and Pryde³ had shown "with some certainty"² that γ glucose is formed from α , β glucose in equilibrium of the intestine, we suggested that the unstable sugar which was found to be present might be γ glucose. We stated further that "The enzyme responsible for the conversion being apparently absent from the blood, and the reaction of the blood being alkaline, suggests that it is impossible for so reactive a sugar to exist free in the blood."

It is obvious therefore that our only reasons for ever suggesting the presence of γ glucose were based on the work of Hewitt and his collaborators. They state⁴:

"It may be presumed on *a priori* grounds that a structural change of the nature referred to has some fundamental reason." The reason was not clear, but a possible explanation was afforded by the differences we showed to exist between the normal and diabetic blood sugars, and more recently by the fact that the nature of the diabetic blood sugar becomes similar to that of normal persons after injections of insulin.⁵

Hewitt² objects: "Polarimetric evidence alone, therefore, must exclude rigorously the presence of β glucose in amount greater than that found normally in aqueous solution of glucose in equilibrium." Control experiments showed that the equilibrium of α , β glucose was hardly affected by the experimental technique; further, after the alcoholic extraction no trace of copper-reducing substance was found to be left behind. Preferential extraction by alcohol or alteration in equilibrium of α , β glucose is therefore definitely excluded. In some experiments the initial rotation was below that which would be given by β glucose alone, and in the case of diabetics above that of α glucose alone. The suggestion that the normal blood sugar contains β glucose, and diabetic blood α glucose, is therefore not valid. Further, in one experiment a laevo-rotation was observed, but as this has not been repeated no stress has been laid on it; in this connexion it is of interest to note that Hewitt and Pryde attach considerable importance to the one laevo-rotation observed in their own experiments.

The fact that we obtained a glucosazone from normal blood may be due to the fact that some of the normal blood sugar is the stable α , β form. It seems probable that normal blood sugar is an equilibrium mixture of α , β glucose, and a reactive sugar.⁵

The question of sterility is disposed of by the following facts: (a) the polarimeter tubes were sterilized; (b) the copper-reducing value was unaltered at the end of three or four days; (c) the fluid was perfectly clear throughout, and no sign of any growth was observed.

With regard to the criticisms on the general technique adopted, though admittedly it is not perfect, we maintain that by it we have been able to obtain a clear and concentrated protein-free filtrate from blood by which we are able to show with ease the very wide differences in the nature of the blood sugar under different conditions. We may merely mention here that the nature of the diabetic blood sugar differs from normal, and that after administration of insulin the blood sugar of the diabetic approaches normal; and further, that after administration of adrenaline the blood sugar differs again from normal. A most striking difference is observed in the blood sugar of normal animals after insulin (unpublished experiments); the blood sugar in this condition, though dextro-rotatory, has no copper-reducing value, as determined by the Wood-Ost method.

The downward change in optical values of the diabetic blood sugar is not always observed, and it would appear that in this pathological condition the blood sugar may be very complex.⁶

Hewitt criticizes the experiments in which we endeavoured to test whether there is present in the blood an enzyme capable of converting added α , β glucose or *D*-fructose into a reactive sugar. The experiments of Cooper and Walker,⁷ directed to the same end, are also not allowed to pass unchallenged. The first point he makes is that the blood was extravascular; criticism on this line is purely destructive and would invalidate a vast amount of work which has been done with blood. A more legitimate objection which he makes against the experiments of Cooper and Walker is that an enzyme, if present, was probably unable to pass through the membrane of the dialyser owing to its physical properties. Our own experiments he dismisses on the ground that the method employed was in every way suitable to destroy γ glucose; he omits to notice that if differences in the blood sugar could be demonstrated by this method, the same method would probably be capable of showing a difference had any change taken place in the nature of the sugar which had been added to the blood. He agrees, however, "that no such enzyme can be demonstrated in the blood,"² and

refers to a paper by De Souza and himself.⁴ They injected sugars into the blood stream, and collected the urine passed immediately afterwards. The one merit of the method employed is that the blood is not extravascular, but as a means of demonstrating the presence of such an enzyme in the blood the method does not lend itself to any conclusions. It is unlikely that more than a proportion of the injected sugar would be changed into a reactive form in the time allowed (five minutes); further, the sugar excreted would presumably be that which was unchanged, and therefore useless to the body during that period; further, if any change was observed it could not be a proof that an enzyme was present in the blood.

After a careful study of the figures given by Hewitt and Pryde in their paper,³ it is reasonable to conclude that the rotations observed with isotonic and hypertonic solutions were of the same degree as those observed with hypotonic. It is stated that in the case of hypertonic and isotonic solutions a specific rotation below $+40^\circ$ was not recorded. No figures are given, but this statement implies that changes of that order were observed. In the absence of fuller details, it would appear from the figures 1 and 2 that only about a quarter of the sugar originally introduced into the intestine was present when the fluid was removed for polarimetric examination. Supposing that a quarter of the sugar remains in the intestine when an isotonic or hypertonic solution is introduced, then the observed alteration in rotation required to change the *specific* rotation from $+52.5^\circ$ to $+40^\circ$ would be 0.34° for the isotonic solution, and 0.50° for the hypertonic solution of glucose employed, or alteration of rotation (not *specific* rotation) very similar to, or even larger than was observed, according to the figures, which are stated to "show the optical changes most markedly." It is a curious fact that marked changes should have been observed in hypotonic, and not in isotonic solutions. The authors state: "Contact with the mucous membrane of the living intestine probably effects the production of γ glucose in excess of any amount normally present in glucose solutions which have attained a permanent specific rotation." Were it a genuine effect due to an enzyme in the intestinal cells, it would be likely that the most marked change would be observed in the isotonic solution in the short time allowed (five minutes), since, presumably, changes in the body preferably occur in fluids isotonic with the tissues—for example, perfusion experiments are always performed with approximately isotonic solutions. On the above reasoning precisely similar alterations in *observed* rotation (not *specific*) must have taken place in hypo-, iso-, and hyper-tonic solutions of sugars, but only in the case of hypotonic solutions do these differences lend themselves to the elaborate treatment which they have received in this paper, owing to the fact that a small difference in *observed* rotation of a hypotonic solution makes a large difference in *specific* rotation. A conclusion derived from neglecting inconvenient facts with the acceptance of others is of questionable use. The conclusions drawn from these facts were only given in the following paper,⁴ in which they state: "It may be presumed on *a priori* grounds that a structural change of the nature referred to has some fundamental reason."

Further doubt is thrown on the work by the admitted difficulty in seeing through the fluid for polarimetric observation, a difficulty which becomes obvious when the wide variations in the successive readings are noted; incidentally, the readings are given to three places of decimals. Small reliance therefore can be placed on the one laevo-rotation observed. No copper-reducing value of the sugar is given; the statement is merely made that it agreed remarkably well with the final rotation.

The one experiment recorded in which fructose was used may be adequately explained by temperature variations, to which this sugar is notoriously sensitive. There is evidence that fructose is more rapidly utilized in the body than glucose, presumably because it is more rapidly converted into normal blood sugar. It is remarkable therefore that Hewitt and Pryde attained such little success with the use of fructose.

Hewitt rightly observes: "The dangers attending the

use of permanganate with biological material are obvious."⁵ The technique adopted by these authors when using permanganate has been briefly referred to by us as being not above suspicion.⁶ They omitted to perform the only conclusive test—namely, a comparison between the rates of decolorization at the beginning of the experiment, and at the end when the sugar had attained equilibrium. The italics used in discussing their experimental method hence have not the importance the authors attach to them.

We may conclude that the experimental evidence put forward by Hewitt and Pryde does not support Hewitt's recent contention² that "they showed with some certainty that γ glucose and γ fructose are formed and can be detected when the stable α and β isomerides are in contact with living intact intestine." We are glad to note that the possible significance of their experiments performed three years ago has not escaped Hewitt, and that experiments are now in progress on the diabetic organism. We feel that it is necessary to point out the above facts in order to correct any doubt on our work which Hewitt's criticism may have caused.

Summary.

1. There is definite evidence that a reactive sugar is present in the blood.
2. There is no chemical evidence that γ glucose is formed in the body.
3. There is definite evidence that some of the blood sugar of diabetics is fundamentally different from that of normal persons.
4. No evidence at present exists that the intestine plays an essential part in the causation of this difference.

REFERENCES.

- ¹ Winter and Smith: *Journ. of Physiol.*, 1922, 57, p. 100. ²Hewitt: *BRITISH MEDICAL JOURNAL*, 1923, p. 590. ³Hewitt and Pryde: *Biochem. Journ.*, 1920, 14, p. 395. ⁴Hewitt and De Souza: *Ibid.*, 1921, 15, p. 667. ⁵Forrest, Smith, and Winter: *Journ. of Physiol.*, 1923, 57, p. 224. ⁶Winter and Smith: *Proc. Phys. Soc., Journ. of Physiol.*, 1923, 57, p. xxxi. ⁷Cooper and Walker: *Biochem. Journ.*, 1922, 16, p. 455. ⁸Winter and Smith: *BRITISH MEDICAL JOURNAL*, 1923, i, p. 711.

THE TREATMENT OF GENERAL PARALYSIS BY MALARIA.

BY

W. L. TEMPLETON, M.B., CH.B.,

ASSISTANT MEDICAL OFFICER, CITY OF LONDON MENTAL HOSPITAL, DARTFORD.

A SHORT account of the treatment of general paralysis of the insane by infection with malaria was given by Dr. E. W. Scripture at a meeting of the Medico-Psychological Association on November 23rd, as reported in the *BRITISH MEDICAL JOURNAL* of December 9th, 1922 (p. 1121). The principle depends on the fact that remissions in chronic disease may occur after an attack of an acute specific fever. Many methods of producing fever artificially have been tried, with varying results, but Wagner-Jauregg, quoted by Pilcz,¹ has drawn up a scale of efficiency of the various methods used as follows: (1) chemical substances—for example, sodium nucleinate; (2) toxalbumins—for example, tuberculin, staphylococci; (3) acute disease; the last is the most efficient. With tuberculin treatment Pilcz² obtained a remission in 26 per cent. of his cases, and eight remained well three years after treatment. The treatment by malaria gives a much higher percentage of remissions. During a recent residence in Vienna as a post-graduate I had an opportunity of seeing some of the clinical results, which are very encouraging.

The method consists in the withdrawal of from 2 to 4 c.cm. of blood from the vein of a patient suffering from benign tertian malaria (whether the patient be in the febrile stage or not), and in its subcutaneous injection into the subject of general paralysis, who will develop within about a week typical attacks of malaria. He is allowed to have ten to twelve such attacks, when the malaria is cured by the administration of quinine in doses of 1 gram for three days in succession; then in doses of 0.5 gram for fourteen days. In no case did the malaria persist; it invariably cleared up after the first few doses of quinine. During the febrile attacks and occasionally after, there is very often an exacerbation of mental