to 8 grains a day in November, and on two or three days none at all was taken; a relapse of subtertian malaria occurred on November 26th, which was quickly controlled by increasing the dose.

The patient left for England on December 1st, and was in good health when we last heard of her.

THE PULSE PRESSURE IN EXOPHTHALMIC GOITRE.

BY

I. HARRIS, M.D.,

HONORARY PHYSICIAN IN CHARGE OF CARDIOGRAPHIC DEPARTMENT,
NORTHERN HOSPITAL, LIVERPOOL.

THE height of the pulse pressure (the difference between systolic and diastolic pressure) is usually influenced under normal conditions by two factors—the height of the systolic pressure and the pulse rate. The higher the systolic pressure the more pronounced is the pulse pressure; the more frequent the pulse rate, the smaller the pulse pressure.

The systolic arterial pressure is usually increased in exophthalmic goitre, although in cases where symptoms of heart failure manifest themselves it may become lower than normal; but the most striking feature in exophthalmic goitre is the relation of pulse rate to pulse pressure. As is well known, the pulse rate is increased in exophthalmic goitre. A frequent pulse in any other condition is usually accompanied by a small pulse pressure. In tachycardia due to bacterial toxins acting on the heart muscle and other forms of myocardial trouble, and in cardiac neurasthenia and heart failure, etc., the pulse pressure is small, usually less than 40 mm. Hg. In rapid heart action due to amyl nitrite and atropine the pulse pressure is likewise diminished.

The following table records the various pressures in cases of exophthalmic goitre:

Initial.	Blood Pressure.			Pulse Rate.		Initial.	Blood Pressure.			Pulse Rate.
	S.	D.	P.				S.	D.	P.	
M.	125	80	45	130		S.	120	80	40	120*
E.	140	70	70	140		W.	138	67	71	160
C.	140	80	60	120		S.	165	95	70	†
J.	125	75	50	100		G.	170	85	80	120
C.	115	50	65	60		R.	170	95	75	90
D.	137	80	57	90		S.	140	80	60	120

S. = Systolic pressure; D. = Diastolic pressure; P. = Pulse pressure.

* Pronounced cardiac dilatation.

† Auricular fibrillation.

So far as I am aware it is only in thyroidism that there is both an increase in pulse rate and at the same time a high pulse pressure. In some cases of irritable heart with manifest signs of thyroidism pulse pressure is also increased. This fact is of value in the differential diagnosis between thyroidism and other conditions. It is also important from the point of view of treatment. I have shown elsewhere¹ that digitalis increases the pulse pressure. Obviously this drug is not indicated in a condition where the pulse pressure is already high.

The electro-cardiograph reveals still another characteristic sign of exophthalmic goitre; the P wave is usually exaggerated in this condition, suggesting hypertrophy of the auricle. A similar electro-cardiographic condition is found in mitral stenosis, and it is significant that exophthalmic goitre, too, predisposes to auricular fibrillation.

REFERENCE.

¹ *Lancet*, 19.1.1, 1972.

THE seventh Sicilian Medical Congress will be held at Palermo from April 20th to 22nd, when the following subjects will be discussed: Deficiency diseases, introduced by Dr. Giuffrè of Palermo; the present state of our knowledge on the etiology and treatment of cancer, introduced by Dr. Parla-vecchio of Palermo and Dr. Fichera of Messina; and the organization of the Italian medical profession, introduced by Dr. Salpietro of Palermo.

Memoranda:

MEDICAL, SURGICAL, OBSTETRICAL.

SUPPRESSION OF URINE AFTER LABOUR.

SUPPRESSION of urine after labour is apparently very rare, and is not even mentioned in most obstetric textbooks. I have inquired of several medical practitioners—some of them of thirty-five years' experience—and find that none of them have ever seen a case. The following may therefore be of interest.

Mrs. G., aged 30, had the following obstetric history. She was married in 1916, had a miscarriage in May, 1917, a full-time child in May, 1918, a full-time child in August, 1919, a two-months miscarriage on January 30th, 1922, and twins on February 4th, 1923. I attended her on all occasions except the first miscarriage; in 1919 I attended her continuously for the last two months of her pregnancy, and feared acute phthisis, but, to my surprise, she improved and got apparently quite well after the birth of the child. During her last pregnancy she complained of vague symptoms, most of them nervous, but there were no physical signs, and she appeared in better health than ever before. Repeated requests for a specimen of urine, however, were made in vain (a not uncommon event in general practice), so that I cannot be sure as to the state of the renal system.

On January 20th, 1923, I examined her and diagnosed a twin pregnancy. On February 4th I was sent for by the midwife because "the placenta had not arrived after forty-five minutes," a child having been born. On examination I found that another foetus was present in the uterus; it was presenting transversely, and there was prolapse of an arm. With the help of a colleague who gave chloroform, I performed internal podalic version. Both twins lived; there was little shock after the manipulations, and the haemorrhage, though rather free from a flabby uterus, was not alarming. On the following morning I was told that the patient had had continuous bilious vomiting, and she was still retching. I hoped that this was due only to the anaesthetic, but I noticed that there was extreme oedema of the eyelids, and the patient told me that "everything was a mist before her." There was no headache—which I mistakenly took to be a good sign. The same day, at 6 p.m., I was informed that the patient had passed no urine, and on discovering that there was no distension of the bladder, the seriousness of the situation was apparent. I passed a catheter, but found, as I expected, that not a drop of urine was obtained. Later in the evening, after appropriate treatment, the patient looked and said that she felt better; the oedema of the eyelids had improved, and she did not complain of headache. The following morning, however, when I saw her at 8.45, she was unconscious, breathing stertorously, the pupils were pin-point, and she died an hour later—forty-one hours after labour was completed. There was no headache nor convulsion at any time. No urine had been passed.

The vomiting was uncontrollable by drugs, and none of the usual measures, such as hot blankets and bottles, hot steam, cupping of the kidneys, pilocarpine, etc., had the slightest effect. No post-mortem examination was obtained.

In view of the feared development of phthisis the possibility of a tuberculous focus in the renal system is suggested, though there was nothing to lead me to suppose this during her pregnancy, when, as I have remarked, she appeared in better health than ever before. However, as I did not have the opportunity of examining the urine, I cannot be definite on this point. The urine was normal on all other occasions, on which I had examined it.

AMBROSE W. OWEN, M.D., B.S.Lond.,

Honorary Surgeon, Aberdare and District General Hospital.

TRAUMA AND APPENDICITIS.

THE reports of cases under this heading (January 6th and 13th) induce me to record a similar case which came under my observation at the Marine Lines War Hospital, Bombay, in 1920.

A young clerk was hit by a ball over the right iliac fossa while playing hockey. He was admitted into my ward the next morning suffering from a typical attack of acute appendicitis. He refused operation at first, but submitted to it a week later, when the pain and swelling had increased. At the operation the caecum and the terminal ileum were found matted into a hard lump, with the tip of the appendix projecting about the middle. This last was removed for examination, and a drainage tube was tied in. The patient made a good recovery after some days of suppuration. Microscopic examination showed a sclerosed appendix with the lumen obliterated.

B. P. SABAWALA, F.R.C.S.E.,
Major I.M.S. (Hon.).

Bombay.