trace of albumin. The diagnosis, therefore, should not be difficult, provided the sputum be carefully examined.

REFERENCES.

- ¹ *Traité de l'auscultation médiate et des maladies des poumons et du cœur*, par R. T. H. Laennec, troisième édition, tome i, p. 151, Paris, 1831.
² Albuminous Expectoration following Paracentesis of the Chest, by P. Horton-Smith Hartley. *St. Bartholomew's Hospital Reports*, vol. lxi, 1905, p. 77.

LACERATED KIDNEY DUE TO INDIRECT VIOLENCE.

BY

JOHN C. JEFFERSON, F.R.C.S.,
HONORARY SURGEON, ROCHDALE INFIRMARY.

THERE are several points of interest in the following case, which will be discussed after the facts have been related.

Mrs. C., aged 53, was first seen on January 1st, 1923. She stated that thirty-six hours previously she had fallen from a chair on which she was standing, and immediately afterwards was seized with abdominal pain. She went to bed, and some time during the night vomiting commenced, and continued up to the time of my seeing her on the morning of the second day after the accident. She then appeared very ill. The pulse was 110 and rather weak, the temperature 100°. There were no signs of external violence. The abdomen was distended and acutely tender in its lower segments, particularly over the right iliac fossa. It exhibited an intense rigidity below the level of the umbilicus; above that level rigidity was not so marked. The liver dullness was not diminished; rectal and vaginal examinations yielded no information. The urine was scanty and high-coloured, but contained no blood. A diagnosis of peritonitis was made, though I was unable to determine its cause.

Operation.

She was removed to hospital, where operation was at once performed. On palpation under the anaesthetic a fixed mass was felt in the right iliac fossa, passing backwards into the right lumbar region and upwards beneath the ribs.

An incision over the right rectus was made, and it was at once seen that there was a considerable quantity of free blood and clot in the abdominal cavity. The mass felt on palpation was found to be a haematoma around the right kidney. This haematoma had tracked downwards along the psoas, had stripped up the peritoneum, and lifted forwards the caecum and ileo-caecal valve; further, it had percolated into the lower part of the mesentery of the small intestine.

To the outer side of the ascending colon there was a small tear in the parietal peritoneum not more than one inch long. Through this blood had evidently made its way into the peritoneal cavity. It was interesting to note that the caecum, appendix, and terminal part of the ileum were deeply ecchymosed, and presented the appearance that they would have shown had they been involved in a recent strangulation. The blood and clots were rapidly cleared out, the rent in the peritoneum sutured, and the abdomen closed.

An incision was now made exposing the right kidney. The haematoma was cleared away, and, on gentle examination of the organ *in situ*, a laceration was felt on the anterior surface of the lower pole extending from the lateral border inwards towards, but not involving, the pelvis. As there was now no bleeding from the laceration its depth was not determined, and I contented myself with packing the cavity resulting from the evacuation of the haematoma and closing the wound as far as practicable around the pack.

After-History: Recovery.

The patient's condition at the end of the operation was grave and continued so for two days, there being much post-operative vomiting. From then on, however, she made good progress. The packing was removed in two or three stages and a drainage tube substituted. Through this urine was discharged in small amount for five weeks. There was never any blood in the urine passed naturally. The patient left hospital at the end of the eighth week.

In the first place, the laceration was caused by indirect violence. In such circumstances the injury to the kidney may arise from acute flexion of the body with corresponding acute flexion of the kidney on its transverse axis (Thomson Walker); or where there is no such flexion but where the accident consists of a fall from a height on to the buttocks, the kidney may be injured by direct impact against the twelfth rib or transverse process of the first lumbar vertebra (Tuffier). In the case under consideration I think the laceration was produced by the first mechanism, for the distance from the seat of an ordinary chair to the ground would hardly be sufficient to cause it by the second method.

Secondly, though lacerations of the kidney usually radiate outwards from the hilum, in this instance the laceration was produced in the cortex only, and, though small, had given rise to much haemorrhage.

Thirdly, there was no haematuria so far as I was able to observe. There may, of course, have been blood in the urine when the patient, who was by no means an intelligent type, was not in a condition to notice it. On the other hand, since the laceration was in the cortex only, there may have been none at any time.

Fourthly, although the laceration was caused by indirect violence the peritoneum was torn and there was accompanying haemoperitoneum.

There is no doubt that haemorrhage had already ceased at the time of operation, but it is highly probable that but for the treatment adopted urine would sooner or later have found its way into the peritoneal cavity, with fatal results.

I employed treatment by packing, rather than suture of the laceration, in order to save valuable time. In this connexion it is interesting to note that Albarran has collected six cases in which packing was resorted to; all recovered (Thomson Walker).

Memoranda:

MEDICAL, SURGICAL, OBSTETRICAL.

LACTIC ACID IN THE TREATMENT OF CHRONIC ENTERITIS.

A NUMBER of cases of chronic enteritis are associated with excessive fermentation of the intestinal content. The condition frequently arises *de novo*, although in a certain proportion of cases it follows an acute stage.

It occurs in babies as a result of the food becoming contaminated by micro-organisms. The motions are frequently loose, green, and malodorous. This is usually associated with abdominal distension, excessive flatus, and attacks of screaming due to colic. There is nothing palpable abdominally. In adults the following syndrome is present:

(1) Colic of either the small or large intestine. The pain is variable in intensity, and usually moves from one part to another. In the small intestine it is usually referred to the region of the umbilicus. The large intestine may be definitely palpable and tender to touch. I have seen one case where the caecum was palpable.

(2) The stools vary in consistency and colour. There may be looseness alternating with normally formed motions or constipation. The stools may be paler than usual, or even greenish. They are frequently offensive, and are likely to be associated with excessive flatus.

(3) There may be an associated gastric dyspepsia. Where the condition has lasted some time mental depression, loss of weight, anaemia, and headache may occur as a result of toxic absorption.

Cases of the types referred to respond very well to lactic acid. The rationale of the treatment lies in the fact that *B. coli*, which is as a rule unusually active in those cases where the motions are unduly offensive, cannot grow in a concentration of lactic acid of more than 1 per cent.¹ The treatment is a result of Metchnikoff's teaching, and given in this way is a satisfactory alternative to curdled milk preparations. The latter cannot always be tolerated by patients,² and involves a certain amount of difficulty where there is not the proper convenience for making bacterial preparations.

The following prescription is useful for adults:

R	Ac. lact.	...	...	...	...	...	3ij
	Syr. aurant.	...	...	...	...	...	3j
	Aq.	...	...	...	...	...	ad 3viij

Sig.: 3ss t.d.s. p.c.

Where there is gastric dyspepsia the following may be substituted:

R	Ac. lact.	...	...	...	...	...	3ij
	Liq. strych.	...	...	...	...	...	mxx
	Tr. card. co.	...	...	...	...	...	3j
	Aq.	...	...	...	...	...	ad 3viij

Sig.: 3ss t.d.s. p.c.

I. H. LLOYD-WILLIAMS, M.B., B.S. Lond.,
M.R.C.S., L.R.C.P.

¹ *Archives of the Middlesex Hospital*, 1916.

² Herschell: *BRITISH MEDICAL JOURNAL*, 1910, i, p. 21.