It does not appear to influence the acute pains in the lower limbs.

The importance of nursing in these cases cannot be overestimated. In fact, without the special care that the patients receive from the nursing staff at Gifford House I doubt whether any of the complete cases would be alive to-day.

Prognosis and General Summary.

In the complete cases the prognosis cannot be other than ultimately fatal, but in the absence of definite uraemic symptoms it is almost impossible to predict an immediately fatal issue. Even where a patient may seem semi-moribund, with foul discharging bedsores, intermittent fever, and rapid general wasting, there is quite a chance that he will make a good recovery and be up in his chair again. A patient with extensive osteomyelitis of the spinal vertebrae following a bed sore gradually pulled round and the condition entirely healed. On the other hand, I have not seen recovery following any uraemic nervous symptoms, even when the patient's general condition appeared quite good. It is always safe to predict rapidly developing coma and death when such symptoms exist. Acute haemorrhagic cystitis is always rapidly fatal; it may come on without any warning at any time during the patient's paraplegic life. I have seen two deaths from this condition, both in patients without suprapubic drainage. The mental factor undoubtedly plays a very large part in the grip that the patient has on life. A greatly depressing shock of some kind, the death of a friend, or serious trouble of any sort, will be sufficient to depress the patient's vital powers. He will become silent, depressed, refuse food, look increasingly ill, begin to develop bedsores, and may die if his confidence is not restored.

The main factors influencing prognosis appear to be:

1. Continued and skilled nursing.
2. Daily massage.
3. Establishment of adequate bladder drainage, preferably by the suprapubic method. Closure of a suprapubic cystotomy opening which the patient has had for some years and which is giving no local trouble is entirely contraindicated. This statement has been put to the test in five of the complete cases, and in all five symptoms have developed, such as bedsores, limb pains, fever, and malaise, which were not manifest previous to the closure.
4. Encouragement of mental confidence and mental occupation. In this respect an auto-wheel chair of special pattern has been found invaluable. It saves the patient any unnecessary friction, and enables him to get about and enjoy life, and generally assists him in forgetting the tragic aspects of his disability.

A NOTE ON THE VALUE OF SYMPTOMS IN THE EARLY DIAGNOSIS OF PULMONARY TUBERCULOSIS.

BY

D. G. MACLEOD MUNRO, M.D., M.R.C.P. EDIN.,

DEPUTY COMMISSIONER OF MEDICAL SERVICES FOR TUBERCULOSIS TO THE
MINISTRY OF PENSIONS.

THESE remarks are addressed primarily to the general practitioner, upon whose failure to diagnose pulmonary tuberculosis in its early stage some quite unjustifiable aspersions have been made from time to time. It is, indeed, encouraging to note how keenly the general-practitioner has recently taken up this question and with what comparative success, especially on the lines of more recent views on physical diagnosis. It is not my intention to discuss this question from any original or novel standpoint, but I am inclined to think that we should seriously consider whether in this connexion the pendulum has not swung rather too far in the direction of diagnosis from purely physical signs.

The time is, I think, ripe to seek an adjustment of values between the two chief factors in the early diagnosis of pulmonary tuberculosis—namely, its symptomatology and physical signs. In a disease in which so many clinical and anatomical phenomena have to be correlated, and in which

the same individual may present a new symptom complex every few weeks, the position of the diagnostician is not always a happy one. It is only by painstaking adjustment of all the contributory elements in such cases and in the use of skilled judgement in interpreting the clinical picture thus presented that success will attend efforts to make a diagnosis in these difficult cases.

I am not aware whether it is still the rule, but until very recently I understand the diagnosis of pulmonary tuberculosis in the army was not regarded as established unless the tubercle bacillus had been found in the sputum; and it is not so many years ago that in civil practice the earliest recognition of the disease was based upon data which at the present time are considered to be the sign of a comparatively advanced stage. In auscultation, for example, it may not yet be generally appreciated that what is to be looked for is a change in the breath sounds of the *inspiratory* phase, which may be noted at a much earlier stage of the disease than alteration in the *expiratory* phase which may be present in many quite healthy chests.

The change of method in the elucidation of physical signs during the past few years has been, indeed, quite revolutionary, and thanks to Riviere and others, certain refinements in physical examination have materially assisted in the earlier recognition of the disease. The utility of inspection and palpation has been especially developed by the American school, and the value of these methods of physical diagnosis greatly enhanced. Hitherto inspection of the chest has been, from the point of view of the general practitioner, rather a perfunctory affair, and the virtues of palpation limited to vocal fremitus—now regarded as of doubtful value as a diagnostic sign in such cases. Now that special attention is being paid to the muscle reflex which results in what is known as "lagging" (the presence of which, especially at an apex, strongly suggests active mischief, although percussion and auscultation may give no indication of this) inspection and palpation assume a more important place. The proper appreciation of this sign requires practice, but this is soon acquired.

The information conveyed by inspection in these cases may be greatly reinforced by light finger-tip palpation, by means of which evidence of localized reflex muscle spasm—especially as affecting the sterno-mastoid, the scaleni, and the upper pectorals—may be detected. The trained finger appreciates in such cases a sense of muscle resistance in contradistinction to the smooth and more elastic sensation of normal muscle tension. Then, too, trophic changes in the muscles of the shoulder girdle and the state of the subcutaneous tissues may be similarly recognized. In the opinion of some experts, indeed, light touch palpation has very largely replaced percussion as an aid to diagnosis.

The technique of percussion, too, has been much improved, and it is now generally recognized that to be of any real value, apart from other points of detail, percussion must be light. But this method of physical diagnosis still presents many pitfalls, and, apart altogether from the question of accuracy, it must be strongly affirmed that local changes in the percussion note, especially at the apex of a lung—even if the existence of such cannot be doubted—convey nothing as to the presence of an active tuberculous lesion, or, indeed, that the lesion, if such exists, is tuberculous at all.

In my opinion, too much attention in the matter of percussion has been focused upon the definition of certain areas of the chest wall called after some eminent authorities. It is questionable, for example, whether in the early cases of pulmonary tuberculosis now under discussion any appreciable change can be found to have occurred in Krönig's area of so-called apical resonance. In rather a different category are the "bands of impairment" in the percussion note first noted by Abrams. These bands are said to be purely reflex in origin and to be one of the earliest indications of active pulmonary tuberculosis. My own experience of this diagnostic sign is not extensive, but I confess that in a number of cases in which I have utilized this method of physical diagnosis I have never been able to satisfy myself as to the reliability of the results.

But if we may gratefully acknowledge that a great deal of useful work has been done by what I may perhaps refer to without offence as the "physical force" school, we must admit, I think, that on the question of the early diagnosis

of pulmonary tuberculosis they have of late held the field too exclusively.

Turning, then, to the symptomatology of the disease, we find that a toxæmic state, revealed in one form or another and in greatly varying degree, is to be regarded as a manifestation of active tuberculosis. Pulmonary tuberculosis, however, cannot be held responsible for every unexplained toxæmia, as the symptom complex of this condition is much the same whatever the source of infection, and the existence of localized infection such as sinusitis, tonsillar infections, oral sepsis, and bronchiectasis must be excluded.

The group of symptoms comprise malaise, lassitude, nervous instability, digestive disturbances, tachycardia, fever, and night-sweats. I propose to deal in some detail only with two of the more prominent of the symptoms of the toxæmic syndrome, both of which on accurate observation will be found to present certain features characteristic of tuberculosis. These are (a) fever, (b) tachycardia.

Fever.

The presence of fever as an indication of tuberculous toxæmia is in this connexion of the greatest value, but certain qualifications are necessary if we are to arrive at trustworthy conclusions. It is unfortunate that temperature records are often taken in so casual a manner as to be practically valueless. It is not necessary, I think, to assume that a rectal temperature record is the only reliable method, more especially in private practice. If certain simple precautions are taken oral temperature records are quite trustworthy, except in the case of mouth-breathers. The thermometer should be retained in the mouth at least seven minutes; the temperature should not be taken immediately after the patient returns indoors after exposure to cold wind nor after partaking of a hot meal. It is not sufficiently recognized that the maximum temperature period varies greatly in different persons; to ascertain this period in a given case it is necessary to take the temperature every two hours. The first recorded temperature for the day ought not to be taken, as is often the case, after the patient has got up and dressed, but as early as possible in the morning before rising. The normal diurnal variation in temperature is about 1.5° F. In a majority of healthy persons the average early morning temperature is approximately 97.2° to 97.4° F., rising to a maximum of 98.6° F. in the early afternoon. This normal average is maintained by an equilibrium between the forces governing heat production and heat dissipation, and the entry of toxins into the blood stream, from whatever source, disturbs this balance. While it is not usual to find much temperature elevation in early pulmonary tuberculosis, it may be found that the early morning temperature is 98° F. and the maximum record for the day 99.2° or 99.4° F. The necessity for a continuous temperature record extending over three or four weeks in any case of suspected early tuberculosis will become apparent when it is pointed out that the toxæmic state is by no means constant; it ebbs and flows. The continuous temperature record therefore gives a fair representation of the degree of toxæmia present.

In chlorosis there is sometimes a slight average increase of temperature, while the premenstrual rise must not be overlooked. In persons of highly strung nervous temperament there are sometimes quite noticeable variations in the temperature curve. It will thus be seen that the process of elimination may have to be thoroughly invoked to clear the ground. Are there, then, any very clear indications, when all this has been done, which may point the way to a tuberculous toxæmia as the responsible source of the fever? I think there are. My own experience has led me to attach much importance to an early morning temperature—taken before rising—which is consistently a little above the average mean. The effect of a definite amount of exercise may be noted; while in a healthy person there is always a temporary rise—sometimes considerable—after exercise, this is usually quite evanescent, and within half an hour or so the temperature returns to normal. This is not so in the case of a tuberculous person. The decline of the temperature to normal limits is delayed and it may remain elevated for two or three hours. Hence the value of post-exercise temperature records in suspicious cases. Then, too, extraneous conditions resulting in emotional disturbance, worry, and anxiety,

which in a healthy person have no appreciable effect on temperature, may cause considerable disturbance in a tuberculous person. These features may be said to be characteristic of the tuberculous process. It will be seen, therefore, that fever, though still to be regarded as one of the cardinal points of the toxæmia of tuberculosis, cannot be accepted without due consideration of the qualifications outlined above, especially if there is no supporting evidence in other directions. It must be noted, however, that the absence of fever does not necessarily rule out the possibility of the existence of pulmonary tuberculosis.

Tachycardia.

From time to time I have seen patients found to be suffering from early pulmonary tuberculosis who have referred to an irregular and rapid action of the heart coming on without any apparent reason, but on the whole it has not been my experience that the cardiac disturbances caused by tuberculous toxæmia are noticed subjectively by the patient.

Tachycardia is of frequent occurrence in early tuberculosis—Fishberg puts it at 60 per cent. of incipient cases; it is hardly ever a subjective symptom. Like fever it is of toxic origin and appears to be the result of excessive stimulation of the sympathetic nervous system. The instability of the pulse in pulmonary tuberculosis would appear to result from a combination of vagus and sympathetic stimulation. Pottenger, who supports this view, considers that tachycardia as such is not a particular characteristic of active tuberculosis. In his opinion it is not regularly present unless the patient is in a depressed state, or has been over-exerting himself. This, I am inclined to think, does not fully represent the case. It is now, I believe, generally accepted that a pulse rate out of all proportion to any febrile condition present is a very common and reliable symptom of early tuberculosis in an active state, and further that the pulse rate may be notably increased on the slightest excitement or even after a fit of coughing. Another well marked characteristic is that, just as in the case of fever due to tuberculous toxæmia the fall of temperature is considerably delayed after exercise, so it is found that the increased pulse rate after exercise falls much more slowly than in health. I have repeatedly seen cases in which there has been persistent tachycardia, not referable to any cardiac abnormality or hyperthyroidism—a condition which is always to be borne in mind in the differential diagnosis—and in which the true diagnosis was only revealed by a sharp hæmoptysis. The physical signs in the chest in such cases may for a long time show no evidence of anything abnormal.

It has to be recognized, however, that pulmonary tuberculosis may masquerade as an affection of any of the great systems of the body with no signs or symptoms in any way directly referable to the lungs.

In none of the types of cases under discussion is it assumed that assistance can be sought by the presence of cough, sputum, hæmoptysis, pleurisy, or indeed any direct overt symptom. In such early cases radiography is of doubtful value and the complement fixation test is at times rather a blind guide. The key to the diagnosis is to be found more commonly in the history of the present illness, and inquiry cannot be too minute or painstaking.

OBSERVATIONS ON A CASE OF MYASTHENIA GRAVIS.

BY

S. L. HEALD, M.D., F.R.C.S. EDIN.,

HONORARY PHYSICIAN, COVENTRY AND WARWICKSHIRE HOSPITAL;

AND

ARTHUR J. WILSON, M.B., CH.B. EDIN.,

HOUSE-PHYSICIAN, COVENTRY AND WARWICKSHIRE HOSPITAL.

THE following case, of a man in whom symptoms of this rare condition developed while he was serving in the army in the late war, and some six years before he attended this hospital as an out-patient, seems of sufficient interest to warrant publication.

The patient, a man 43 years of age, a painter by trade, first came under observation as an out-patient on December 13th, 1922, when he was complaining of muscular weakness and exhaustion and inability to swallow properly.