of the maxillary sinuses prevented stagnation in these cavities and its consequent back-pressure and reinfection of the deeper sinuses. The frontal sinuses drained spontaneously and were the first to clear. The drainage from the sphenoidal sinuses was encouraged by daily, or twice daily, introduction of strips of gauze (between the middle turbinal and septum) soaked in equal parts of 5 per cent. cocaine and adrenaline. I was very glad that I had refrained from any operative treatment on these deeper cavities. The traumatism entailed might only—in such an acute infection, and in the narrow and sensitive nose—have resulted in further blockage to the outflow and might possibly have spread infection to other areas through the blood or lymph streams. Pain, insomnia, fever, and other symptoms were treated on general principles. Local warmth (by woollen nightcap and radiant heat) gave relief.

I have never seen a fatal case resulting directly from acute sinusitis, diagnosed in good time, and treated on the lines described in this paper. In the two fatal cases of sphenoidal sinusitis which I have referred to, secondary infection of the meninges or cavernous sinus had developed before the patients came under observation.

The care taken of the present case was successful in fending off all complications, except the curious acute thyroiditis. This, I imagine, was an infection via the lymphatics, from the pharynx or tonsil, due to the pus descending from the sphenoidal sinuses.

The moral I venture to draw from this case is that all cases of acute sinusitis should be regarded with much more consideration than they generally receive. If carefully treated there are very few which would not recover completely. Most of the chronic cases seen in practice and requiring operative treatment commenced originally as acute cases. Had they then been treated carefully, they would not have drifted into the chronic infections which entail so much ill health and often demand prolonged attention and sometimes anxious operative measures.

The treatment of acute sinusitis is largely in the hands of the family physician. Except for exploratory puncture and lavage of the maxillary sinuses, operative treatment is rarely called for in an acute case.

REFERENCE.

¹ Transactions of the Medical Society of London, vol. xxix, 1906.

OPTIC NEURITIS OF SPHENOIDAL SINUS ORIGIN: OPERATION: CURE.

(With Special Plate.)

BY

SIR STCLAIR THOMSON.

THERE are various causes of retrobulbar neuritis. I am able to give a very full report of a case in which the onset of this affection was rapid and the cause was undoubtedly suppuration in the sphenoidal sinuses.

Previous History.—A young man of 24 reported that, as a boy, he had worn glasses, but that on joining the army, during the war, he was told that glasses were not necessary. He served for three years in the army; chiefly on the Italian front, and during his service he developed a yellow-green post-nasal catarrh with occasional cacocsmia. After demobilization in 1919 he noticed that he had more need of handkerchiefs and had to

the left sphenoidal sinus, owing to the prominence of the middle turbinal. There were numerous carious teeth, and the patient was at once sent to Mr. W. Hern for attention to them.

Ophthalmic Report.—On the same day he was examined by Mr. Treacher Collins, who reported as follows:

"Has had aching pain in left eye for three days, which then ceased. Eyeball still tender to pressure, but not so much so as at first. Experiences sharp pain on movement of his left eye upwards. Vision of right 6/9 partly; -1 cyl. 120°=6/6; 1 letter. Left 6/60 unimproved. Left pupil acts to light but does not maintain contraction 'hippus.' Ophthalmic examination shows marked swelling of the left optic disc, blurring of its margins, and obscuration of retinal vessels. The swelling of the disc = 6 D. Right optic disc normal. Field of vision in left eye somewhat contracted up and in (vide Chart 1)."

Treatment.

The same evening the patient entered a nursing home, where he was given radiant heat baths to his head, for twenty minutes three times a day, with steam inhalations of menthol and eucalyptus every two hours, and a few doses of aspirin. The

LEFT I.

CHART 1.—Optic neuritis of sphenoidal sinus origin. November 28th, 1922.

wash his nose out. When fatigued he was subject to headache over the left eye.

Sudden Onset.—On November 20th, 1922, while on business in Germany, he felt depressed all day, and in the evening was surprised to find that he could not see anything clearly from the left eye. He consulted Professor Römer of Bonn on November 23rd, who found that there was inflammation of the optic nerve of the left eye, negative Wassermann reaction, and that an x-ray photograph showed some opacity of the accessory sinuses. A written report advised that the trouble was serious and threatened the eyesight, and that it was a question for a rhinologist whether the condition would abate with conservative treatment or require operation on the sinuses.

Nasal Condition.—On November 28th I found pus on the floor of each nostril and the roof of each choana. Exploratory lavage of the maxillary sinuses showed them to be clear. X-ray examination by Mr. Coldwell revealed marked obscurity of the frontal and sphenoidal sinuses. It was impossible to catheterize

LEFT II.

CHART 2.—Optic neuritis of sphenoidal sinus origin. February 17th, 1923.

left eye was kept covered so as to exclude all light. This treatment was continued for three days, but, as there was no improvement in his eyesight, I proceeded to operate on the sphenoidal sinuses on December 2nd.

Operation.—The area of the nose was prepared half an hour beforehand with strips of ribbon gauze soaked in equal parts of 5 per cent. cocaine and adrenaline, and tucked in between the middle turbinal and the septum. This rendered the nose so ischaemic that, when placed under a general anaesthetic, it was not difficult to introduce Killian's long duck-billed speculum between the septum and the middle turbinal, so as to snap and prise outward the latter body. This proceeding avoided the removal of a healthy and useful middle turbinal, with the traumatism attendant on its amputation. The ostium of each sphenoidal sinus was then well opened with punch forceps.

Immediate Results.—Four days were allowed to elapse for the nose to recover from the inevitable reaction. During this time there was no change in the eyesight, although the treatment

by radiant heat, steam inhalations, and rest was continued. On December 6th both sphenoidal sinuses were washed out with warm normal saline, and the patient at once noticed a remarkable improvement in his vision. Two days afterwards Mr. Treacher Collins reported:

"Vision of left eye = 6/18. Still much swelling of the optic disc. White lines in the retina to the outer side of the optic disc."

Progress.—Daily lavage of both sinuses was continued, clotted yellow pus being expelled. On December 20th the radiant heat baths were given up; the patient was sent out into the fresh air, and the washing of the sinuses was continued every two or three days. The discharge was examined bacteriologically and reported to contain—

"a large number of Gram-negative bacilli with a small number of cocci and diphtheroid bacilli. On culture these proved to be the pneumobacillus of Friedländer, *Staphylococcus aureus*, and a diphtheroid bacillus of the Hoffmann type."

Progress was continuous. By January 2nd Mr. Treacher Collins reported:

"Vision of left eye = 6/9 partly. Swelling of optic disc = 1.5 D. Two white dots now only seen in retina."

By January 16th he thought the swelling of the disc was less, and no retinal changes.

Cure.—By February 17th his report was as follows:

"Vision of his left eye is now better than that of his right. With -0.5 cyl. 90° he sees part of 6/6 and reads J.I. fluently. His field of vision shows only very slight contraction up and in (vide Chart 2). The swelling of the optic disc has subsided, but it still presents rather ill-defined margins, and has a filled-in appearance, no sign of physiological cup showing. The retina is quite clear."

The sphenoidal cavities had been washed out at decreasing intervals, and their interior had occasionally been painted with 25 per cent. argyrol. All suppuration in them has now ceased, the greenish-yellow post-nasal discharge has disappeared, and, as a result, the patient has gained in weight and feels better than he has done since the war.

In 1906 I published two cases in which I was able to demonstrate by *post-mortem* examination that ophthalmic complications (hazy discs and diplopia in one, with optic neuritis and complete ophthalmoplegia in the other) were due to suppuration in the sphenoidal sinuses.¹ This led me to a study of the literature of the subject, and I was surprised to find no fewer than forty other cases in which it had been shown by *post-mortem* examination that sphenoidal sinusitis had been the cause of cerebral death. Many of these cases had been accompanied by orbital or ophthalmic symptoms, and changes in the fundus had been noted in half of them. Fortified by this research I ventured to urge that "in all cases of apparently inexplicable retinitis, papillitis, and optic atrophy the nose requires exploration."

In the eighteen years which have since elapsed I have had a good many cases referred to me. In the majority, I must confess, I found no symptoms of suppuration in the sphenoidal or other accessory sinuses to explain fundus changes. On the other hand, in cases of affection of the orbit (so-called "orbital phlegmon," unilateral proptosis, etc.) the *causa causans* could, in the large majority of cases, be found in the nose or its accessory sinuses.² This led to my forming the opinion that while the nose was frequently the source of orbital affections, yet retinitis, papillitis, and optic atrophy were only rarely caused by sepsis in the accessory sinuses. In this opinion I was confirmed by certain cases of retrobulbar neuritis in which I opened the sphenoidal sinuses and found them free from suppuration. The observation that, in many of these cases, the neuritis disappeared only made me think that the cause of it must have been elsewhere. I did not think of claiming the cure as a consequence of opening a healthy sinus.

But, lately, other views have been put forward. It is now held that "the mere presence of pus in the sinuses may undoubtedly be the cause in some instances, but is by no means the usual or only one"; that "in only a small percentage of cases has pus been found"; and that "deflected septa with crowding of the chronically enlarged middle turbinate are the most usual findings."³ We are even assured that "poor ventilation and faulty drainage are all important predisposing factors," and that "it is futile to longer believe purulent sinus disease the one and only etiological factor."⁴ Quite recently nine cases of retrobulbar neuritis were treated by sphenoidal drainage.

In only one case (sixteen months' duration) no improvement had resulted. Yet in only two out of these nine cases was pus found.⁵

These and similar records would seem to warrant the exploratory opening of the sphenoidal sinuses in cases of retrobulbar neuritis, even in the absence of evidence of suppuration in these cavities or the neighbouring ethmoidal cells—when no other cause for the retinitis can be found, and the patient is threatened with blindness.

To suggest the opening of healthy accessory sinuses is a doctrine not altogether free from danger. I have been associated with cases of retrobulbar neuritis which recovered without operation on an accessory sinus and without any definite diagnosis of the cause of the condition. I would like to learn more of the disease from ophthalmologists and to hear from them what is the usual prospect of recovery. It seems a subject very suitable for a joint discussion from this point of view.

In any case, there can be no question as to our duty in opening and draining suppurating sphenoidal sinuses when associated with optic atrophy. In the case reported above operation was strongly recommended by two distinguished ophthalmic surgeons, and it had a double benefit—it averted blindness, and cured the patient of a chronic suppuration which had been poisoning him for some years.

REFERENCES.

- ¹ Cerebral and Ophthalmic Complications in Sphenoidal Sinusitis, *Trans. Med. Soc. London*, xxix, 1906, and *BRITISH MEDICAL JOURNAL*, September 29th, 1906, p. 768.
- ² St. Clair Thomson: The Frequency of Orbital Manifestations of Nasal Sinusitis, *The Ophthalmoscope*, April, 1908.
- ³ Leon E. White: Aeration of the Posterior Accessory Sinuses in Acute Optic Neuritis, *The Laryngoscope*, xxxii, May, 1922, p. 382.
- ⁴ *Idem.*, *ibid.*, June, No. 6, p. 438.
- ⁵ Gavin Young: *Journ. of Laryngology*, December, 1922, and April, 1923.

CYST OF THE RIGHT SUPRARENAL CAPSULE REMOVED BY OPERATION.

BY

SIR HAMILTON A. BALLANCE, K.B.E., C.B.,
M.S., F.R.C.S.,

SURGEON, NORFOLK AND NORWICH HOSPITAL.

(With Special Plate.)

AN operation for the removal of a cyst of the suprarenal capsule is a rare procedure, and a cyst of this organ large enough to produce pressure symptoms is very uncommon; the following case, therefore, seems worth recording.

In August, 1921, I saw, with Dr. Wortley Quait of Mundesley and Dr. Burton-Fanning of Norwich, a lady, aged 49, who for many years had suffered from "indigestion." The trouble had been much worse for the last five years and was mainly characterized by pain in the back passing round the waist to the epigastrium. The pain had no relation to meals, and would often come on at night, when a little food would bring relief. Of late the diet had chiefly consisted of fish and milk foods, and there had been much flatulence and some loss of weight. The patient had been backwards and forwards between this country and the tropics for twenty years, and often when abroad she would be quite free from indigestion troubles for six months at a time. For the last six weeks there had been frequent vomiting, but a week might pass without any sickness, and after such an interval the patient would bring up a considerable quantity. There had never been haematemesis. Constipation had been very troublesome. The stomach was seen to be distended and visible peristalsis was present; there was no tenderness anywhere and no tumour could be felt. An x-ray examination showed that even after eight and a half hours a bi-smuth feed had not entirely left the stomach.

First Operation.

The abdomen was opened and an old ulcer of the first part of the duodenum, just beyond the pylorus, almost encircling the bowel and narrowing it, was found. The stomach was dilated and hypertrophied. A posterior gastro-jejunostomy was performed and the appendix removed. The gall bladder was healthy.

During the examination of the abdominal contents a large cystic tumour was found in the right hypochondrium occupying the concavity of the diaphragm and pushing the liver to the left and downwards, so that the gall bladder was in the middle line of the body and the edge of the right lobe of the liver on the level of the umbilicus. The kidney was displaced downwards, but the cyst was only slightly attached to the upper pole of this organ. At the conclusion of the gastro-jejunostomy it was deemed advisable to defer the removal of the cyst, and the abdominal wound was therefore closed. The patient made a satisfactory recovery from the operation, and examination of the right side of the abdomen a few weeks later did not reveal any abnormal swelling

SIR STCLAIR THOMSON: OPTIC NEURITIS OF SPHENOIDAL SINUS ORIGIN.

Skiagram showing condition of sphenoidal sinus before operation. Inset: Sketch showing the relation of the parts. T, Sella turcica. S, Sphenoidal sinus. E, Perpendicular plate of the ethmoid.