Case 13.

A man, aged 69, had noticed dryness of the throat for four months; the teeth were very foul and he smoked about 3½ ounces of tobacco a week. An epithelioma was found involving the left tonsil and extending to the anterior pillar and soft palate; the tongue showed leucoplakia. The growth was diathermized, an enlarged gland in the neck was removed for examination, and the external carotid was tied; the gland was found to be inflammatory. There was no recurrence one year and eight months after operation.

Case 14.

A very old gentleman, aged 75, was found to have a large epithelioma of the right tonsil, extending on to the mucous membrane covering the lower jaw. The patient refused to have the glands removed, but he consented to preliminary ligature of the external carotid, which was carried out, together with a thorough diathermy of the primary growth. The patient made a speedy recovery, and there was no recurrence one year and seven months after operation.

Case 15.

A woman, aged 71, had an epithelioma of the right side of the tongue and adjoining portion of the floor of the mouth, adherent to the lower jaw. There were enlarged glands in the neck. The growth was thoroughly removed by diathermy, and subsequently the glands were removed from both anterior triangles. There was

no recurrence one year and four months after the operation. It will be noted that this was a case of carcinoma of the tongue occurring in a woman.

Case 16.

A man, aged 38, was found to have a carcinoma occupying the whole of the right antrum. The bone was so extensively involved that it was found impossible to remove the tumour completely, and the growth was therefore scraped away with a sharp spoon, and the whole of the cavity subjected to diathermy. A fistula subsequently resulted, and it is hoped at a later date to close the fistula by a plastic operation. There was no recurrence over two years after diathermy.

Case 17.

A man, aged 64, complained of difficulty in swallowing for a month, and of pain in the region of the back of the tongue. He smoked 1½ ounces of tobacco a week, was a heavy drinker at one time, but had been temperate for the last ten years; he was in the habit of drinking very hot tea; his teeth were very septic. A carcinoma of the uvula and soft palate was found, involving the upper extremity of the anterior faucial pillars on both sides. The growth was freely destroyed by diathermy, and although no enlarged glands were felt, the anterior and posterior triangles on both sides were cleared out by "block" dissection. There was no recurrence two years after operation.

REPORT ON A CASE OF DIABETES TREATED WITH INSULIN.

BY
E. I. SPRIGGS, M.D., F.R.C.P., D. V. PICKERING, M.B.,
AND

A. J. LEIGH, B.Sc., A.I.C.

(From the Duff House Laboratories, Ruthin Castle, North Wales.)

THE following observations were made in co-ordination with the scheme of work arranged by the Medical Research Council at various hospital centres.

A man aged 47 was selected as a moderately severe case of diabetes mellitus, who had been observed with some detail over a period of eight years. He was an officer in a Highland regiment and noticed thirst and polyuria a month after the outbreak of war; but he could, at that time, march twenty-five miles. Four months later (January, 1915) he went to Belgium, was speedily exhausted, sent home in February, and came to Duff House, Banff, in March. Since then he has been readmitted five times. He is accurate and conscientious, and has, in the intervals, followed the régime prescribed by his medical advisers here and at home.

In the eight years this patient has been observed under each of the three methods of treatment which have been in vogue in the present generation—namely, (1) restriction of carbohydrate, without restriction of other foodstuffs; (2) the fasting treatment, with subsequent restriction of all foods; and (3) insulin.

Table I gives the weight, state of the urine, the blood sugar, and the calories and carbohydrates in the food at the beginning and the end of each of his former visits to Duff House.

The ascertained maximum tolerance, without glycosuria, for calories and carbohydrate at each visit is shown in Table II.

TABLE II.—Showing the Maximum Tolerance for Carbohydrate and Calories on the same Five Occasions.

Year.	1915.	1918.	1919.	1921.	1922.
Calories ...	3,190	3,116	2,334	2,304	£35
Carbohydrate	56	61	166	127	25
Glycosuria ...	0	0	0	0	0

The table shows that the complaint was at first of a mild type and responded well to treatment; but the deterioration of tolerance is clearly seen. In the last two years especially the patient has found increasing difficulty in keeping anything like sugar-free. His present occupation is not exacting, entailing mainly the duties which fall to the lot of a country gentleman of moderate means and quiet tastes who looks after his garden and animals. But he had been less able to do work and had become progressively more miserable and depressed of late.

The insulin used at first for this patient was prepared here by one of us (A. J. Leigh), who had the advantage of seeing the method carried through at the National Medical Research Council laboratories at Hampstead, and would like to acknowledge his indebtedness to Drs. Dale and Dudley. The mode of preparation was that of Collip, modified by Dale and Dudley. In one point we varied the procedure; this was that we got the use of a small building at the slaughterhouse and minced the fresh pancreas directly it was taken out of the animal, adding the alcohol on the spot and transferring the mixture to the laboratory afterwards. On May 9th and

TABLE I.—Showing the Condition on Admission and on Leaving on Five Previous Occasions.

Date.	Weight in Kilos.	Glycosuria. Grams in 24 hours.	Ketonuria as Acetone. Grams in 24 hours.	Blood Sugar Percentage.	Diet.		Treatment.	Remarks.
					Calories in 24 hours.	Carbohydr. Grams in 24 hours.		
1915. March 24 ...	72½	15	Trace	0.17	2,800	25	Strict diet till glycosuria remained steady; then carbohydrate tests	
May 19 ...	74	4	Small amount	0.16 (May 17th)	3,000	80		
1918. March 16 ...	67	0	0	0.15	3,000	25	Returned for a week for observation	With steady restriction of carbohydrate urine had become sugar-free.
March 21 ..	67	0	0	0.12 (Mar. 2nd)	3,064	41		
1919. July 4 ...	74½	Trace	0	0.12 (July 6th)	2,904	73	Returned for further tests. Carbohydrate gradually increased	Tolerance had improved still further.
August 19 ...	70½	0	0	0.17	2,197	138		
1921. May 2 ...	62½	12	Trace	0.13	1,535	27	Fasting and limitation of total food	In interval fasting treatment had been started owing to much glycosuria and feeling out of sorts. Tolerance reduced, but good.
June 23 ...	62½	0	0	0.16 (June 24th)	2,393	105		
1922. March 21 ...	63	74	Trace	0.14	1,947	11	Limitation of diet (less drastic towards end)	More ground had been lost. Patient insisted on increase of diet beyond his tolerance, contrary to advice.
April 14 ...	63	86	0	0.26	2,556	63		

10th the insulin used was that supplied from the British Drug Houses. We tested this material on rabbits for comparison with our own insulin.

The details of the experimental period of administration of insulin are shown in the accompanying charts.

Experimental Period.

The patient had been upon a fairly constant diet. A similar diet was continued for these observations, and

day, but allowance must be made for the fact that the general level of the blood sugar was lower. Such a flooding of the blood with sugar immediately after carbohydrate food is a normal phenomenon which is exaggerated in diabetes. It is hardly to be expected that insulin, at all events in a small dose, would affect it. There was, of course, also a rise of blood sugar after meals of protein and fat, but it was less high and much slower. In view of these facts it

CHART A.

CHART B.

for the first eleven days (April 19th to 29th) it was fixed both in quantity and quality, except that carbohydrate was added experimentally on two days. The diet contained carbohydrate 17, protein 133, fat 98; calories 1,430. The carbohydrate addition consisted of 60 grams of white bread on the 20th and 23rd. The blood sugar was estimated hourly from 8 a.m. to 9 p.m. for the first five days—that is, fourteen times a day. Later it was taken three times a

day, but allowance must be made for the fact that the general level of the blood sugar was lower. Such a flooding of the blood with sugar immediately after carbohydrate food is a normal phenomenon which is exaggerated in diabetes. It is hardly to be expected that insulin, at all events in a small dose, would affect it. There was, of course, also a rise of blood sugar after meals of protein and fat, but it was less high and much slower. In view of these facts it

appeared wise to make additions to the diet in the form of protein and fat and not of carbohydrate.

3. The sugar in the urine (Chart C) fell from round about 50 grams a day to zero, reappearing when insulin was discontinued and falling again when it was resumed.

4. The acid bodies, expressed as acetone (estimated by van Slyke's method), totalled about 4 grams a day. With 12 units of insulin daily this was reduced to less than 1 gram.

In the records of the case the following points are to be noted.

1. A reduction of the general level of the sugar in the blood after insulin was given, noticeable with 9 units (3 rabbit doses) in three hypodermic doses a day (see Chart A), considerable with 18 units in three doses, and later with 12 units in two doses a day. The effect was greatest four hours after a dose. With repeated meals the sugar was rising high again eight hours after a dose, though not so high as it had been on the same diet without insulin. The effect of the insulin was also definite the following morning, as the blood sugar, on a constant diet, was not so high when insulin had been given the day before. (Chart C; compare 8 a.m. blood sugar on 28th and 29th with that on the 19th, 20th, and 27th.)

2. On the addition of 30 grams of carbohydrate (60 grams bread) to the mid-day meal (April 23rd) immediately after the mid-day dose of 6 units of insulin (Chart B) the percentage of blood sugar rose sharply and remained high for three hours. The rise was of the same character as that obtained by giving the same dose of carbohydrate on a non-insulin day; it was, indeed, though not absolutely so high, relatively more pronounced on the insulin

blood sugar and of sugar in the urine to about the same level as had been reached without insulin on a diet of 1,400 calories. There was also an increase, though a smaller one, in the total acid bodies. The nitrogen balance continued positive up to as much as +5 grams a day. After the above-mentioned excess of food was given, in a further fortnight of observation, not shown on the chart, the insulin was increased and the total food was reduced to about 2,500 calories.

8. The patient's weight increased steadily; 8 lb. were put on in six weeks of treatment.

9. The blood sugar was never brought below 0.11 per cent., and no symptoms of hypoglycaemia were observed.

10. The respiratory quotient was determined several times, but in this case similar figures were obtained with and without insulin. The reflection occurs that the case might not be severe enough or the dosage of insulin large enough to produce considerable changes in the quotient; especially as no attempt was made to enable the patient to assimilate much exogenous carbohydrate.

11. The investigation showed that a suitable régime for this patient is a diet of about 2,500 calories with low carbohydrate, and 30 units of insulin daily.

12. The change in appearance and attitude to life after insulin was given was unmistakable, and was remarked by the members of the staff, who, of course, know the patient well. He grumbled sometimes, as anyone might, at having to have injections and still having to look after his diet, but was interested in his various pursuits and had a look of health and keenness.

It was advised that 20 units of insulin be given just before breakfast and 10 before the evening meal, the smaller dose being given in the evening to avoid reducing the blood sugar too much in the night. This patient when at home sometimes prefers to omit lunch and make two main meals a day, with a little tea in the afternoon. In that case it was advised that the insulin should be given in two equal doses just before the two meals.

The fasting days formerly practised were omitted, but he was advised as a safeguard to take half quantities of food on one day a week. Progress will be watched, and an occasional fast day prescribed if needed.

The following is an example of a day's diet of about 2,500 calories as taken by this patient.

Diet taken on May 26th, 1923.

	Grams.					Cal.	Grams.		
	B.	L.	T.	D.	Total.		Carb.	Prot.	Fat.
*Duff House casein bread	30	15	30	15	90	196		47	
Milk	105		105		210	156	11	7	9
Butter	20	10	20	10	60	476		1	51
Cheese		30		30	60	270		17	22
Eggs	2	1		2	5	400		34	29
Fish—lemon sole ...				120	120	107		25	1
Bacon	60				60	320		10	30
Meat—lean beef—stewed		50			50	293		36	16
Filletted steak ...				60	60				
Vegetables—									
Cabbage	280				280	58	8	5	1
Tomato (raw) ...	150				150	101	16	5	2
Lettuce (raw) ...			300		300				
Kale		400			400	76	4	14	
Asparagus				400	400	92	10	12	
K. iller's sugarless marmalade or jam	30		30		60	9	2	1	
Tea	340		340						
Clear meat soup ...				200					
						2,554	51	214	161

* *Lancet*, 1921, ii, 960; *BRITISH MEDICAL JOURNAL*, 1921, ii, 707; also *Duff House Papers*, vol. i, Oxford Medical Press, 1923.

It will be noticed that in most cases the weights are given in multiples of 30 grams or cubic centimetres, so that the corresponding number of ounces can be approximately calculated by dividing by 30.

Summary.

In a detailed study of a moderately severe case of diabetes in which tolerance had been diminishing for the last three years, it was found that with the help of two daily hypodermic injections of insulin, on a measured diet, the excess of sugar in the blood, the sugar in the urine, and the acidosis could be controlled; also that the patient could take more than double the amount of food he had previously been able to take without the appearance of sugar in the urine. Regulation of diet is still needed, for if an excess of food was taken above the new and higher level of tolerance, the hyperglycaemia, glycosuria, and acidosis reappeared.

The treatment led to improvement in the nutrition, physical capacity, mental outlook, and appearance of the patient.

THE BEHAVIOUR OF THE BLOOD SUGAR UNDER THE ACTION OF INSULIN AND OTHER AGENTS.

BY

G. S. EADIE, M.B.

(From the Departments of Physiology and Biochemistry, University of Toronto.)

IN view of the failure to demonstrate any action of insulin on the oxidation of glucose *in vitro* either in presence of defibrinated blood, muscle juice, a combination of these, or pus,¹ it seems likely that its action is indirect. Winter and Smith² have recently published some experiments which led them to the conclusion that there is present in normal blood a kind of sugar differing from an equilibrated mixture of the α , β varieties in having a lower specific rotation. They fail to find any evidence of this in the blood of diabetic patients, and consider that the diminished carbohydrate metabolism of the diabetic is due to the lack of this more reactive form. It became of interest, therefore, to see if their results could be repeated and to study the effects of epinephrine and insulin. The following is a preliminary account of some of these experiments. No interpretation will be attempted.

Method.

Both dogs and rabbits were used. The rabbits were killed by knocking on the head and bled from the carotid artery; two were used in each experiment. In the case of dogs, blood was removed, either from the heart or femoral artery. The blood so obtained (usually 100 c.cm.) was treated as described by Winter and Smith—namely, diluted with seven volumes of water and precipitated with one volume each of 10 per cent. sodium tungstate and 2/3 N. sulphuric acid. It was filtered first through coarse filter paper, then through a Whatman No. 40 or 42. It was then evaporated to dryness under reduced pressure (water pump), using a capillary to prevent bumping, the temperature of the water-bath surrounding the boiling fluid being kept between 30° and 40° C.; 100 c.cm. of 85 per cent. alcohol was then added (proportionately less when less blood was taken) and the mixture let stand for twenty minutes. It was then filtered into a smaller distillation flask, and the extraction repeated with half the volume of alcohol. This was filtered in the same way, and both filtrates were united and evaporated to dryness as before. The residue was then taken up in distilled water, filtered through a No. 50 Whatman paper, and placed in a polarimeter tube. Readings were made on successive days.

The polarimeter used was a half-shade instrument made by Schmidt and Haensch. Successive readings did not vary more than 0.04°. The average of ten readings was taken. Sodium light was used. The tubes were 1.894, 2, and 2.2 cm. in length.

The copper-reducing value was obtained by the method of Shaffer and Hartmann, using the fluid in the polarimeter tube after the completion of the experiment. Blood sugar estimations were made by the same method. The epinephrine and insulin were administered subcutaneously.

Results.

A general table of the results is given. No experiments are omitted except those in which the polarimeter reading was impossible on account of the opacity of the solution.

(a) *Normal*.—With the rabbit three experiments were done. In one of these the rotation gradually rose to a value corresponding to the reducing power. In the others there was little or no change, but the rotation found was markedly lower than that calculated from the reduction. With the