III.

There are many subtleties of a biological nature which I would have liked to touch upon, such as climate and environment generally, and their influence upon the physical make-up of individuals and races. But I must confine myself to one or two questions, fraught with difficulties and subtleties, which are forcing themselves upon our attention with ever-increasing insistence.

Of psychology we have heard a great deal during the last ten years. Much of what we have heard has been very illuminating and very helpful, but a great deal of it appears to some of us to be the purest balderdash. In an honest endeavour to sift the wheat from the chaff I have been struck with the responsibilities which have been fastened upon the unconscious or subconscious mind. At first I was inclined to think that the subconscious was a pure assumption, a convenient but rather desperate endeavour to explain the otherwise inexplicable. I have, however, been able to convince myself that the unconscious has a real existence, and that it is therefore legitimate to speculate upon its workings. And not only has it a real existence, but a definite and anatomical local habitation. To appreciate this fact we must once more return to first principles.

On looking back through the ages it is not difficult to realize that mind must have existed in the scale of evolution not only before man was moulded, but even before the vertebrates emerged. The ganglion cells which constituted the nervous system of the invertebrates were the ancestors of our autonomous or sympathetic nervous system. These ganglion cells were the organs of a mind, a very primitive and rudimentary mind no doubt, but quite adequate and effective so far as the needs of its possessor were concerned. The function of these ganglion cells was to control breathing, circulation, feeding, excretion, and reproduction, and the ganglion cells of our sympathetic nervous system are their lineal descendants, performing the same functions for us as the original ganglia performed for our invertebrate ancestors. Now, closely associated with these original ganglia, and practically coeval with them in the scale of evolution, are the endocrine glands. The system thus evolved was the organ of a mind. It received an impression and responded thereto by an act, which, by constant repetition became customary or automatic. If, for example, you inadvertently touch an object which is very hot, you remove your hand with great rapidity, long before the sense of undue heat has reached your consciousness. You perform, in fact, a reflex. The whole of our purely animal physiological life is dominated by these reflexes, situated in the sympathetic or vegetative system. The workings of our viscera, which normally never reach our unconsciousness, are the most orderly and best disciplined things we possess. It is therefore obvious that they must be under the guidance and control of something which for want of a better term we are compelled to call a mind.

Now this unconscious mind is not only automatic, it is also constructive. Arbuthnot Lane has shown us how, in order to counteract undue mechanical strain in the adult skeleton, supporting bands of considerable ingenuity in design and complexity in structure are developed. Their manufacture is quite unconscious, but no one who has studied them could for one moment doubt that they are the result of thought. They are not haphazard; they are beautiful illustrations of the laws of mechanics. Moreover, if there be no such thing as an unconscious mind, how are we to explain the hypertrophy of organs in response to a special demand? How else are we to explain the phenomena of intuition, the process by which people, chiefly women, arrive at a correct conclusion by a jump, as it were? Such a person will tell you that she knows; she cannot tell you how she knows; she says she just knows. And, the devil of it, she does know! The reason is, of course, that the primitive, the visceral, the autonomous or unconscious mind gives information to the conscious, but like a wise judge it announces its conclusions but withholds its reasons. I feel that I must in passing emphasize the fact that our vegetative nervous system, this organ of our unconscious mind, includes practically the whole endocrine system. The ganglia of the sympathetic have the closest possible rela-

tionship with the suprarenals and, through these, with each and all the ductless glands.

Now, I am no psychologist, still less am I a psycho-analyst, but events during and since the war have brought home to me the fact that ordinary physicians cannot afford to neglect the operations of the unconscious mind. In my young days our elders spoke gravely of the influence of mind over matter. They did not know what they meant, and our intolerant youthful scepticism convinced us that the words meant nothing at all. Nevertheless, they do mean a great deal, and they are rapidly coming to occupy an honourable place in the domain of therapeutics.

When M. Coué came to London I went to hear him, and subsequently read the book by M. Baudouin which sets forth the principles of the New Nancy School, of which M. Coué is the originator. To the experienced psychotherapist there is nothing new in this teaching, but to the ordinary physician there is. Or rather should I say that the ordinary physician will find in this teaching the orderly presentation of many facts and considerations of which he was previously cognizant, but so vaguely cognizant as to render his knowledge useless.

We are rapidly passing out of the sector of drug therapeutics in which most of us were brought up, and are being forced to take our bearings in fresh areas. Among these the area of rational psychology is by no means to be neglected. We know something about the anatomy of the unconscious mind; let us study its physiology. Suggestion is a strong living force, and auto-suggestion an even stronger. These forces should not be left to quacks and charlatans to play with. They should be studied, harnessed, and exercised by responsible medical men as a part of the ordinary therapeutic armamentarium. No one has a better opportunity than the general practitioner of utilizing these forces for the public benefit, because no one finds his patients in a more receptive or suggestible condition. Rational psychology should no more belong exclusively to the psychologist than digitalis should be the monopoly of the cardiologist.

REFERENCES.

¹ See H. W. C. Vines, *Proc. Roy. Soc. Med.*, 1922, vol. xv. ² BRITISH MEDICAL JOURNAL, October 21st, 1922, p. 763. ³ *Ibid.*, January 20th, 1923, p. 130.

AN OPERATION FOR ATROPHIC RHINITIS (OZAENA).

(Preliminary Note.)

BY

JAMES ADAM, M.A., M.D.,

SURGEON FOR DISEASES OF EAR, NOSE AND THROAT, STOBHILL HOSPITAL, GLASGOW, AND GLASGOW ROYAL INFIRMARY.

THE pathogeny of atrophic rhinitis is still a matter of dispute; this is due possibly to want of attention to classification of cases and need not now be discussed. What is not in dispute is that, apart from epithelial metaplasia, undue width of nostril—that is, of the whole nasal passage from vestibule to choana, whether this width be primary or secondary—is an important factor in stagnation, drying, putrefaction, and fetor of the nasal secretion. Where the disease is unilateral it is in the wider nostril, or worse in it.

To prevent the formation of the stinking crusts characteristic of the disease various operative measures have been devised. Submucous injection of paraffin sometimes succeeds in narrowing the nasal passages sufficiently to enable the patient to cleanse his nose by blowing and so lessens crusting and fetor. But frequently, and especially in extreme cases, the mucosa will not hold much paraffin, even though the paraffin injection is preceded by one of anaesthetic fluid so as to create a pocket; the paraffin perforates the membrane when more than a little has been injected. Moure tries to meet this by prior vibratory massage of the mucosa for months so as to revivify it. In my hands the paraffin method has been on the whole disappointing. Translation of the inner antral wall toward the septum (Halle) seems rather a formidable measure, hardly likely to be generally adopted.

While paraffin may continue to be used for the less

severe cases, for others a more satisfactory method than either of those mentioned was suggested to me by a case in which the crusts were confined to the left nostril. In this girl, aged 14, the right nostril was occluded by a cone-like deviation of the quadrangular cartilage. The left nostril was extremely wide, at least an inch between septum and antral wall below the anterior end of the middle turbinal; turbinal atrophy was marked. The cone was resected through the right nostril with the idea of replacing it with its apex inverted towards the left nostril. Unfortunately the cone broke, but the pieces were pushed under the muco-perichondrium of the left side of the septum. The improvement that resulted was probably due to the freeing of the right nostril, as the left is still too wide, and it has been agreed to do the operation now to be described—transplantation of rib cartilage.

Flakes of cartilage are, with a wide, sharp gouge, taken from the seventh or eighth costal cartilage, and after incision and elevation of the muco-perichondrium of one side of the septum over the greater part of its extent and similar elevation on the inner antral walls between middle and lower turbinates, these flakes are introduced by suitable forceps—the swan-billed septal punch is excellent—into the pockets thus created, which are packed full. The incisions are closed by careful packing with bismuth gauze, which is left for four or five days.

Attention to the following points helps to guard against risks of septic infection, an important consideration in such cases with a closed wound.

Local anaesthesia should, if possible, be used. An intranasal douche with normal saline should be given before the anaesthesia till no trace of pus can be seen. The nose and face must be cleansed with spirit, and the vestibule painted with iodine tincture. The costal wound is closed with an anchor dressing before the nasal operation is begun. If local anaesthesia is used, then, before the costal operation is begun, the nose is anaesthetized with cocaine and an injection, at the vestibule, of eucaine or novocain solution. The pieces of cartilage as removed are laid on

sterile gauze and not touched except with instruments. They may be an inch or more long but neither broad nor thick; for to secure primary union and retention of the grafts the nasal wounds are best not more than 1 cm. long, placed vertically and just posterior to the skin margin, where they can be wiped with iodine.

I have done this operation in three patients, aged 15, 18, and 60 respectively—once under chloroform, twice under local anaesthesia—with primary union and retention of all the grafts in each case. All the patients expressed themselves pleased with the results in respect of comfort and diminution of discharge; but I am not satisfied, as the cavities are still too wide, though the result is better than could be got by any other method known to me. It is surprising how much cartilage can be packed into the pockets; the cubic capacity of the nasal airway is relatively great. To avoid necrosis of the septum the perichondrium can be safely elevated from one side only at one operation; this leaves the other nostril too wide. This defect will be met in future by storing the excess of costal cartilage in a subcutaneous pocket, whence it can be easily taken a month later for transplantation on to the second side of the septum.

Where only one nostril need be narrowed the operation would probably be better done from the other, narrower, nostril; a bevelled incision, longer than that indicated and carried right through septal cartilage but stopping short of perichondrium of the wider nostril, would allow complete elevation of the latter, thorough packing with transplants, and complete closure of the septal window owing to its bevelled edges and elasticity.

Any treatment of atrophic rhinitis, the ciliary brush being gone, can be only palliative. All that is claimed for the method here suggested is that in well developed cases it is more efficient than paraffin injection and less formidable than Halle's operation. The cases are too few and recent to judge results. The method is published that others who have more opportunity than I now have may try it and report.

ACUTE OTITIS MEDIA WITH JUGULAR BULB THROMBOSIS.

BY

E. WATSON-WILLIAMS, M.C., CH.M., F.R.C.S.E.,
REGISTRAR, EAR, NOSE, AND THROAT DEPARTMENT, BRISTOL ROYAL
INFIRMARY.

THE following case presents some very uncommon features, especially the absence of any of the usual signs of mastoid disease. The notes given explain the main features.

C. G., a girl aged 10, on February 11th, 1923, complained of sore throat, vomiting, and some right earache, beginning suddenly on the previous day. I saw her on the following day; she then had flushed face, dry lips, very furred tongue, and slight tonsillitis; both tympanic membranes were pink but not bulging. There was a little pain in the right ear, and on deep pressure in the postaural sulcus some tenderness, such as an acute myringitis might produce, but no tenderness over the mastoid process, no redness nor swelling there. Constipation was present, the vomiting had ceased. Her appearance was suggestive of scarlet fever. Temperature 102°, pulse 126, respiration 24.

On February 13th, the right ear was discharging. There was epistaxis in the morning and some vomiting; constipation continued. The child looked and felt better. There were no mastoid signs; the tenderness in the post-aural sulcus had disappeared. Temperature 99°.

Left myringitis occurred on February 15th, with rise of temperature to 102.6°. There was discharge from the ear on February 16th; on this day the vomiting ceased.

On February 18th, vomiting began again, with some diarrhoea, general abdominal discomfort, and tenderness, but no distension.

Both ears were discharging; there was occasional slight right earache; no mastoid signs. The tongue and lips were very dirty and dry. The child was not well although the temperature was down to 99°. The case appeared to be possibly abdominal, and enteric was considered.

On February 21st, the temperature, after remaining down one day, rose again, and the child had a rigor in the morning. Still there were no signs of mastoid abscess; but after consultation, in the absence of any other reasonable possibilities, I decided to operate on the right mastoid, as there had once or twice been earache on that side. With the patient under ether, I performed the Schwartze operation. The mastoid cells and bone were absolutely normal; the antrum contained a very little mucoid material. I therefore exposed the ascending part of the lateral sinus; it was normal in colour and contained fluid blood; the wound was packed and left open. Lumbar puncture gave perfectly clear fluid

under slightly raised tension: 15 c.cm. were withdrawn. Pathological report: Slight increase of cells and globulin, no organisms, culture sterile.

On February 24th the temperature remained high and another rigor occurred. Ether was therefore again given, and I exposed the internal jugular vein; it was perfectly normal. The lateral sinus, however, was filled with recent clot to a distance of two inches behind the

genu. This was curetted out, as also the jugular bulb, and free bleeding was obtained from both ends. The sinus was packed off, and the jugular vein tied and divided. A small perisinus abscess was found superficial to the sinus, close to the bulb, and was drained. The bone surface was lightly rubbed with "bipp" and the wound packed. The pathological report stated that both clot and abscess contents grew streptococcus. Later that day the patient's condition was good.

Although subsequently the temperature rose again and I feared that the mischief was going to extend, it fell later by lysis, and convalescence was quite uneventful. Hearing was good in both ears. The child was discharged from the infirmary ward on April 3rd.