subjected, and in a person with an approaching attack of gout it is easy to understand why it is that a joint that has been under such continuous strain for about five hours

Fig. 1.

should be the spot selected at five o'clock in the morning for the exhibition of the most typical symptom.

One member of the profession, on hearing my suggestion as to the cause of ingrowing toenail, facetiously suggested

Fig. 2.

that we had better go to bed with our boots on; but, although we cannot all follow him in that habit, I venture to think that ingrowing toenail might be warded off, especially in children, by the wearing of a specially made toecap.

THE EFFECTIVE TREATMENT OF ACUTE AND SUBACUTE RHEUMATISM.*

By D. B. LEES, M.D., F.R.C.P.,

CONSULTING PHYSICIAN TO ST. MARY'S HOSPITAL AND TO THE HOSPITAL FOR SICK CHILDREN, GREAT ORMOND STREET.

[WITH SPECIAL PLATE.]

THE title of this paper is intended to imply that the treatment of acute and subacute rheumatism, as advised in the textbooks and as usually carried out in practice, is inadequate and unsatisfactory. It is true that a great advance has been made since the days when it was maintained that cases of this disease did as well upon mint-water or upon lemon-juice as with any powerful drugs. In those days, as some of us can remember, there was no great exaggeration in the story of the physician who was asked "What is good for rheumatism?" and who replied, "Six weeks." At that time it was common for rheumatic patients to remain in hospital for weeks, and the very painful and persistent arthritis, with the profuse acid sweats, formed a clinical picture which is now not seen in hospital wards. Fatal hyperpyrexia, now extremely rare, was then not uncommon. There was, however, even then reason for thinking that treatment by alkalies was of some distinct advantage, in spite of the fact that the alkali employed was generally a salt of potash, and therefore not free from depressing effects, and that it was administered only in small quantities. An investigation of the clinical records of Guy's Hospital, which I made in 1874 as the basis of my thesis for the Cambridge M.B. degree, seemed to me to show clearly that the alkaline treatment certainly gave the best results. Yet these results left much to be desired, and the ineffectiveness of the treatment of acute rheumatism was one of the reproaches of medicine.

* Read before the Therapeutical Section of the Royal Society of Medicine, December 1st, 1908.

It is clearly seen by us all, how great an improvement has been effected by the introduction of salicylate of soda. The rapid relief of pain, the fall of temperature, and the shortened duration of the illness are results which follow, we all agree, the administration of this drug. Yet the remedy, it must be confessed, has some side-effects which are annoying and which tend to limit its usefulness. Without much investigation whether these hindrances could be removed by some modification of the method of administration, physicians have been inclined to desert the simple salicylate in favour of some of the organic compounds of salicylic acid provided for us by the ingenuity of German chemists. But the most commonly used of these, acetyl-salicylic acid, the so-called "aspirin," cannot be combined with an alkali, hence we are at once deprived of a useful ally of sodium salicylate; and though free from some of the side-effects of salicylate, it may produce the most dangerous symptoms of salicylate poisoning, as proved by the case of the girl which I have elsewhere narrated who became comatose with typical "air hunger" after seven 15-grain doses of aspirin given in two and one-third days.

The use of sodium salicylate is also hindered by the superstition still prevalent among us that it is a depressant to the heart. The origin of this mischievous notion was probably due, in the first place, to impurities in the drug when a large demand suddenly arose through its early reputation as an antirheumatic. But the drug as now supplied to hospitals is of great purity, and careful observation of its effects for many years enables me to say deliberately that, when given to a rheumatic patient, it does not cause any weakening of the cardiac action. But it must be remembered that the rheumatic toxæmia itself always causes more or less cardiac weakening, and that this may sometimes become specially manifest. If at the time salicylate is being taken, the remedy is almost certain to be blamed for what is in reality one of the effects of the disease. It must be borne in mind, first, that a similar cardiac failure occurs in diphtheria and in influenza, quite apart from the action of any drug, and, secondly, that in every case of acute or subacute rheumatism the cardiac muscle suffers more or less. Endocarditis occurs only in a certain number of cases, and pericarditis is still less common, but the existence of a dilatation of the left ventricle can always be proved by careful percussion. Since I brought forward evidence of this fact before this society in 1898 I have never seen a case of acute or subacute rheumatism, in a child or in an adult, in which dilatation of the left ventricle did not exist. It appears to be invariably present in this disease. For a recent confirmation of this fact I may refer to the instructive article on rheumatic myocarditis contributed to the last number of the *Quarterly Journal of Medicine* (vol. ii, No. 5), by my friend and former house-physician, Dr. Carey Coombs, of Bristol.

The idea that sodium salicylate was a cardiac depressant, conjoined with the frequent difficulties in its administration, have caused physicians to give it only in small doses, and to discontinue its use as soon as or soon after the painful arthritis had subsided and the temperature fallen. It is a common experience that with this method relapses of rheumatic symptoms, in one form or another, have been frequent. It is also a common experience that often the temperature does not fall permanently to the normal, but that the evening temperature persistently rises to 99° or 100°, sometimes for a considerable period, especially in children; this should never be disregarded, for it means that the rheumatism is still active. It is therefore clear that the present treatment of rheumatism is inadequate, for though it controls it does not completely arrest the disease.

But it is not merely inadequate; it is also unsatisfactory, for it aims almost entirely at the cure of the painful arthritis, and pays little or no heed to the heart. It despairs of any real influence on the condition of this organ; it has no hope of diminishing a rheumatic endocarditis, and it usually ignores altogether the affection of the cardiac muscle. This is largely due to the fact that the earlier descriptions of acute rheumatism were the result of observations of the disease not in a children's hospital, but in hospitals for adults. Hence the present absurd arrangement of the textbooks, in which a description of its characteristics in adults is followed by an

article describing what the writers are pleased to call its "peculiarities in childhood." Yet those who have worked simultaneously both in a hospital for children and in one for adults know well that it is in children that the disease is most frequent and most virulent; that in a considerable number of children the first attack is fatal, and that it fatally cripples the heart in a large number of the cases which survive. If every medical student of the past, and especially the gentlemen who now devote themselves to work in pathological laboratories and no longer visit the wards, had been compelled to study for three months in a children's hospital, we should have been spared such phrases as "rheumatism—that is, arthritis," and it would never have occurred to any one to describe as an "attenuated pyæmia" a malady which is as definite and distinctive, and as certainly microbic, as pneumonia or typhoid, which is one of the most virulent diseases of childhood, and which never produces pus.

We must give up the conception of acute rheumatism as a form of arthritis of adults, with occasional metastasis to the heart, and with certain "peculiarities" when it occurs in childhood. We must insist on the conception that it is a microbic toxæmia most virulent in early life, in which the heart is invariably affected to a greater or less degree, but the joints slightly and often not at all, with the peculiarity that when it occurs in adults the most prominent symptom is often a painful arthritis. It was from this point of view that I wrote the article on Acute and Subacute Rheumatism in the first volume of Dr. Allee's *Manual of Medicine*, a few months before the epoch-making discovery of the rheumatic diplococcus by Poynton and Paine, and subsequent experience has only confirmed the conviction that this is the right way of presenting the facts.

If we are to seek for an improvement in the treatment of acute rheumatism, it must surely be based on the great and universally acknowledged curative effect of sodium salicylate in cases of rheumatic arthritis. This drug is not a cure for all kinds of arthritis; the remarkable results obtained by it in rheumatic arthritis are not observed if it be used for a tuberculous, pneumococcal, or gonorrhoeal joint inflammation. The prompt subsidence of a rheumatic arthritis under its use can only be due to a definite antagonism to the rheumatic process or microbe—in other words, its action is specific. The general unwillingness of physicians to admit this really unavoidable conclusion is due to the fact that the doses given are usually so small that only the more easily checked manifestations of acute rheumatism—the arthritis and the pyrexia—are fairly controlled. But if the inference is sound, may it not be possible, by a considerable increase of the dose, to produce a definite arrest of the disease in place of the imperfect control exerted by the small doses given at present? The question seems reasonable, but the possibility of an affirmative reply must depend upon the successful avoidance of the unpleasant symptoms which salicylate is apt to produce. Fortunately these are much less troublesome in children than in adults; indeed, among the toxic symptoms observed in adults vomiting is almost the only one which gives trouble in childhood.

The addition of sodium bicarbonate to each dose of the salicylate, in an amount always double of the amount of the salicylate, is a very effectual means of diminishing these side-effects, and the unpleasant taste may be covered by glycerine and peppermint-water or chloroform-water. The use of sodium bicarbonate in combination with the salicylate has two additional advantages. It helps to neutralize the acid toxins of the microbe, which Dr. Ainley Walker and Mr. Ryffel found, by special culture of the diplococcus, to be partly formic and partly acetic acids. And when we remember Dr. Gaskell's discovery, related in the third volume of the *Journal of Physiology*, that dilute lactic acid causes dilatation of the isolated frog's ventricle and finally arrest in diastole, while dilute sodium hydrate causes increased contraction with diminishing size until the ventricle stops in systole with its cavity reduced to nil, we may be reasonably sure that in man the administration of large doses of sodium bicarbonate must tend to cause a lessening of the pernicious cardiac dilatation of acute rheumatism.

Potassium salts should not be used, for the potassium base is really a cardiac depressant.

It is desirable to administer the combined salicylate and bicarbonate in frequent, but at first in moderate, doses. If the frequency be every two hours from 6 a.m. to 10 p.m., with one additional dose during the night, the total daily dose of each drug will be ten times the amount of the single dose. The amount of each should be increased daily, or every second day, until any unpleasant side-effect such as vomiting, deafness, tinnitus, vertigo, or a tendency to delirium is observed. In children, drowsiness, an acetone-odour of the breath, acetonuria, and a slowing and deepening of the respiration must always be looked out for; if neglected, these may lead to "air-hunger" and fatal coma. Such symptoms are especially liable to occur if the child is costive, and may then be produced by quite small doses of salicylate, especially if the urine is too acid. They may be entirely prevented by careful relief of constipation, conjoined with the administration of a sufficient amount of sodium bicarbonate. Enough must be given to render the urine alkaline, and a surprising amount is often needed to effect this in a rheumatic child. Twice as much as the dose of salicylate is usually sufficient, but occasionally in young children it is not quite sufficient, and extra doses of the bicarbonate alone are necessary to make the urine alkaline. It is also a good rule, suggested by Dr. Reginald Miller, Medical Registrar at the Hospital for Sick Children, never to increase the dose of the salicylate unless the patient's bowels have acted on that particular day.

If, through neglect of these precautions, any of the above-mentioned symptoms of acid poisoning occur, it may be necessary to omit the salicylate altogether, and to increase the amount of the bicarbonate—even to a drachm hourly. With these precautions the administration of large doses of salicylate is quite safe. The only fatal case in my own experience occurred seven years ago, in a child to whom the salicylate was given *without any bicarbonate* and who was costive. The dose of salicylate was very small, only 20 grains daily (four 5-grain doses), but by an oversight it had been continued for ten weeks, and the importance of the prevention of constipation was not then recognized. The credit of first pointing out the danger of constipation in children taking salicylate belongs to Dr. Langmead, formerly Medical Registrar at Great Ormond Street, who published a valuable paper in the *Lancet* for June 30th, 1906.

As soon as any unpleasant symptom due to salicylate occurs in an adult, or if vomiting occurs in a child, the administration of the drug should be suspended for a few hours. Two, or three, doses may be omitted; even a 12-hour interval may be allowed. But the elimination of salicylate in the urine is rapid, and the interval should not be long. After a few hours the medicine should again be given, but in reduced amount, one-half or two-thirds, according to the intensity of the symptoms observed. In many cases even the same dose as before will not produce a recurrence of the toxic effects, and I have sometimes suspected that they may be due in part to a bactericidal influence of the drug, destroying a large number of diplococci so quickly that their internal toxins are thrown in large amounts into the circulation. But it is wiser to reduce the dose of the remedy when it is again given, and to increase it the next day and continue to increase it as before. Almost always it is found possible in this way to reach a much larger dose than caused toxic symptoms at first, sometimes even three or four times as much. If in the course of this increase of dose any fresh toxic symptom occurs, the same plan of arrest and temporary reduction should be adopted as before. There are very few patients who cannot be trained in this way to tolerate large amounts of sodium salicylate, and there is really no need for any of the German substitutes. But one must expect to find every now and then a patient who is specially susceptible to this drug, as others are to mercury, quinine, iodide, or bromide, or even to articles of diet such as eggs.

The initial dose of sodium salicylate may be:

For an adult, 15 grains, or 150 grains daily.
For a child of 7 to 12 years, 10 grains, or 100 grains daily.
For a child younger than 7, 5 grains, or 50 grains daily.
Given in each case with twice the amount of sodium bicarbonate.

The daily increase of dose may with advantage be from 2 to 5 grains in the individual dose—that is, from 20 to

50 grains in the total amount given daily. The increase in the dose should be progressive until the evening temperature is and remains normal. The amount needed for a child may be nearly as great as for an adult, for in a child the infection is more virulent and renal elimination is more active. In some mild cases of acute and subacute rheumatism it may be sufficient to increase the dose of salicylate to 150 grains daily, but in many it is desirable to increase it to 200 or 250 grains, and in some severe cases to 350 or 400 grains. Chorea often requires 250 to 350, or even 400 grains. The doses usually given in this disease are much too small; large doses often cause rapid improvement. But in chorea time is often a necessary factor in the cure; damaged cortical cells require time for the restoration of their nutritive equilibrium after the arrest of the deleterious action of the rheumatic diplococci. A chorea treated early is often promptly arrested by large doses of sodium salicylate and bicarbonate, but a case which is already chronic will require a longer time.

In the chronic rheumatism of adults I have seen improvement commence when the daily dose reached 450 or even 500 grains, while small doses may produce no apparent effect.

The maximum dose in my own experience was given to a boy of 15 years, with severe pericarditis, endocarditis, a much dilated heart, and a tremendous crop of subcutaneous and subperiosteal nodules, many of large size. In this case the dose was raised without any difficulty, and quite rapidly, to a maximum of 600 grains of salicylate with 1,200 grains of bicarbonate daily. This dose was given for two days without any ill effect. After a day free from medicine, it was resumed at the lower rate of 500 grains of salicylate and 1,000 grains of bicarbonate, given for six days in each of the three following weeks. This was followed by a gradual reduction. An icebag was also applied over the heart for the cure of the pericarditis. The final result was most satisfactory.

What are the observed effects of these larger doses of sodium salicylate in acute and subacute rheumatism? Rapid relief of pain and subsidence of arthritis; fall of temperature to the normal with little or no tendency to subsequent rise; rarity of relapse; marked diminution of the torpidity and depressed look of the rheumatic child; and as regards the heart, an easily-observed reduction in size of the dilated left ventricle, while the cardiac impulse becomes stronger and more localized. It would be unreasonable to expect that a murmur due to endocarditis should immediately disappear, but the improvement in the condition of the muscular wall of the heart is often most striking. The rheumatic nodule is specially resistant to treatment, for its abundance of fibrous tissue tends to hinder a bactericide from reaching the diplococci which it contains. Yet under the influence of large doses of salicylate and bicarbonate even large nodules, such as were at one time said to be "equivalent to a sentence of death," will melt away. In the boy already mentioned as many as 216 nodules, some of large size, were counted on one day; yet when he left the hospital not a vestige of them remained. Examples of such nodules will be found in the special plate.

Is it possible to assist the curative influence on the heart by the use of any other therapeutical measure? The answer to this question is very decidedly in the affirmative. In rheumatic pericarditis the extraordinary value of the local application of ice is one of the most certain facts in therapeutics. I advocated the use of the icebag in pericarditis at the Nottingham meeting of the British Medical Association in 1892, and I have used it for the treatment of this condition ever since. Its curative influence is very great. But in using it two points must always be borne in mind. The patient's lower extremities must be kept constantly warm by hot-water bottles, and any considerable dilatation of the right auricle must be relieved by leeches before the ice is applied—the indications for leeches being

an extension of the cardiac dullness in the fourth right intercostal space to two fingerbreadths, rapid respiration, restlessness, and some degree of cyanosis of the lips and face.

The beneficial action of the icebag in pericarditis is clearly a local one, and it suggests the probability that this measure may be useful also in rheumatic myocarditis and possibly even in endocarditis. I think there is no doubt that the application of an icebag to a dilated rheumatic heart (with the precautions already mentioned) certainly causes a more rapid reduction in size of the left ventricle and an increase in strength of the cardiac impulse. And in some cases in which evidence of endocarditis has been present, I have observed under its influence a notable retrogression of the auscultatory signs. The earliest indication of an endocarditis is usually a systolic apex murmur, to which a doubling of the second sound at the apex is often soon added, and in some cases the second element of this double second sound becomes changed into a short mid-diastolic murmur. Under the local influence of ice I have noticed first the disappearance of the mid-diastolic murmur, and the reappearance of the double second sound, soon followed by the disappearance of the doubling, leaving only a systolic murmur and a second sound, and at a later stage the systolic murmur has tended to disappear.

I have gradually made more frequent use of ice in the treatment of cardiac rheumatism apart from pericarditis, and can recommend it as certainly helpful. But the two precautions already mentioned must be always remembered. The combination of ice to the heart with the use of large doses of sodium salicylate and bicarbonate has yielded very remarkable results in practice, and I speak from the

experience of careful trials carried on for three or four years in a hospital service of 60 beds, consisting of 25 beds at the Hospital for Sick Children, with 5 beds for children and 30 for adults at St. Mary's Hospital.

My conviction is that if these methods were adopted universally, the lives of many rheumatic children would be saved, and that an enormous reduction would be effected

in the prevalence of rheumatic heart disease in the adults of the next generation. How much suffering would thereby be prevented can be readily imagined by those who have watched the miserable downward progress of cases of chronic heart disease, the result of former rheumatism.

I commend this method of treatment to your favourable consideration. I submit no statistics, for statistics presented by the advocate of some special treatment are justly suspected as tainted by an involuntary bias. But I ask you to try the method for yourselves. And if to some there seems to be danger in these large doses, I would beg you to remember that an excessive timidity in dosage, due to the fear of doing harm, may often be more detrimental to the patient than a cautious boldness. In medicine a sin of omission is more lightly regarded than a sin of commission, *but it may be even more disastrous*. In advocating a bolder course I have endeavoured to state clearly the precautions which are necessary. Let no one adopt the former and yet neglect the latter.

DESCRIPTION OF PLATE.

Fig. 1 shows many nodules around the elbows, and four or five subperiosteal nodules along the ulna.

Fig. 2 shows large nodules on the occiput, the scapular ridges, and the vertebral spinal processes.

Fig. 3 represents very accurately the many small nodules on the flexor tendons at the wrist in the same patient, but this photograph was taken some years ago from a patient under the care of Dr. J. A. Coutts, and is published with his permission.

A COLLECTIVE investigation on cancer within the boundaries of the State of Hamburg has been made. The material gathered will be edited by Dr. Koerber.

AN Italian Committee for the Study of Cancer has recently been founded. It is under the patronage of the King of Italy, and its Honorary President is the Sindaco of Milan, where are the head quarters of the committee. It is constituted on the same lines as the International Cancer Research Association founded at Berlin in 1908, and its objects are to encourage research on cancer and the publication of the results; to collect statistics on a uniform method; and to promote the establishment of institutes for investigation of cancer.

Fig. 1.

Fig. 2.

Fig. 3.