operation. In those in which the stomach was previously dilated it was found to have returned to the normal, or nearly the normal size. If the stomach tube was passed from one to three hours after a meal, the food contents were found to be still present in the stomach, showing that the gastro-enterostomy opening behaves in the same way as the normal pylorus. The contents were usually found to be neutral in reaction, and showed traces of bile constituents, usually bile salts. If the tube is passed in the fasting condition, a small amount of bile-stained mucus is usually evacuated.

The exceptions to the above ideal results are comparatively unimportant; one, for example, has a ventral hernia, one is still a little anaemic, one has fullness after meals, one suffers from flatulence and occasional attacks of bile (the earliest operation case in the series), and one has a feeling of distension, and his gastric contents after a test meal show butyric acid and the absence of free HCl.

I beg to record my sincere indebtedness to my medical colleagues on the staff of the infirmary, and particularly to Dr. Francis Boyd, for their invaluable co-operation in the clinical examination of the cases here recorded.

PRIMARY DIFFUSE PNEUMOCOCCIC PERITONITIS: TREATMENT BY DRAINAGE AND PNEUMOCOCCIC VACCINE: RECOVERY.

By H. BETHAM ROBINSON, M.S.LOND., F.R.C.S.,

SURGEON TO ST. THOMAS'S HOSPITAL; CONSULTING SURGEON TO THE EAST LONDON HOSPITAL FOR CHILDREN, ETC.

Not many years ago pneumococcic peritonitis was an unrecognized disease, but of later years the attention of the profession has been drawn to its occurrence through the writings of Jensen, von Brunn, and others on the Continent, and by the paper of Annand and Bowen in particular in this country.

The comparatively limited number of cases so far recorded, and the uncertain recognition of the disease, call for the publication of any further experience. I therefore offer the following contribution, which was a case of diffuse and primary pneumococcic peritonitis with an acute onset simulating perforation of the appendix.

Dorothy W., aged 8, was admitted into the hospital on the evening of April 22nd, 1908, under the care of my colleague, Dr. Sharkey, and was transferred to me the next morning.

Personal and Family History.

The father was said to suffer from some chest trouble, and there was a history of tubercle in the mother's family.

She had always been a nervous child, subject to bilious attacks. Three years before she had had measles, and two years before influenza with symptoms very similar to those of the present illness, abdominal pain and vomiting. From the previous November to February she was kept from school on account of her eyes and headache. She had many times complained of abdominal pain, which was generally relieved by micturition; she had rather a habit of holding her water.

History of Illness.

On the day before admission she had been seemingly in perfect health. At 8.30 a.m. on the 22nd she was suddenly taken ill with free vomiting, faintness, and slight abdominal pain centred round the umbilicus; she was very thirsty and feverish. In the evening, fifteen hours after the onset, she was admitted looking ill and complaining of severe abdominal pain and vomiting. The face was very flushed, the eyes bright but not sunken, and the tongue furred generally but rather dry at the tip. The pulse was 130, regular and of small volume; temperature 103.8°, and respirations 36. The chest moved well and equally, and there were no abnormal physical signs detected in heart or lungs. Abdominal movement was restricted but not absent, especially below the umbilicus; there was definite distension, but it was everywhere resonant. Palpation proved an almost universal tenderness, more marked round the umbilicus and in the right lower quadrant; here there was some increased resistance, but no distinct lump could be felt. There was no alteration in the position or size of the liver; the spleen and kidneys seemed normal, and the urine was unaltered except for an increase of urates. Per rectum there was a boggy feeling, especially on the right side, as if there was something in Douglas's pouch.

I was asked to see the child at 11 next morning, her pulse then being 150 and temperature 103.4°. The abdomen moved fairly well on respiration except on the right side below; here and across the bladder region she was tender. There was a

little rigidity over the right rectus both below and above the umbilicus. An immediate operation was decided on, as her condition suggested a perforative peritonitis dependent probably on a gangrenous appendix.

Operation.

A vertical incision was made over the right rectus below the umbilicus, opening the sheath and displacing the muscle inwards. On opening the peritoneum a greenish-yellow fluid escaped, of gummy consistency, with a few flakes of dirty-white lymph in it. Although this was generally diffused through the cavity, and the intestines and omentum were smeared over with it, yet they did not otherwise appear to be in any way altered from the normal; the appendix was absolutely healthy. A good deal of fluid had collected in the pelvis and in the right kidney pouch. There was no sign anywhere of glueing together of the intestinal coils by early adhesions. The right Fallopian tube was very definitely injected. (Subsequent inquiry elicited no evidence, past or present, of any vulval or vaginal discharge.) The glands along the iliac vessels and at the bifurcation of the aorta appeared to be a little enlarged. The peritoneal fluid was quite odourless. Drains with wicks were placed down into the pelvis and up into the kidney pouch, and the main part of the wound was closed in layers. She stood the operation very well, and was put back to bed in the Fowler position, and given $\frac{1}{2}$ grain of morphine. Saline was ordered per rectum.

After-History.

After the operation there was a gentle drop in the temperature to 99° at 8 o'clock the next morning, with a corresponding marked reduction in the pulse-rate. She was feeling much more comfortable, and there had been very little pain and no vomiting. She was given milk and water frequently in small quantities. On April 25th her condition was very satisfactory, but both pulse and temperature had again risen a little. As her bowels had not acted, she was put on magnesium sulphate in the evening. The next morning there was a very slight result after an enema. The wicks were removed from the tubes, and a considerable discharge of turbid fluid came away. Calomel, 2 grains, given at night, followed by magnesium sulphate in the morning; this resulted in four actions during the day. On this day (April 27th) the temperature again began to go up, and, as the clinical report showed that the peritoneal effusion was a pure culture of the pneumococcus, it was decided to use pneumococcic vaccine (see chart). Beyond the variations in pulse and temperature there is now not much to report except her steady progress. On May 5th both drainage tubes were removed and gauze drains substituted. On the 6th, as on removing the gauze a fair amount of pus could be pressed up from the lower abdomen, Bier's cupping for ten minutes was started; this soon produced a marked diminution in the amount of discharge. This, however, did not cease altogether, and apparently came up from the loin, so on May 27th a counter-opening was made here and a piece of gauze pulled through from front to back. This soon had a marked effect both on the discharge and the temperature. On June 6th the front wounds were healed, and on the 10th the drain was left out behind. By this date her temperature had been normal for a week, and she was beginning to get up. She was able to leave the hospital on June 15th.

She was readmitted in capital health on October 19th to have a small ventral hernia through the drainage hole dealt with. Some adherent omentum was set free, the peritoneal bulging cut out, and the freshened fascia brought together in layers. She went out on November 5th with a sound scar.

The clinical types of this disease vary as to whether the peritoneal effusion becomes localized or is diffuse. The localized form starts with acute symptoms, and tends to become chronic; at the same time there is walling-in of the effusion, which usually occurs in the lower abdomen in the region of the umbilicus. This is the type of case which in the past was probably considered to be a tuberculous abscess, and, pointing at the umbilicus, ultimately burst or was intentionally opened, with in many instances recovery.

In the diffuse form the symptoms are more markedly acute and remain so, and the signs those of a generalized peritonitis. Jensen and others have commented on the similarity of the onset to acute perforative peritonitis dependent on the appendix, and this case is another example of such.

These clinical forms of peritoneal involvement are found again, whether the peritoneum is primarily or secondarily affected. In a large number it is secondarily implicated, the primary lesion being most often a pneumonia, or a throat or ear lesion, or a pneumococcal abscess elsewhere; the blood infection from these foci is responsible for the subsequent peritonitis.

In the primary cases of peritonitis the possible origin may be traced in a few cases to an appendicitis or to an enteritis, the initial intestinal symptoms, especially diarrhoea, supporting the latter; neither of these causes appeared to be responsible in this case. As stated before, there was in this patient very great congestion

of the right Fallopian tube, in marked contrast to the surrounding peritoneum. It has been thought that the infection may ascend from below, but here there had never been observed, nor was there present, any vulvo-vaginal discharge. A point that has been urged in favour of this possibility of origin is the curiously undue proportion of cases in females, but this excess is regarded by most as only accidental. The most accepted explanation, however, is that, as in the former group, the infection comes from the blood, and this is infected by absorption from a mucous surface without any discoverable lesion. No further pneumococcal lesion was found in this case, nor was the blood examined for organisms.

The symptoms of onset agreed with those described by previous observers. In a child of apparent good health there was a history of sudden severe abdominal pain, free vomiting, but no diarrhoea, with rapid development of peritoneal signs. The localizing of the pain, first about the umbilicus and then in the right lower quadrant with the most marked tenderness and some rigidity in the same region, seemed to point to the appendix as the probable source of the trouble; in fact, it is difficult to see how one could exclude this view, bearing in mind the many cases seen of a severe first attack and where there had been previously no warning that the appendix was not healthy. In this child there was some previous history of abdominal pain and vomiting, suggesting appendix trouble.

The character of the peritoneal fluid at the operation was quite distinctive; it was odourless and of a yellowish-green colour, of gummy consistency, and with a limited number of greyish-white flakes of fibrinous coagula in it. The fluid was very little turbid, and this is explained by the early period at which the operation was performed; it later became typically purulent, but remained odourless throughout.

The treatment employed was by incision and drainage. Whether it would have been wiser, after mopping out the fluid, to have closed the cavity, as recommended by some, must remain unanswered; there are not sufficient recorded observations to support one view or the other. It may certainly be urged that the closed cavity runs little or no risk of staphylococcal contamination from outside, which is very likely to occur, and did do so in this case. If the fluid should re-collect, or circumstances should otherwise determine it, the wound could be easily opened again and then drained. In another early case I shall be disposed to adopt this line of action.

With the advice and assistance of Dr. Dudgeon, pneumococcal vaccine was employed in the treatment of the case, which was considered a favourable one for such a course. It was started on the fifth day after operation, when, after an initial reduction of pulse, temperature, and symptoms, there was a marked increase again of the pulse and temperature. The injections, eight in all, were given in increasing doses (see chart), at intervals of five or six days. The satisfactory influence of the first two is well shown in the chart; after that the pus became contami-

nated, and it was from this cause that the well-marked rises of temperature occurred, masking the effect of the vaccine. On counter-drainage in the loin the temperature rapidly subsided, and she made a speedy recovery. It must be considered, I think, that the use of vaccine contributed in a large degree to the successful issue of the case.

INCISIONS FOR OPERATIONS ON THE UPPER ABDOMINAL ORGANS.

By ALEXANDER DON, M.B., C.M., F.R.C.S., M.A.,
SURGEON TO AND LECTURER ON CLINICAL SURGERY IN THE ROYAL INFIRMARY, DUNDEE.

THE incisions practised for operations in the upper abdominal organs are numerous, which indicates that none of them are entirely satisfactory. Kocher's and Mayo-Robson's are the most used in this country, and those described by Bevan, Kehr, and Czerny less frequently. The first three have the merit of giving very free access to the biliary apparatus, the duodenum, the pylorus, and the head of the pancreas, while similar incisions on the left side are available for gastric, splenic, and pancreatic operations. The objection to these incisions is that they cut off the nerve supply of the whole upper half of the rectus and overlying skin. A central quadrilateral area undergoes atrophy in consequence. The family doctor knows best that this is no imaginary sequel. It would be of little moment were the patients to remain invalids, but to an active man or woman the injury is serious. Hernia is not common, but the anaesthesia of the skin leads to abrasions, ulcerations, and other injuries, which are more serious from their being less painful, and therefore liable for a time to escape notice and treatment. Kehr's incision leaves a part of the rectus intact, and is in this respect an improvement on the others, but he cuts off more of the blood supply, while Czerny's merely produces another fibrous intersection in the rectus, and leaves a weak median scar.

The following incisions have no doubt been often employed by others, and there seems nothing against their universal adoption. They are anatomically correct, give ample room, and, when healed, produce a firm cicatrix. The blood and nerve supply is intact, the only vessel severed being the deep epigastric artery near its anastomosis with the internal mammary.

The first incision through the rectus is used by most surgeons for exploratory laparotomies. The skin and anterior sheath are divided a fingerbreadth to the right or left of the middle line from opposite the seventh costochondral junction to the umbilicus. The sheath is dissected off the muscle on the inner side, and the muscle freed by blunt dissection from its posterior sheath, and retracted outwards as far as possible. The posterior sheath and peritoneum are then incised as far out as