THE
British Medical Journal.

THE JOURNAL OF THE BRITISH MEDICAL ASSOCIATION.

LONDON: SATURDAY, JANUARY 7, 1905.

A Lecture

ON

THE PAINS OF TABES.

Delivered at the National Hospital for the Paralysed and Epileptic, November 2nd, 1904.

By Sir WILLIAM R. GOWERS, M.D., F.R.S.,

Physician to the Hospital, and Consulting Physician, University College Hospital.

GENTLEMEN,—It is often well to let occasion wait on opportunity, and as I chance to have an unusual number of cases of locomotor ataxy under my care in the hospital at the present time I propose to take some features of this disease as my subject to-day. Its manifestations are many and various, and far transcend the mere inco-ordination which Duchenne recognized as the main symptom of the disease, and expressed in the name he gave to it. This name was apt enough, so far as his perception was concerned, but the extension of our knowledge of its manifestations is so great that we have fallen back on the older Hippocratic term, revived by Romberg, of "tabes dorsalis." It is a word which connotes almost nothing, and so may be made to denote almost anything—a convenient characteristic in medical terminology.

The sensory symptoms of the disease are those which I propose to consider to-day, and especially those which give to the malady a special terror for too many of its subjects—the pains of tabes. They are far more common than ataxy; they begin earlier in the course of the disease, and often make life almost unendurable in cases in which there is no other subjective symptom.

But it is not possible to discuss any manifestation of tabes without alluding to the morbid process which underlies the symptoms of the disease. What this is, indeed, is at present a matter of theory, but the evidence on which the theory rests is strong. I cannot now describe the evidence in detail, but I must give you, in a few words, an outline of what is believed. You may know it quite well, but it is not well to assume knowledge which may not be possessed by all.

The most remarkable feature in the pathology of the present time is certainly the rapid increase in our perceptions of the part played by chemical processes and chemical substances. We are compelled to believe that the symptoms of tabes are due to the influence of a chemical toxin, which results from syphilis and probably arises in a way similar to that by which the toxin that causes diphtheritic paralysis has been proved by Martin to be caused by the organisms of diphtheria. This is due to the indirect action of a ferment produced by the organisms, which influences the albuminous bodies in the process of their production, so that they are altered in constitution, perhaps very slightly, and yet enough to convert them into nerve poison instead of a food. The analogy is the more pertinent because, sometimes, the toxin of diphtheria causes precisely the symptoms of locomotor ataxy. The difficulty in the case of the cause of tabes is the long interval which elapses before it manifests its influence, and the strange way in which this is augmented at times, without any perceptible cause. Still, we may remember that one effect of specific diseases, immunity from recurrence, is lifelong. It is conceivable that a chemical process once set

wrong may be enduring in its perversion, and it is also conceivable that the pathological vitiation may be augmented by causes quite distinct from those which give rise to it.

One fact that I would especially impress upon you is that the effects of this toxin vary in different cases, in a way which is most strange. It compels us to believe that there must be minute differences in the chemical character of the toxin which fit in, as it were, with chemical differences in the constitution of the nerve elements, so as to enable it to act upon some and not upon others, even of similar nature—similar, but not identical.

The toxin, which we are compelled to assume to be the cause of tabes, exerts its influence chiefly on the elements of the lower sensory spinal neuron, which includes the nerve fibres of the posterior roots and the up-bearing (afferent) nerves, all of which depend for their vitality on the cells of the posterior ganglia. These cells determine the life of the sensory fibres from the limbs, and the posterior roots of the spinal cord, some of which end in the grey matter, not far from their entrance, while others ascend the cord in the posterior median column. The morbid influence seems to be exerted on the elements of this neuron, in greatest degree on the extremities of the fibres, which are the parts farthest from the vital centre, and where, therefore, the vital influence may be least. Thus we find the degeneration of the posterior root is greatest adjacent to the cord and within it, and the changes which are found in the distal sensory nerves are chiefly at their extremities. But the most marked degeneration is in the fibres of the posterior median columns, which seem to be the continuation of the fibres from the muscles, and are probably at the medulla connected with the cerebellum. The changes in the fibres which pass from the muscles to the spinal cord, and subserve the muscle-reflex action, on which depend both the knee-jerk and spinal co-ordination, must be assumed as the explanation of the early loss of the knee-jerk, and the peculiar features of the ataxy when this is developed. This brief reference to the pathological causation of the disease must perforce suffice to-day. I hope to return to them another time. I pass now to the symptom which is to-day our special subject.

CHARACTER OF THE PAINS.

The pains of tabes are generally an early symptom of the disease, and they are often most distressing. In rare cases they are trifling, or so slight that they are hardly noticed, but this is the exception. You should know, however, that their occurrence may be denied by a patient who suffers much from them. Only this morning I saw a patient with loss of the knee-jerks, and asked him if he had had pains in his legs. "No," he replied, "I have had no pains, but I have suffered a great deal from rheumatism." On inquiry as to the character of the "rheumatism," it was ascertained that it consisted of the sharp darting pains characteristic of tabes. They had occurred specially on changes in the weather, as these pains so often do, and this had impressed the mind of the patient, and possibly the mind of his doctor, so that their rheumatic nature was considered beyond question. This is a mistake that is common, and one against which you should be on your guard. It is only in the case of pains of moderate intensity that this error can arise. It is not likely in the cases, unhappily frequent, in which their severity is such as to strain to the utmost the patient's power of endurance. Their tendency to occur in paroxysms, unheralded, and of uncertain duration, still

further induces a constant dread which cannot be shaken off, and no doubt increases the actual suffering. All pain that is much feared is felt more severely, for the process of apprehension seems to lower the resistance in the brain and augment the conscious suffering. You may have realized this yourselves, if you have ever had the opportunity of comparing the pain caused by the extraction of a tooth when you dreaded and anticipated it, and when you were resolved to ignore it and resist it. Something of this kind is probably at the root of the endurance of suffering by oriental nations. But this power becomes worn down by recurring pain, and the suffering is then always great.

It is useful to classify symptoms which are various in character. Nature, indeed, is prone to ignore our divisions, and to blend that which we distinguish. Nevertheless, most of the pains of tabes will be found to fall into one or other of the following forms, and their recognition is useful by giving you a better conception of the symptom, and also by aiding your observation. You are compelled to note more carefully when you endeavour to discern features and distinguish forms.

A. The first division of pains consists of those which are brief, commonly momentary, but succeeding each other after a short interval in the same place, commonly for hours, sometimes for days. Of these brief pains there are two important classes, the superficial and the deep.

1. The superficial pains seem to be on the surface or just beneath the surface. They are usually felt at one spot, often only a few inches in extent, sometimes larger, extending down part of a limb. They are most common on the limbs, especially on the lower legs and feet, but frequent also on the thighs, and on the arms and hands, especially on the inner side, that is, on the ulnar region, of the forearm and hand. Sometimes they are felt on the head; in one of the patients before you they occur on any part of the head, on the ears, on the face, and especially about the mouth. They are less common on the trunk than those that are deeper. Each pain is usually extremely brief, a stab or flash of pain, gone as soon as felt, but recurring. It is these pains which have given rise to the common term "lightning pains," an expression which so aptly conveys their character that once heard from a doctor's lips it is employed ever afterwards, and I have often known it to be spontaneously adopted when it has not been learned.

These superficial pains have one remarkable effect: they make the skin tender. After they have recurred a few times, sometimes almost as soon as they begin, the skin in that area becomes so tender that if touched the patient feels as if the place was an open sore. If the finger is drawn gently over it, or the bedclothes are moved upon it, the pain produced is often hardly bearable. A remarkable fact is that this occurs where sensibility to pain is lost. Sensibility to touch is lost in tabes much less frequently than that to pain, or it is only diminished when that to pain is lost. When this is the case, even when the tenderness that attends these pains is at its height, the prick of a pin may occasion no pain whatever. It is evident that the painful sensation is produced through the nerves of tactile sensibility, which indeed can alone be excited by such a surface stimulation. It is an illustration of the fact that almost any nerves, when intensely excited, may give rise to a sensation of the character of pain. You know how true this is of nerves which, under a normal stimulus, cause hardly any appreciable sensation, such as the afferent nerves of muscle, from which such extreme pain arises by their excessive stimulation in cramp. So it is also with the nerves of the pleura, peritoneum, and intestinal wall. Hence it is probable that these superficial pains may themselves be due to a morbid action in the tactile nerves—a fact of some importance. The tenderness of the skin caused by the pains slowly passes away after the pains have ceased, but may last a day or two.

2. The second variety of brief pains is deep-seated. They are not felt in or near the skin, but the sufferer is not able to refer them to any definite structure. Sometimes, however, they are described as being distinctly about the joints, not within them but in their immediate neighbourhood. They present the same features as the superficial pains, since pain after pain recurs at the same part, separated by a short interval. But the pains themselves are not so momentary as those on the surface, they generally last several seconds. One of the patients before you gives a description of them which is often heard—he compares them to a knife in the flesh turned round and round. Another says they are like hooks in the flesh, pulled on for a moment and then slowly

released. Two other of the patients here compare them to a sudden dragging on the tendons, on the side of the foot in one case, in the other on the outside of the leg. Not only do these deep pains die away less quickly than those which are superficial, but after they have gone on for a little time some degree of pain is felt in the intervals. The character of the pain varies very much. Not only is it, as I have said, burning or darting, but it is sometimes described as precisely like the pain of cramp, when there is no muscular contraction. Such pain must be due to the nerve structures through which cramp is felt, the afferent muscle nerves. Remember, however, that any of these sudden pains may cause a reflex start of the limb in which they are felt, but this is purely secondary. In the abdominal wall also there may be secondary muscular spasm, sometimes very distressing.

These deep-seated brief pains are chiefly felt in the limbs, and especially in the legs. They occur in the arms, sometimes in the trunk, but less frequently on the head. They are not attended by the peculiar hyperaesthesia of the skin just mentioned, although sometimes there is a persistent increased sensitiveness in certain muscles, whereby pain is caused by their contraction.

B. The second division of the pains of tabes includes those which are prolonged, lasting for some time—hours, days, or even longer, in the same place. They often present variations in intensity, but not the complete intermissions or considerable remissions presented by the pains of the first class. They are more common in the trunk than in the limbs, but are met with in both the legs and arms.

The best-marked pains of this variety are deep. Prolonged local superficial pains are very rare, although, as we shall see, diffuse surface sensations are not uncommon. But one form must be recognized which is often, though not always, superficial—a girdle sensation at some position on the trunk, commonly at the level of the epigastrium. It is then often accompanied by a change in cutaneous sensibility in the same place, sometimes in a limited zone around the body. When the girdle sensation has a deeper character there is no corresponding change in the sensibility of the skin. Lightning pains are seldom met with in the zone. The fixed trunk pains are often unilateral, and then may give rise to difficulties in diagnosis; they may be thought to be intercostal neuralgia, or evidence of liver disease or of renal colic. The last is not an infrequent mistake when there are exacerbations. Another form which may also lead to error is that which is met with in the sciatic nerve, and sometimes closely simulates the pain of sciatic neuritis. Indeed, not only may it be difficult to discriminate between the two, but it is probable that actual sciatic neuritis does occur, since there is not only pain along the course of the nerve, but definite tenderness of it. I saw such a case a few months ago, and I might possibly have passed it as simple sciatica but for the habit of testing the knee-jerks in all such cases. Such a clinical habit is very useful; the misfortune is that, after finding nothing nineteen times, you neglect it in the twentieth, and find the omission has resulted in a grave error of diagnosis. This patient's knee-jerks were absent. Further inquiry elicited the fact that pains had occurred elsewhere, sharp and darting. His pupils had no light action. It was a case of stationary tabes. Then I learned that his life had been rejected just before at an assurance office, after a very careful examination by the physicians, who doubtless had discerned the nature of the case. No proposer would be rejected for simple sciatica, but no office would accept a case of tabes on account of the many possible perils the future may have in store for even an early and apparently stationary case. This is a digression; but a step down a by-path often gives you a view of some point more useful than you would otherwise obtain.

These persistent deeper pains have many forms. Indeed, the varieties of character they present are such as to baffle description. The enduring sensation is often varied by brief exacerbations, which seem to constitute a link with the intermitting pain. I have already mentioned one common variety, a sense of distressing tension on the muscles or tendons, varied by a sudden pull, and referred by the patient to the "leaders." It seems quite distinct from the cramp-like pain already referred to. It may be felt in the sole, calf, or hamstring muscles. When the deep, persistent pain reaches great intensity it has often a burning character and may be most severe, indeed, hardly endurable. It is often varied by a sensation as if the deep, red-hot iron were pulled and twisted. Such pain may last for twenty-four or forty-eight hours, unless relieved by treatment, and is one of the

most agonizing forms of tabetic pain. Even when an attack is so severe, it is seldom that any exciting cause can be traced.

The second form of fixed pains consists of widely diffused sensations, various in character, seldom amounting to severe pain, but often most distressing, on account of their persistence. They continue for long—for weeks, and even months, varying in degree, but never ceasing. The mind has no rest, and the powers of endurance are taxed by the duration rather than the degree of discomfort. Such sensations are sometimes deep, but more often superficial in character. They are described as tingling, numbness, "swelling," and the like—not rarely as sensations of heat or cold. Some of you may remember a man I showed you last summer, whose life was rendered one long misery by such sensations of painful coldness, especially in the left leg and left side of the trunk. Sometimes an actual sensation is disordered, as when a sensation of pebbles beneath the feet is felt whenever they are in contact with the ground. In other cases there is a deep-seated feeling, which is described as like that which would be "numbness" or "tingling" on the surface, only it is not superficial.

All these tabetic sensations vary much. An almost endless list may be made of them; many classes might be established, but they would serve no useful purpose. The simple divisions I have made will be found, I think, to have practical value, and will, as I have said, give that point to observation which is needed for precision.

TABETIC NEURALGIA.

From the pains of ordinary tabes I pass now to a form of the disease in which the pains dominate the aspect of the malady. If a patient came to you and complained of sharp darting pains in the legs, you would and should think at once of locomotor ataxy, and test his knee-jerks. If you found them normal, it is not unlikely that you would dismiss the idea of tabes from your mind. You might be right, but you might possibly be altogether wrong, and this is my reason for emphasizing the facts I wish to put before you. There is one variety of tabes in which pains are the only prominent symptom; the knee-jerk is not lost, and there is no ataxy. They may be regarded as cases of partial tabes; the symptom which is generally present and characteristic, loss of the knee-jerks, is absent. It may be said that these are simply cases in which the disease is in an early stage and is developing. This does not seem true of the majority of the cases. In those I have had the opportunity of watching for a time, the disease has advanced no further, so far as the ataxic symptoms are concerned, and in only one was there a further change in one knee-jerk. On the other hand, the pains had existed for years when the knee-jerks were still perfect, and we know that they are found to be lost, as a rule, as soon as the pains suggest a suspicion of the disease. Hence I think that these cases must be regarded as the result of a peculiar character of the toxin. They are not merely a stage in the development of ordinary tabes; they are a special variety.

The great difference in the symptoms and course of these cases can only, I think, be explained by assuming a difference in the character of the causal agent. We have another illustration of the varying incidence of the poison in the frequency with which ataxy does not develop when the toxin has a special tendency to act on the optic nerves. The fact has been absurdly misrepresented. The freedom from ataxy has been described as the result of the atrophy of the optic nerves. The only conclusion that is justified is that which I have mentioned, and of which the cases I am now considering afford another illustration. I think that the term "tabetic neuralgia" is an appropriate and unobjectionable designation for these cases.

This form of the malady is not common, but I have notes of eleven cases. Before I summarize their features I may briefly describe to you the last case I have seen. He is a man of 42, and contracted a chancre about twenty years ago, the nature of which cannot be doubted, since he went to Aix for a course of mercurial treatment.—Let me mention, in passing, that you should always be very careful in the way you make inquiries regarding this remote though real cause of the disease. It is easy to ask questions in such a way as to indicate to the patient that it is the ultimate cause of his suffering, but it is not wise to do so. The knowledge may fill him with distress and self-reproach, and in most cases can do no good. Remember that you may convey this knowledge inadvertently, but not less effectively, merely by making a pointed inquiry regarding it as soon as you have

discerned the leading symptoms. Take care to avoid this. Ask about it in the course of inquiries regarding other preceding diseases, inquire casually but carefully; you may easily ascertain all you need without raising any suspicions in the patient's mind.—This patient was sent to me on account of his pains, which began three years ago. They had been felt only in the legs and feet, in various parts; they were sharp momentary stabs; often an attack began as most trifling pains, frequently repeated, which he likened to the ticking of a watch; they gradually grew more intense and less frequent. Often they were felt at several places in the same attack, which lasted for hours, sometimes for days. They varied in severity, but even when least in degree they prevented sleep. They were always superficial, and wherever they were felt the skin became so tender that he could not bear it to be touched. The intervals of freedom varied, but he averred that a change in the weather to damp cold usually seemed to induce an attack. He presented no trace of unsteadiness, and his knee-jerks were active and equal. Sensation on the legs was perfect to touch, but he could not feel the pain of a prick, as such, on the feet and lower legs above the ankles, or on the lower half of the thighs—the prick there seemed only like a touch. Elsewhere pain was felt, although it was delayed near the limit of the analgesia. His pupils were not small, about $3\frac{1}{2}$ mm. in diameter. The right presented no light reaction even to a match; the left responded slightly. Otherwise his condition was normal. He had seen several doctors, and his pains had been ascribed to phlebitis (because he had some varicose veins), to neuritis, and to gout. The doctor who sent him to me had thought of tabes, but had dismissed the idea when he found the knee-jerks were normal.

I may now briefly summarize the facts of the 11 cases I have seen. All were in adult men. A clear specific history was present in about the same proportion as with ordinary tabetics, in 9 out of 11. In no case could the disease be excluded. The interval between the primary disease and the onset of the pains varied from 7 to 19 years, the average being 12 years. The knee-jerks were normal in degree, and equal on the two sides, in 8 out of the 11 cases. In 2 cases they were unequal, less on one side than on the other, but in only one was a further diminution observed. In this, when first seen, the jerk was good on the right side, but on the left could only be obtained with "reinforcement;" six months later it had disappeared on that side, while remaining good on the right, but he did not present the slightest trace of unsteadiness.

Considerable significance attaches to the presence, in these cases, of other symptoms often associated with ordinary tabes. The action of the pupil was not recorded in some of the cases, but in 2 the light-action was lost, in 2 others it was very slight but not absent, in 2 it was completely lost only on one side; in 1 there was instant contraction followed by dilatation to the former size. The exposure to light continuing unchanged, a peculiar reaction which I described many years ago. A significant fact is that one patient presented characteristic optic nerve atrophy. In this case the knee-jerks were normal, and no change occurred during a year the patient was under observation; there was no trace of ataxy, and he could walk ten miles with ease. In him sensation on the skin was quite normal, and it was so also in most of the other cases, but in 1 sensation to pain was lost on the feet and lower legs and lower parts of the thigh; in 1 sensation to pain was delayed on the legs and lost on some parts of the trunk, and in another it was lessened on the left leg and foot. One patient suffered from characteristic gastric crises, and he, with one other, complained of a girdle sensation.

The pains were the symptom for which all the patients sought relief, except the subject of optic nerve atrophy. They presented many of the variations met with in ordinary tabes, and were generally severe. Seven of the cases had superficial pains—intense, momentary, darting—and in six it was expressly said that they made the skin tender, so that, as one patient said, the contact of the bedclothes could not be borne. Another described the feeling "as if fish-hooks were stuck in the skin, and pulled on every few minutes." In three cases of intense darting pains tenderness of the skin was denied. Several patients had deeper pains, one about the joints, and one, who had both superficial and deep pains, expressly distinguished the former as making the skin tender, and the latter as not having this effect. They occurred in various parts—legs, arms, trunk, and in one case were troublesome in the head. In several they were said to be influenced by changes in the weather, as in ordinary tabes, being especially intense on a change to damp cold, as in the case first described.

The duration of the pains before the patient was seen was in three patients one year, in one 2 years, in three 3 years, in one 4 years, and in three 10 years.

The practical importance of these cases is their recognition, and perhaps also their future. In not one of the cases I have seen has the nature of the pains been suspected. They had been ascribed to gout, or to rheumatism, or else to neuritis, which has become a frequent diagnostic refuge. When pains cannot be referred to other causes it is easiest to refer them to a cause that is least generally familiar. But if you are aware of the occurrence of these cases of tabetic neuralgia, and look carefully for the other symptoms which are found associated with tabes; if you test sensibility on the limbs, and ascertain carefully the past history, you will seldom be in doubt as to the nature of the case. Especially important is a minute scrutiny of the character of the pains. Other causes may produce pains resembling the deep, dull, prolonged pains of tabes; but the real "lightning pains," momentary, stabbing the skin, as it were, or flashing along it, and varying in seat, are, I think, hardly ever caused by any other morbid process.

The recognition of the nature of these cases is also important in its influence on your opinion regarding the future. The fact of the tabetic nature of the pains may incline you to a grave prognosis. But I think that the danger of the development of ataxy in such cases is very small. In none of the cases I have seen has there been any trace of unsteadiness. The knee-jerks must disappear before the danger of ataxy arises. Even their loss does not necessarily involve a future defect of co-ordination, for it seems to attend a much slighter degree of impairment of the muscle-nerves than is needed to produce ataxy. We often see, and are able to watch for years, cases in the stationary first stage of tabes, when the knee-jerks are lost, but no trace of ataxy develops. Hence the future of the cases of this tabetic neuralgia, in which the knee-jerks are normal, or at most one only is impaired, may be regarded with equanimity, so far as the danger of inco-ordination is concerned. It is most unlikely that this form will ever deserve the name of locomotor ataxy. But the persistence of the pains, at least in some degree, must be regarded as probable.

SOURCE OF TABETIC PAINS.

Can we discern the structures from which these pains of tabes proceed, or form a reasonable conjecture regarding their source? We must refer them to some part of the sensory neuron which undergoes degeneration in tabes, the neuron which depends for its vitality on the ganglion cells of the posterior nerve roots. We may dismiss the cells themselves from consideration; all evidence goes to prove that the nerve fibrillae simply pass through them and that no nerve energy arises in them. There remain the fibres of the sensory roots which pass into the cord and apparently end there, by contiguity to the structures of the reflex centres or to those which continue the sensory path up the cord. It is not probable that the fibres which degenerate in the posterior median columns conduct any sensory impulses which have a direct effect on consciousness. Lastly, there are the extremities of the peripheral nerves as a possible source of the pains. This origin certainly agrees best with the facts and enables us to understand them most readily. It is in the extremities of the nerves that the sensory impulses normally arise. We know that these nerve endings are the first to fail in their nutrition, perhaps because farthest from the vital centre, perhaps because the toxin has a special influence upon them. Indeed, their degeneration has been adduced as an objection to the origin of the pains in them, because they are found visibly degenerated only in late cases, and the pains are an early symptom. But truly this is what we should expect. The degeneration must begin by minute changes in molecular nutrition, and these must attain a degree relatively colossal before it can amount to alterations that can be detected by the microscope.

Other facts also suggest this source. As we might expect, the pains which are superficial and seem to be in the skin, afford us what seems the clearest evidence of their origin. The fact that their occurrence is attended with the remarkable tenderness which I have described, clearly suggests a peripheral origin. If the pain were produced in the spinal cord it is much less easy to understand the concurrence of the soreness at the spot. Another point which supports the same view is that in some cases a stroke on the tender skin not only causes pain but excites the same flash of intense pain which had occurred spontaneously. Would it not need considerable and weighty

evidence to make us regard this local effect as produced through the medium of the change in the spinal cord?

Evidence to the same effect is afforded by the changes in nutrition which have been observed to attend the superficial pains. We may put aside the grave effects which have been described, such as rapid ulceration of the skin. I have never seen it, and I desire to adduce only what I have observed. One of the patients before you has local darting pains, and the area of the skin in which they are felt almost immediately becomes slightly reddened, and upon it are numerous small white spots, about a sixth of an inch in diameter. Dr. Holmes, our Resident Medical Officer, has seen them, but he has not been able to ascertain whether they are raised or not, on account of the significant fact that the tenderness of the skin is so great that the patient cannot endure the slightest touch. The change is probably of vasomotor origin, and perhaps proves no more than the local soreness, because the effect might be produced from the spinal cord. But a trophic influence of more significance was presented by a patient who was under my care here some years ago. He had attacks of superficial pains on the scalp, each lasting for a day or two. A week after each attack it was found that the hairs at that spot bent suddenly about an eighth of an inch from the scalp, and many of them broke off there. Clearly, coincident with the pains, there was an altered nutrition of the hair, whereby it had not a normal consistence and strength, which were regained when the pains had ceased. The evidence is strong—though I cannot speak of it now—that the trophic influence of nerves is one of direct continuity of their tissue with the structures they influence. The effect on the growth of the hair, coinciding with the pains, shows that there was a local alteration in the function and nutrition of the nerve endings, transient, causing the pains and causing also the imperfect nutrition of the hair. If you consider carefully the facts I have mentioned, I think you will agree with me that they constitute, when combined, strong evidence of the peripheral origin of the pains of tabes.

It may, perhaps, be said that the argument destroys itself. If the pains result from minute changes in nutrition, far below that which the microscope can reveal, they should cease with the establishment of the great alteration which is revealed by visible degeneration. But there may be nerve-endings little changed among those which are visibly degenerated, and also those which subserve the sense of pain may be much changed, and those of tactile sensibility little altered. At least we should expect this condition from the common preservation of the sense of touch when that of pain is lost. Histological research is still backward in the province of pathology. Tactile loss is comparatively rare in considerable degree, but is occasionally met with. I do not, however, remember to have met with superficial pains in a region in which the sense of touch was lost. Thus the degree of change in some nerve endings does not exclude the peripheral origin of the pains in others which are only slightly affected.

If we regard them as due chiefly to the distal nerve endings, the question arises whether other structures take part in their production. I fear this topic would lead us too far into the fogs of physiology. But it is a question whether pains may be produced from the conducting fibres when the degeneration ascends to them. We know that the function of the fibres differs from that of the endings chiefly in degree; they also can respond to a stimulus—that is, in them also other forms of energy can be converted into a nerve impulse. At present we can say no more. The origin of the pain-producing impulses within the cord is also a difficult question. It seems to me improbable that the brief surface pains can arise there, but the deeper pains of tabes, as we have seen, vary so much in character that we may consistently regard them as possibly varying in source, although it may be hard to distinguish those which arise in different parts of a sensory tract through which all sensory impulses pass. Moreover, the share the centres may take in the production of peripheral symptoms will need consideration also in connexion with the loss of sensation, which I have to leave unconsidered to-day.

The pains that may be felt in the trunk of a nerve, especially in the sciatic, may be the result of changes in the nerve-endings in its sheath. These are abundant; they are the source of the pain in sciatic neuritis, and the frequent sciatic pains of tabes show that they suffer with the other afferent nerves. Whether pains may be generated in the conducting fibres of the distal nerves, as degeneration ascends them from their extremities, we cannot say. Such pains

would be referred to the distribution of the fibres, just as are those which are generated in the stump left when a limb has been amputated. We know well that nerve impulses may be generated in conducting fibres, although a stronger stimulus is needed than suffices in the special transforming structures in which they end. A touch on the skin is enough; firm pressure may be needed on the nerve trunk. In the augmented sensitiveness which seems to attend degeneration, we can conceive that pains may be generated, and therefore cannot dismiss the hypothesis as inconceivable, although it may be incapable of proof. Some evidence, at least suggestive, might be afforded by a case of tabes in a patient who had lost a leg, but such a case has never come under my observation.

TREATMENT.

What can we do for these tabetic pains? If you do not ask the question now, you will ask it, and anxiously, when a case comes under your care. The treatment of the pains is partially separate from that of the disease, and so I may briefly speak of it; but prognosis and treatment are intertwined, and there is one fact which it is well to make clear to the patient. Although the pains can often be lessened, they can seldom be altogether arrested. The tendency to them continues, as a rule, in some degree. They come and go, but an increase in the pain does not necessarily indicate any increase in the underlying morbid process. It is important that this should be made clear, for the patient is prone to think that he is essentially worse whenever an attack of pain occurs more severe than usual; but the pain has no necessary proportion to the disease. The altered structures seem to be brought into a state of instability so great that they are susceptible to other morbid influences—gouty, "rheumatic" (as we say), and those due to digestive derangements and constipation, which often have a conspicuous relation to the attacks of pain.

Only the superficial pains can be relieved by local measures. Chloroform, sprinkled on lint, with oiled silk over it, generally gives some temporary relief. More lasting is the effect of a hypodermic injection of cocaine at the upper part of the area in which the pains are felt. A quarter or third of a grain usually stills the pain for one or two hours, and often the attack ceases. It is probable that some effect would be produced by the electrical administration of cocaine. If rubbed into the skin, in whatever combination, it has hardly any influence on sensibility. Voltaic electricity has no influence on the pains whichever pole is applied. But if the positive electrode of a voltaic battery is saturated with a solution of cocaine, 6 or 10 per cent., the sensibility of the skin is abolished to both touch and pain in ten minutes, and after a longer application the effect continues for about twice the time the current has been allowed to pass. This was ascertained some years ago, and I have lately verified it, although independently. With the negative electrode no such effect is obtained. I am sure that this deserves a careful trial in cases of tabes with superficial pains and tenderness. I have found it give speedy relief.

Applications to the surface have not much effect on the deeper pains. A deep injection of cocaine has little influence on severe pains which are in the softer substance of a limb, perhaps because it cannot be directed accurately to the nerves involved. But where the tissues are less in amount, as in the foot, pains that seem deep and do not cause tenderness of the skin may be stilled in a few minutes by an injection at the upper part of the region to which they are referred. As a rule, these deep pains are amenable to internal remedies, which are also useful for those which are referred to the skin. Of the various pain-stilling agents, the well-known coal-tar derivatives stand above all others, at least for pains of moderate degree. It is indeed strange how high a position these agents have attained in general estimation. Unknown twenty-five years ago, they are now among the agents most frequently used. They were inventions rather than discoveries; they were new combinations, artificial products, which, as far as we know, had never existed before. The first purpose for which they were used was to reduce the temperature in pyrexia, as the names of two of them still imply. But their power to relieve pain was soon discerned, and quickly superseded their antipyretic influence. It is almost wholly as pain-stillers that they are now used. They lessen many kinds of pain without causing sleep.

Moreover they maintain their place. Many things come into fashion, obtain for a time, and pass away, forgotten. Survival is a sure indication of value. That which is really useful, stays.

Phenacetin, antipyrin (or phenazone), and antifebrin (or acetanilide) are those which are most used and merit most confidence. They are of great service in most tabetic pains, unless they are of great intensity. One may succeed when another fails. The order in which I have mentioned them is the order in which they are most used, but the order of their efficiency is, I think, the reverse. Acetanilide is certainly the most effective and the least prescribed. This is possibly because it is more potent and has obtained the reputation of being less safe. I have not myself observed any evidence to support the opinion. Certainly, however, 10 gr. of acetanilide may be regarded as equivalent to 15 gr. of antipyrin. It is an element of more than one patent medicine for the relief of pain. Of each of these agents it may be said that the effect of a given quantity is greatest if it is given in a single dose. Its influence is also greater if given at the beginning of an attack than when the pains have become well developed. Thus employed they are most useful. Sometimes, however, the suffering is so intense as to resist every agent except morphine, and the patient's suffering may be so severe as to necessitate this agent, and to render even large doses necessary to quell the pain. It is strange that the influence of morphine seems to be "used up" in stilling pain. A dose, say, of $\frac{1}{2}$ gr., which would be hardly safe in health, so deep a sleep does it induce, may cause no sleep whatever if its sedative influence has been expended on severe pain. But such intensity as to compel morphine is rare, and neither morphine nor opium should be allowed to be taken at the patient's discretion.

Besides the relief of the tabetic pain when it occurs, something may often be done to lessen the tendency to its occurrence, to make it less frequent and less severe. The agent which I have found most effective for this purpose is the chloride of aluminium, given in a dose of 5, 8, or 10 gr. three times a day. It is readily soluble in water and may be combined with most other drugs. I can give no explanation of its action. I tried it just simply because chloride of ammonium also sometimes has a similar, although feeble, effect—a reason which you will say, and I admit, is not rational therapeutically. Yet that which is purely empirical sometimes deserves esteem. A similar influence may be obtained, in some cases, by the salicylate series of drugs, the alkaline salicylates, and also the salicyl-acetic acid called aspirin. The latter has the disadvantage of insolubility, but may be given suspended with mucilage. The influence of these agents is curious in connexion with the relation of the pains to changes in the weather.

The use of iodide of potassium or mercury belongs to the treatment of the disease rather than to that of the pains. I shall hope to refer to it on another occasion, but need only say now that I have seen no clear evidence of good from either. At present, moreover, I cannot say more than that we have still to gain experience regarding modern therapeutic appliances which have been thought, or may be thought, to have some capacity of relief. I refer to high-frequency electric currents, the so-called light baths, and radium. The last requires care. One patient here fancied he derived more benefit than we could discern, and when he left the hospital he purchased some. In spite of warning, he put the tube containing it against his skin at night and had some troublesome ulcers in consequence. Remember that local sores and inflammations may develop in these patients without the warning which pain supplies in health. It is indeed strange that this danger signal should be absent in a disease that is one of the most pain-giving. But the subject of the sensory defect in tabes, and its relation to the special symptom, ataxy, I hope to consider when we meet again.

RUPTURE OF THE SPLEEN.—An inquest was held at Aldershot on December 28th, 1904, on the body of a trombone player belonging to the mounted band of the Royal Artillery. He was admitted to the Cambridge Hospital on December 21st and died on December 26th. He was brought in unconscious, having become so in the course of a kind of fight with a band-boy. The latter in self-protection hit the deceased with a Wellington boot, and the deceased continued to fight until he became unconscious. The medical evidence showed that death was due to a rupture of the spleen. The deceased, it was stated, had served in India, and had there suffered from malarial fever. The jury returned a verdict of death from misadventure, and added a rider to the effect that the band-boy had used no undue violence in defending himself from attack.