widened, or not, softened, torn, and dilapidated by ulceration, gangrene, or unavoidable injury in fingering. Occasionally a concretion is found within, or escaped by perforation into the abscess or exudation. Concretions of any kind are exceptional, but are more often of faecal composition, and more rarely calcareous. The steps for removing the injured appendix are similar to those described above, allowance being made for the altered circumstances. The foul surfaces are stuffed with gauze before the protective packs are removed. The wound is unsewn, or only slightly so, and the plug is left from one to three days, according as it is loose or adherent. The unaffected peritoneum becomes sealed around the plug. A local cleansing before plugging may be attempted by applying pure liquid carbolic acid after wiping the foul surfaces dry, and followed immediately by the application of a rag wetted with methylated, rectified, or absolute alcohol. The carbolic acid acts instantly, and the alcohol removes what remains of it and prevents damage by it. The after-treatment demands a diet for a day or two of little more than drinks of water or tea with or without a little weak alcohol; sometimes only sips of these liquids. Additions of nutritious fluids are in severe cases only cautiously and gradually made, as appetite returns and the state of the bowels permits. The dressings require removal every day, and the dry gauze pack, after its first removal, is to be changed at intervals of a few hours or a day or two, according to the amount of discharge. The use of x-roform powder aids the disinfecting and healing process, under such circumstances.

AN OPERATION FOR APPENDICECTOMY.

By A. W. MAYO ROBSON, D.Sc., F.R.C.S.,
Senior Vice-President, Royal College of Surgeons of England.

So many different methods of performing appendicectomy have been described that one almost hesitates to add to the list, yet, as I have been told by several experienced surgeons both at home and abroad who have witnessed the operation I am about to describe, that they had never seen this particular method, I think perhaps a knowledge of the ease with which it can be carried out, and of its efficiency, may afford some excuse for my describing the operation that I have done in chronic appendicitis or in the intervals of acute attacks some hundreds of times with universally satisfactory results. Moreover, by the adoption of this method, it is safe to promise that the patient will be, under ordinary circumstances, on the sofa in ten days, and able to return home within the fortnight without risk of hernia or abdominal weakness.

Fig. 1.—To show line of incision.

An oblique incision of 3 to 4 in. (Fig. 1) is made about 2 in. internal to the right anterior superior spine of the ilium down to the aponeurosis of the external oblique, which is then split in the direction of its fibres to an extent equal to the external incision. The fascia over the internal oblique is divided in the direction of its fibres, and the internal oblique and transversalis are then split in the same direction, so as, when the margins are forcibly stretched, to make a lozenge-shaped opening with the peritoneum as its floor. As a rule there is

no bleeding, but if any vessels bleed they are at once ligatured before opening the peritoneum.

The peritoneum is then caught up by two pairs of pressure forceps and brought to the surface, when it is incised between the forceps and the opening is increased to the extent of an inch to an inch and a half (Fig. 2). Through the

Fig. 2.—Peritoneum exposed in floor of space obtained by splitting the muscles.

opening the caecum is seen and drawn up. By following the longitudinal band the appendix is soon discovered, and any adhesions are separated by the finger, or, if needful, ligatured. As soon as the appendix is brought to the surface any protruded caecum or colon is returned into the abdomen, leaving the appendix and the apex of the caecum protruding through the lozenge-shaped aperture in the abdominal wall (Fig. 3).

Fig. 3.—Caecum and appendix projecting through lozenge-shaped space obtained by splitting the muscles.

The mesentery of the appendix is then seized with pressure forceps and divided between the forceps and the appendix, a ligature being applied to the stump. If the mesentery is very long it may need to be ligatured in two, or even three, portions; but, as a rule, one ligature is sufficient (Fig. 5).

The root of the appendix close to the caecum is then seized between the blades of a pair of ordinary Spencer Wells pressure forceps and crushed (Fig. 4), the forceps being taken off and reapplied at right angles, in order to make the crushing more complete, so as to leave only the peritoneal

coat in the crushed portion. To this crushed base of the appendix, close to the caecum, an ordinary thin catgut ligature is applied and cut short (Fig. 5). A purse-string suture is then inserted around the root of the appendix (Fig. 6), but not tightened until the appendix has been removed, which is effected by seizing it with the pressure forceps just above the crushed portion and cutting it through between the ligature and the forceps. The ligatured stump of the appendix, consisting merely of a little nipple of tissue, is depressed by means of a probe or a pair of Lister's sinus forceps and the purse-string suture tightened.

Fig. 4.—Forceps applied to root of appendix.

Fig. 5.—Crushed portion of root of appendix with ligature applied close to caecum.

This entirely buries the ligatured root of the appendix, making the surface of the caecum appear as in Fig. 7. For additional security another purse-string suture may be applied, this taking up the root of the mesentery so as to make it form an additional cover to the tip of the caecum. The ligature of the crushed stump of the appendix is not absolutely necessary, as the crushing quite seals the lumen of the appendix. If thought desirable, it may therefore be omitted, but personally I usually apply the ligature.

Fig. 6.—Purse-string suture applied but not yet tightened.

Fig. 7.—Dimple seen after tightening the purse-string suture.

The caecum is then returned, and the peritoneum closed by a circular suture, after which the muscles, none of which have been divided, fall together. Two or three interrupted catgut sutures are applied to unite the separated margins of the internal oblique, and then several sutures are in like manner applied to the edges of the aponeurosis of the internal oblique muscle to secure accurate apposition, after which the skin is united by several interrupted sutures.

A dry aseptic dressing is then applied, and not disturbed for ten days, at which time the patient may be allowed on the sofa. The damage to the abdominal wall is so slight that no belt is necessary or even desirable.

In acute cases, where the root of the appendix cannot always be safely brought to the surface, I have found that crushing the root of the appendix and applying a catgut suture to the stump has been sufficient to secure freedom from a faecal fistula; but in the usual operations in the quiescent period, as described above, I always apply the purse-string suture in addition.

It will be seen that the first part of the operation, the splitting of muscles, is similar to the one advocated by Dr. McBurney, but the treatment of the appendix which I have employed for years was original on my part, though I believe it does not differ much from the method pursued by Dr. Ochsner of Chicago and by Dr. Doyen of Paris, both

of whom probably arrived at the operation as I did, independently of one another.

If there are no special complications to cause delay, the operation can be readily completed in from ten to fifteen minutes. Where the diagnosis is doubtful, and it may be necessary to examine the abdominal contents more thoroughly than is generally necessary in a simple operation on the appendix, a vertical incision through the outer part of the rectus sheath is employed and the muscle is displaced inwards, when the posterior sheath of the rectus and the peritoneum are divided together in a vertical direction; in this way a free opening can be made large enough to admit the fingers or even the hand without dividing any muscle. When the operation is finished, the posterior and anterior parts of the rectus sheath are united separately, and a strong abdominal wall is left. Through this incision the appendix, the pelvic organs, the kidney, and the gall-bladder region can be readily explored, and if needful to do any further operation on these organs the incision can be extended without division of muscle or serious weakening of the abdominal wall.

UNUSUAL CASE OF SUPPURATIVE APPENDICITIS.

By GEORGE A. CLARKSON, F.R.C.S.ENG.,
Leicester.

THE appendix being liable to considerable variation in position, the abscess resulting from inflammation of that body is liable to extend in many directions. In the following case the abscess not only produced intestinal obstruction by pressure on the anterior wall of the rectum, but subsequently having perforated the pelvic fascia forming the sheath of the psoas, the pus passed below Poupart's ligament in the sheath of that muscle.

A man, aged 32, was taken ill in February, 1901, with a mild attack of appendicitis. With rest and treatment the symptoms gradually subsided, and at the end of a month the patient was able to get about again.

At the beginning of May, 1901, there was a relapse, and before long difficulty was experienced in getting the bowels to act, and the patient suffered much discomfort and pain. Aperients and enemata had less and less effect, and by May 19th there was almost complete obstruction. This was due to pressure on the anterior aspect of the rectum by a fluctuating swelling, which could easily be felt by the finger and which was obviously an abscess. The walls of the rectum were completely flattened against the sacrum. The amount of swelling in the right iliac region was very slight. On May 20th the abdomen was opened in the Leicester Infirmary, and an abscess containing extremely fetid pus was found behind the bladder. Nothing could be seen of the appendix owing to adhesions. The abscess cavity was drained, and the wound closed in fourteen weeks.

After a lapse of two years—that is, in July, 1903—a fresh attack occurred, and a large inflammatory mass formed in the right iliac region. This gradually subsided, but by the middle of September a large mass was again present, accompanied by severe constitutional symptoms, great pain, and consequent loss of flesh and strength. The pain, which was so intense as to necessitate the free use of morphine, extended down the right thigh as far as the knee. The thigh then became gradually flexed on the pelvis, and the least movement of it caused great suffering.

As the patient desired that an attempt should be made to relieve his condition, on October 1st I opened the abdomen over the inflammatory mass. The caecum and adjacent coils of small intestine were found densely adherent in all directions, and the active trouble apparently lay behind and internal to the caecum. As an attempt to separate the adhesions nearly resulted in the bowel tearing, the wound was closed. Following the operation, the right thigh continued to become more and more flexed on the pelvis, the severe pain persisted, and profuse sweats occurred.

Early in December swelling and fluctuation was obvious below Poupart's ligament, and an abscess was opened in the sheath of the psoas containing more than 20 oz. of very foul-smelling pus. This gave great relief to the pain, but the patient continued to emaciate, and the thigh remained in the same flexed position. Large quantities of pus were discharged daily, and the patient died of exhaustion on April 29th, 1904.