A skiagram showed the dislocation of the right hip into the sciatic notch, and also the fixation of the left hip in the acetabulum in a position of abduction, a position which compensates for real shortening by a tilting down of the pelvis on that side.

[Note.—The dislocated right hip was reduced by manipulation under an anaesthetic on April 2nd, and put up in a position of eversion and abduction.]

CONCLUSION.

I would summarize the points which I have learnt from this case as follows:

1. The danger of trusting to an "aseptic" system of surgery when dealing with the region of the groin. I now always use 1 in 2,000 biniodide of mercury as lotion for these cases when asepsis is doubtful.
2. The absolute importance of an exact bacteriological diagnosis in every case of suppuration.
3. The great gravity of *Bacillus pyocyaneus* infection.
4. The potency of vaccinal treatment of this infection.

A CASE OF TUBERCULOUS MENINGITIS WITHOUT TUBERCLES.

By F. W. HIGGS, M.B.LOND., M.R.C.P.LOND.,

SENIOR ASSISTANT PHYSICIAN, BELGRAVE HOSPITAL FOR CHILDREN;
MEDICAL REGISTRAR (LATE ASSISTANT CURATOR OF MUSEUM),
ST. GEORGE'S HOSPITAL.

[From St. George's Hospital and Clinical Laboratories.]

THE following case of tuberculous meningitis seems worthy to be recorded, as it exhibited several unusual and important features:

C. M., a boy aged 8 years, was admitted to St. George's Hospital under Dr. H. D. Rolleston. The history obtained from the patient's mother was as follows: Onset gradual and indefinite. The child had "two severe frights" six weeks and two weeks respectively before admission to hospital. Since the latter the patient has been "morbid," and has complained of pains in the head and has vomited frequently. During the last few days it has been increasingly difficult to get the child to take his food. No history of any head injury. No previous illness except measles. History of tuberculosis in family of maternal grandmother.

Condition on Admission.

Child looks very ill. Lies quite listless, but cries when disturbed. Occasionally shows signs of understanding what is said to him. Some external strabismus of right eye. No voluntary movements of eyes performed. Edges of optic discs blurred, but no definite optic neuritis. Some paresis of left side of face. Occasional twitching movements of orbicularis oris, and clonic movements of the muscles of the upper extremities and of the hamstring muscles on both sides. No note of any head retraction or stiffness of neck. Muscles well nourished. No definite loss of power in any group of muscles. No hypertonicity. The legs are kept flexed at hip and knee, but can be readily extended passively. No loss of pain sensation. Knee and ankle jerks absent. Plantar reflex gives flexor response on both sides. Superficial reflexes absent. Incontinence of urine and faeces. Respiration rapid (36 per minute). Pulse 128. Temperature 102.5° F. Urine acid, specific gravity 1020; no albumen; chlorides normal. No abnormal signs in chest and abdomen. Lumbar puncture was performed on the day after admission, and one and a half test-tube full (about 30 c.cm.) of turbid flaky fluid was withdrawn. The patient became progressively worse, and died comatose two days after admission with temperature 100.6° F. and respirations 48. The bowels had acted normally once or twice a day while the child was in hospital, and no further vomiting had occurred.

Post-mortem Examination.

This was performed nine and a half hours after death. Fairly nourished body. Rigor mortis present.

Meninges.—Some excess of turbid fluid at the base of the brain. The membranes generally are hyperaemic and "sticky" in appearance. No exudate or tubercles on the convexity. A thick layer of fibrino-purulent exudate is present on the inferior surface of the pons. Similar exudate in rather less amount is present in the interpeduncular space. The superior surface of the medulla is adherent to the middle lobe of the cerebellum. There is also similar exudate to that at the base around the vessels in the pia mater deep in the Sylvian fissures. A careful search reveals no tubercles anywhere on the vessels of the meninges or in the exudate. Brain 47 oz., hyperaemic and rather soft. Moderate dilatation of the ventricles, which contain excess of slightly turbid fluid. No tubercles are visible in the choroid plexuses, which appear normal. The brain is not otherwise abnormal. There are no tuberculous foci in it.

Spinal Cord.—The spinal meninges contain a slight excess of turbid fluid, especially in the lumbar region. The cord itself looks oedematous but otherwise normal. No tubercles are visible in the spinal meninges and there is no fibrinous or purulent exudate thereon.

Thorax.—Pleurae normal. Lungs (right 10 oz., left 8 oz.) are congested. There are no signs of miliary or other form of tuberculosis in them. Around the left main bronchus is an actively caseating tuberculous gland, the size of a Barcelona nut. Its capsule appears intact.

Larynx, trachea, and thyroid, normal.

Pericardium normal; heart 6 oz., normal, except for moderate dilatation of the right ventricle and tricuspid orifice.

Abdomen.—Peritoneum and mesenteric glands normal; no tubercle. Alimentary tract normal; no ulceration or tubercles. Liver and kidneys normal. Gall bladder and ducts normal. Pancreas congested. Spleen 2 oz., congested; consistence normal. Suprarenals normal. Bladder normal, distended; no obstruction to outflow of urine.

No surgical tuberculosis present. All the organs were searched for miliary tubercles, but none were seen.

Microscopical Examination.

Sections of the cortex of the brain were made, showing also the meninges and exudate. Acute inflammation of the meninges and superficial brain substance was present. The meningeal exudate consisted of fibrin and round cells. The cells were mainly large and small lymphocytes, but some polymorphonuclear leucocytes were present. The cells were not spread uniformly throughout the fibrinous mass; in some parts they were thickly aggregated in large irregular clumps, while in other parts there was fibrin with comparatively few cells scattered in it.

All the arteries were surrounded by collections of round cells, and the arterial walls themselves were infiltrated with leucocytes, but no definite endarteritis was present. Many of the cells of the exudate took the stain badly, and appeared necrotic, but nothing suggesting miliary tubercle formation was seen; there were no giant cells and no epithelioid cells seen in the sections examined. Similar sections stained by the Ziehl-Neelsen method showed the presence of a considerable number of tubercle bacilli scattered throughout the cortex and exudate (each field containing 12 to 20 bacilli).

The organisms were especially numerous in the close vicinity of the blood vessels.

The examination of the cerebro-spinal fluid removed a few hours before death (the report on which was not received until after the necropsy) showed that the cells were chiefly polymorphonuclear leucocytes, there being only a few lymphocytes. A considerable number of tubercle bacilli were found in the fluid after centrifugalization, and no other organism was present. The lungs and other organs were unfortunately not examined for isolated tubercle bacilli.

The unusual features of the case are: (1) The presence of tuberculous meningitis with much fibrino-purulent exudate at the base of the brain without any miliary tubercle formation; (2) the absence of miliary tuberculosis of the lungs and other organs; and (3) the great predominance of polymorphonuclear leucocytes over lymphocytes in the cerebro-spinal fluid.

The child only lived two days after admission to hospital, but a diagnosis of tuberculous meningitis was made from the history and physical signs. At the necropsy, however, doubt was thrown on this diagnosis, as, although there was an actively caseating bronchial gland to serve as a primary focus, there was no generalized miliary tuberculosis and no tubercles in the meninges, which, moreover, showed much thick fibrino-purulent exudate, mainly posterior-basal in distribution. On the other hand, the general "stickiness" of the meninges and the presence of some exudate around the vessels in the Sylvian fissures favoured the diagnosis of tuberculous meningitis. The doubt as to the diagnosis was accentuated by the cytological examination of the cerebro-spinal fluid. The bacteriological examination of the centrifugized fluid and the sections of cerebral cortex settled the question.

The presence in the sections of cortex and meninges of "diffuse leucocytic infiltration, predominantly perivascular," as described by Delille, is noteworthy, as also is the presence of inflammatory cells scattered diffusely throughout the walls of the arteries. There was, however, no obvious tendency to the formation of circumscribed nodules consisting only of round cells without giant cells or epithelioid cells, such as is sometimes seen in miliary tubercles in the lungs and elsewhere in the more acute cases of generalized tuberculosis.

A case reported to the Société Médicale des Hôpitaux de Paris by MM. Sireday and Tiné¹ is similar to the present case in that there was tuberculous meningitis without tubercles, and microscopically there were numerous tubercle bacilli with cellular exudate, "chiefly perivascular," but in their case there was little fibrinous or purulent exudate in the meninges, and "diffuse tuberculous granulations" were found in the lungs. Further, the cerebro-spinal fluid contained chiefly lymphocytes, as is usual in cases of tuberculous inflammation. The

absence of miliary tubercles in the meninges in tuberculous meningitis, though a recognized pathological occurrence, is comparatively infrequent, and especially so when associated with their absence also from the lungs and other organs; and the predominance of polymorphonuclear cells in the cerebro-spinal fluid is certainly rare in tuberculosis.

With regard to the reason why tuberculous granulations are not always found in cases of generalized tuberculosis, it may be considered that while general tuberculous infection usually takes the form of a *pyaemia* it may occasionally be only a *septicaemia*, and that the primary tuberculous focus in the first case discharges into the circulation clumps of bacilli with tissue debris, and that these clumps, becoming impacted in the smallest arterioles, cause the formation of miliary tuberculous granulations or "tuberculous pyaemic abscesses" around the blocked vessels, but that in the second case separate bacilli only are discharged into the circulation in considerable numbers, and that these isolated bacilli do not get lodged in the small vessels. The diffuse leucocytic infiltration of the cerebral cortex and the cellulo-fibrinous inflammation of the meninges may be explained as being the result of the irritation of the toxins, and of the bacilli, which have found their way from the blood vessels into the cortical substance and into the cellular tissues of the meninges, these parts especially easily reacting to the tubercle bacillus.

I wish to express my thanks to Dr. H. D. Rolleston for permission to publish this case, and to Dr. Chas. Slater for furnishing the report of the bacteriological examination.

REFERENCE.
¹ *Lancet*, April 13th, 1907.

OBSERVATIONS ON AN OUTBREAK OF MEAT POISONING AT LIMERICK.

By E. J. MCWEENEY, M.A., M.D., R.U.I.,
 D.P.H.IREL. CONJT., F.R.C.P.I.,

PROFESSOR OF PATHOLOGY AND BACTERIOLOGY, CATHOLIC UNIVERSITY
 MEDICAL SCHOOL, DUBLIN; BACTERIOLOGIST TO THE LOCAL
 GOVERNMENT BOARD FOR IRELAND.

In view of the economic and hygienic importance of the subject of food poisoning, I have thought it well to bring the leading facts revealed by my study of this disastrous outbreak (73 cases with 9 deaths) before a wider circle of the professional public than could be reached by the *Journal of Pathology and Bacteriology*, in which the fuller communication which I have prepared on the subject will appear.

I.—THE OCCURRENCE.

On Tuesday, November 3rd, 1908, at 6 p.m., an alarming attack of illness broke out amongst the inmates of an industrial school for girls at Limerick. The symptoms, ushered in by severe headache, were those of acute gastrointestinal disturbance—nausea, abdominal pain, uncontrollable vomiting and diarrhoea accompanied by tenesmus, and followed, in the severer cases, by the most profound collapse. By midnight, some twenty-eight of the girls were stricken down, and on the following day nearly fifty more became affected, though for the most part not so severely. The first death (that of S. K., aged 13) took place at 7 on the Wednesday morning, after little more than twelve hours' illness. Another child succumbed at 9 that night. On the following day, Thursday the 5th, no fewer than 7 more deaths occurred in rapid succession, the last (that of M. K., aged 13) ensuing at 6.35 p.m., about forty-eight hours after the commencement of the outbreak.

II.—CLASSIFICATION OF THOSE AFFECTED.

The pupils numbered 197, and were divided into four classes. The first or senior class comprised 67 girls between the ages of 13 and 17, and on it fell the brunt of the attack, with 55 illnesses and 8 deaths. The second class, 48 in number (ages between 10 and 13), yielded 10 cases with 1 death. In the third class there were 51 little girls between 7 and 10 years old, of whom 8 sickened—5 decidedly, 3 somewhat doubtfully. The fourth class comprised "babies" between the ages of 3 and 7. There were 31 of these, none of whom took ill. Nor were there any cases amongst the Sisterhood in charge, teachers in training, or servants. Of the 73 cases,

9 died as above stated, and the remainder rapidly recovered, after several days' mild fever, accompanied in several cases by herpes labialis, in others by desquamation of the cuticle, more especially of the fingers. There was no rash. All the recovered cases that were tested (over 20) had agglutinins in their serum, specific for the causative organism.

III.—PATHOLOGICAL ANATOMY.

Assisted by Drs. P. F. Graham (the medical attendant to the institution) and Dr. J. F. Shanahan, I performed, on November 6th, the autopsy of 3 of the cases—namely, S. K., aged 13, the first to die; L. O'D., aged 12½, the only second-class girl who succumbed; and M. K., aged 13, the last to die. The notes were kindly taken by Captain Elsner, R.A.M.C. Eliminating all non-essential details, the following were the chief findings:

Alimentary Canal.—Empty throughout, save for a little creamy fluid; not inflamed, nor ulcerated, nor even hyperaemic, save for a few streaks of capillary engorgement on the gastric mucosa. The mucous membrane of stomach and intestines presented a peculiar pale, swollen, velvety aspect, with here and there patches of white, prominent granules, each as large as a pin's head, projecting under the mucous membrane. They consisted of nodules of tumid lymphoid tissue. Peyer's patches were not tumefied.

The *liver* showed characteristic though not exactly obvious changes in all 3 cases. It was large, soft, hyperaemic, and sprinkled throughout with little, indistinct, pale-yellowish areas or spots. Microscopic examination of the fixed and hardened preparation revealed the most intense fatty degeneration of the liver cells at these places. In one case (L. O'D.) there were areas of haemorrhagic breakdown as big as a pea. The spleen showed no constant change. The kidneys were in a state of cloudy swelling.

The *cardio-vascular system* showed but little change. Endocardium and aortic intima not stained. The volume of the blood was much diminished and the clots were tarry in character. The *lungs* showed only some scattered patches of hyperaemia. The anatomical diagnosis was *cholera nostras* due to food poisoning.

IV.—BACTERIOLOGY.

From practically all of the viscera examined (for details, I would refer to the complete paper), as well as from the discharges obtained from the recovering cases, I succeeded in isolating, either by direct culture or after preliminary enrichment in malachite-green broth, a typical strain of the *Bacillus enteritidis* of Gaertner. The medium used for plating out was Drigalski's litmus-nutrose-lactose-crystal-violet-agar. On it the Gaertner colonies were readily distinguishable by their bluish translucent appearance, vigorous growth, and marked tendency to become confluent. The same colonies were obtained from all three cases examined *post mortem*. Avoiding technical details, I will merely say that morphologically this organism is identical with the typhoid bacillus, and is quite as actively motile. It differs from the typhoid bacillus *inter alia*, in its more rapid growth on gelatine, its ability to clear up milk, and to ferment glucose with formation of acid and gas. It differs from an actively motile strain of *B. coli* in the delicate character of its gelatine colonies, its inability to coagulate milk, to ferment lactose, and to form indol. I could find no constant morphological or cultural difference between this strain of Gaertner's bacillus and the paratyphoid *B. bacillus* (two strains isolated by myself from Irish cases). Neither does it seem to differ morphologically or culturally from the bacillus of Ratn.

Serologically, however, it stands nearer to typhoid than to paratyphoid *B.*, and by comparative agglutination tests I find it to belong to the true Gaertner type (that responsible for the outbreaks of food poisoning at Rumfleth, Haustadt, Morseele, etc., described by van Ermengem and his pupils) and not to the morphologically identical, but serologically distinct, "Flügge-Kaensche" or "Aertryke" type.

I reserve details as to its behaviour on the various coloured and other test media for the complete paper. As regards its pathogenicity for animals, the Limerick strain is very virulent for rats and mice, setting up in mice a fatal epizootic disorder accompanied by yellowish mucoid