induction of anaesthesia in a child's iron cot with high barred sides is very difficult, and the child is most apt afterwards to associate its experiences with its own sleeping bed instead of with a table which has then been taken quite away.

The mothers of these children are frequently absurdly lacrymose, and unnerve the child by their presence at the beginning of affairs. The surgical nurse or doctor should therefore take them into the room and the mother should go downstairs until called for after all has been completed.

Sebaceous Cysts.

These cannot be completely removed under nitrous oxide gas owing to venous oozing obscuring the cyst wall, and the C_1E_2 mixture is the most satisfactory anaesthetic for the patient and operator. These cysts are not very sensitive unless inflamed, and there is no need for a very deep anaesthesia. Their position on the scalp is often somewhat inaccessible when the patient is lying down, and a table should be used instead of a bed, so that the head may be properly reached from its upper end. If the patient has to be laid partly on his face he should be anaesthetized lying on his side and then rolled into position, care being taken that the chest walls can freely expand in respiration.

Dislocations.

Patients meeting with an accident causing dislocation of a limb are unprepared by abstinence from food for an anaesthetic, and are not likely to take it well. A jaw gag must be at hand in case of semi-solid vomiting, and, whatever anaesthetic is used to make the patient unconscious, ether should afterwards be given to obtain muscular relaxation. A great many deaths have occurred under chloroform unwisely pushed for this purpose. C_1E_2 followed by ether or gas and ether are the most suitable and safe anaesthetics to employ.

Diagnostic Examinations.

Chloroform may be wisely administered for examinations, except in rectal cases, where dilatation of the sphincter and causes reflex laryngeal spasm and ether should be used.

As chloroform requires some days for its complete elimination from the blood, it is wise not to employ it for a lengthened examination the day before a contemplated operation. After an interval of four days less after-effects result from the second anaesthetic.

Just as much careful watching of the patient must be practised as during a surgical operation; but two great factors, namely, the inhibitory reflex effects produced by surgical stimuli and haemorrhage, are absent.

Parturition.

One observation only is needed in conclusion. It should not be forgotten that all the conditions during confinements are extremely favourable for the administration of chloroform—the lateral posture, the higher blood pressure, the deeper respiration, all assist materially in the maintenance of circulatory equilibrium. The absence of difficulties and depressing effects from chloroform at confinements must not be looked for in other patients during other operations.

RESECTION OF THE TRACHEA FOR CICATRICAL STENOSIS.*

By G. GREY TURNER, F.R.C.S.,

TANT S RGEON, ROYAL VICTORIA INFIRMARY, NEWCASTLE-UPON-TYNE.

So far as I can gather, it seems usual in cases such as that about to be described to attempt to dilate either through the larynx or a tracheotomy opening, but these methods are exceedingly tedious and admittedly unsatisfactory. In a *System of Practical Surgery*, by Professor E. von Bergmann, vol. xi, p. 220, it is stated that "extensive circular scar stenoses or complete occlusion form an indication, according to the experience of Kuster

and v. Eiselsberg, for circular resection with subsequent suture of the trachea." In my case the trachea rapidly became obliterated by a considerable mass of new tissue, and it seemed to me in accordance with surgical principles to excise the involved segment when dealing with a non-collapsible tube of large calibre like the trachea. The occurrence of so serious a complication as secondary haemorrhage may appear as a fitting rebuke for adopting an audacious proceeding, but it appeals to me rather as a warning against the close suture of the external wound in cases where the area has become septic from the constant presence of a tracheotomy opening.

The patient was a man of 25, who inflicted the injury in a fit of depression following alcoholic excesses.

He was admitted to the infirmary on April 16th, 1908, in such a collapsed condition that his recovery was despaired of. There was a fearful gash in the neck almost completely severing the trachea, though the great vessels had escaped.

Next morning he was so much better that chloroform was administered and the injured parts thoroughly examined by the house-surgeon. It was then found that the trachea had been divided just below the cricoid cartilage, and that the sterno-mastoid and infra-hyoid muscles had been partially divided. The wound was thoroughly cleansed and the trachea drawn up to the cricoid with three catgut sutures, the muscles being separately repaired with catgut. Tracheotomy was not considered necessary, but the superficial wound was left partly open.

From the time the patient first began to talk the voice was characteristic of recurrent laryngeal paralysis, and this condition never altered. Unfortunately suppuration occurred, and the wound only healed slowly by granulation. The sutures holding up the trachea gave way, and it was found necessary to introduce a tracheotomy tube. As cicatrization proceeded all attempts to dispense with the tube gave rise to much distress, and it was obvious that the trachea was becoming stenosed.

On May 15th an anaesthetic was again administered, and just above the tracheotomy opening the trachea was found so much stenosed that it would only admit a fine probe through a channel at the summit of a funnel-shaped narrowing. It was found possible with a pair of sinus forceps, followed by artery forceps, to easily dilate the stricture, and an attempt was made to keep it patent by a silver tracheotomy tube passed upwards into the larynx through the top of a large rubber tracheotomy tube introduced in the usual way (in imitation of Störk's tracheal cannula and dilator). As might have been expected, this plan was not successful, and, in spite of all our efforts to keep the trachea open, it became necessary to interfere again, or let the case drift into one of permanent stenosis with its necessary tracheotomy.

On June 12th the last operation was performed, again under general anaesthesia. This time complete obliteration was found, the lumen of the trachea being filled by a mass of scar tissue; it was therefore decided to resect the involved portion. The cricoid and the upper part of the windpipe were accordingly thoroughly exposed through a vertical incision, and a complete segment of the trachea excised. The resulting gap was very wide, but the trachea was freed without fear until it could be easily approximated to the cricoid, to which it was fixed by several No. 1 chromic catgut stitches passed around the upper ring of the trachea and through the cricoid, care being taken to tie the knots on the outside. The posterior and lateral surfaces were brought into perfect apposition, but there was a little irregularity of the union in front. A vertical opening was made into the trachea about $\frac{1}{2}$ in. below the line of suture for the introduction of a small silver tracheotomy tube. The skin incision was then carefully closed by silk worm stitches. The part removed consisted of a complete transverse section of the trachea with two rings, the lumen being completely obliterated by a mass of scar tissue.

For five days the patient went on very well, but there was then a little suppuration with rise of temperature, and the skin wound gave way. On the eighth day, without any warning, he had a sudden profuse haemorrhage. The bleeding clearly came from the left side of the wound from a recess by the side of the trachea, and, after resisting all the efforts of the house-surgeon, it stopped spontaneously. As the source of the haemorrhage was doubtful, and its spontaneous arrest probably only temporary, I thought it wiser to give an anaesthetic, and, if possible, deal directly with the bleeding point. As a matter of fact, I was only able to curette a small cavity, which I packed with gauze soaked in turpentine. As one or two of the sutures in the front part of the tracheal wound had given way, two silk worm stitches were introduced and subsequently removed.

After this the patient made a straightforward recovery and was soon able to leave the hospital, the wound healing completely in the sixth week following the resection operation. When shown at the meeting in December, 1908, he was very well and had no difficulty in breathing except on severe exertion or when suffering from cold. The scars in the neck were not very obvious, but on palpation the trachea was found to be a little narrowed from before backwards and this was well shown on x-ray examination.

Dr. S. S. Whillis kindly examined the larynx and reported paralysis of the left vocal cord, general hyperaemia of the laryngeal and tracheal mucous membranes, but no evidence of stenosis as far as could be seen.

* The case that forms the subject of this note was shown at a meeting of the Newcastle-upon-Tyne Clinical Society.