die from loss of blood, but rather from an acute septicaemia, with high temperature and a steadily-increasing pulse. The epistaxis could always be controlled by plugging.

7. The presence of a purpuric eruption on the body, as virtually the only sign for several weeks.

8. As is usual in the acute lymphatic cases, the white cells are almost exclusively large lymphocytes.

We thus see that acute leukaemia is generally of the lymphatic type, occurs in young subjects, is generally fatal in from two to four months, and is uninfluenced by treatment.

NOTES ON THE MICROSCOPICAL EXAMINATION OF BONE MARROW.

(METHOD AND CLASSIFICATION.)¹

By CECIL PRICE JONES, M.B.LOND.,
Pathologist, East London Hospital for Children.

[From the Bacteriological Laboratories, Guy's Hospital.]

THE interest attaching to the red bone marrow as a probable factor in the causation of the so-called blood diseases, and as an agent controlling the course of many other diseases not specially regarded as blood diseases, has created a demand for a more thorough knowledge of the sources and causes of variation of the different blood cells, and for a more systematic examination and study of the cytology of the red bone marrow.

The subject is complicated by several technical difficulties arising from the readiness with which the marrow cells undergo *post-mortem* changes; from the unsatisfactory results obtained by the smear preparation and sections now generally employed, whereby the true appearance of the cells is most liable to be destroyed or masked by artifacts; and also from the absence of a suitable scheme for classifying the varieties of cells. The method of preparing and the system of classification to be described in the following notes, and which have been adopted in the examination of a very large number of specimens of bone marrow obtained from various sources, including besides those taken at human *post-mortem* examinations, those prepared from freshly killed dogs, cats, kittens, rats, rabbits, guinea-pigs, mice, chickens, pigeons, frogs, etc., in both adult and fetal states, have been found, in great measure, to overcome these difficulties and to establish the study of the bone marrow on a basis comparable with that of the cytology of the blood.

METHOD.

Marrow is obtained as soon as possible after death, from a rib or vertebra, by squeezing it out of the bone by means of strong forceps or scissors; in the animals above mentioned it was obtained from the femur. It is conveniently collected on a platinum loop, and transferred at once to a watch-glass containing the dissociating reagent. There results a more or less complete emulsion, and the cellular elements of the marrow are dissociated.

To obtain this result repeated trials were made with a number of different solutions, including distilled water, salt solution, various strengths of alcohol, formalin, Eau de Javelle, solutions of urea, etc., but owing either to their insufficient dissociating action or to the production of chemical changes that interfered with the subsequent fixing and staining of the cells, these were all discarded in favour of a 10 per cent. solution of neutral glycerine.

This dissociating reagent is prepared by diluting glycerine with ammonia-free distilled water to form a 10 per cent. solution, and titrating this against decinormal sodium hydrate, using phenolphthalein as an indicator. The initial reaction of this solution varies from +0.1 to +0.5 (Eyre's scale). This reagent has a specific gravity of 1.029 at 15.7° C.

A loopful of 10 per cent. glycerine is then placed on a clean cover slip, and to this a loopful of the marrow emulsion is added and very gently spread over the surface of the slip. The film thus prepared is allowed to dry in the air, without heating, until a uniform ground-glass appearance is produced; this usually results within five minutes, unless the film is too thick. The film is then fixed and stained with a combined eosin and methylene-blue solution (Jenner's blood stain), for two and a half minutes, washed with ammonia-free distilled water, dried thoroughly and mounted in xylol balsam. It is in fact treated exactly as a blood film.

On microscopical examination the various cells should be seen well separated from each other, so that they can be individually studied and classified. A field showing eight to

ten cells is to be aimed for: experience soon enables one to judge on this point. Care must be taken to avoid making the emulsion too concentrated or the films too thick, lest the cells are crowded together and the object of the film be defeated.

CLASSIFICATION.

Red bone marrow consists of connective tissue, fat, blood vessels, lymphatics, nerves, and particular cell elements characterized by a peculiar transitional nature and a tendency to produce atypical forms. This atypical transitional constitution is the great obstacle to the satisfactory classification of the marrow cells, since it is frequently very difficult to decide into what class the various intermediate cell forms should be placed.

In film preparations the cells to be classified are the myelocytes or marrow cells proper, and those cells that belong to the blood. There are two chief divisions into red cells and white cells.

Red Cells.

The red cells comprise (a) the non-nucleated red cells of the blood, (b) the nucleated red cells of the marrow. The varieties of these cells will not be discussed in these notes, but it may be mentioned that the nucleated red cells in normal marrows number about 10 to every 100 of the white cells, though this proportion is often considerably increased in diseased conditions.

White Cells.

The white cells are classed into two divisions, the non-granular and the granular.

The non-granular group comprises:

- (a) Giant or mother cells.
- (b) Lymphoid or primitive cells.

The granular group comprises:

- (c) Coarse oxyphile cells.
- (d) Fine oxyphile or neutrophile cells.
- (e) Amphiphile cells.
- (f) Coarse basophile cells.
- (g) Fine basophile cells.
- (h) Polymorphonuclear leucocytes.

Without entering upon the question of the functional and developmental nature of these cells it is desirable to shortly describe the appearances relied upon for purposes of recognition:

(a) *The giant or mother cells* have the appearance of zoogloic masses with rounded, irregular, bulging margins. They vary in diameter from 20 to 50 μ . The protoplasm appears to be indistinctly granular and stains faintly neutrophile. In the depth of the cell are three to five or more nuclear bodies about 4 μ in diameter, staining faintly basophile, and containing one or two small deeper staining nucleolar-like bodies. These cells constitute about 1 per cent. of all the white cells.

(b) *The lymphoid or primitive cells* exhibit many varieties of transitional forms. Commonly they closely resemble the above-mentioned nuclear bodies of the giant cells, being round, homogeneous masses about the size of a red-blood cell, staining faintly basophile, showing no distinction of nucleus and surrounding cell substance, and containing one or two nucleolar-like bodies that stain more deeply basophile. Transitional forms extend in several directions. On the one hand, they present all grades of cells, from those showing a scarcely perceptible rim of differentiated cell substance to the fully developed cell with deep-stained margin of hyaline protoplasm, indistinguishable from the lymphocyte found in the blood. On the other hand, this early differentiating rim of cell substance takes a faintly oxyphile stain, and all grades can be observed between this and cells that are not to be distinguished from the nucleated red cells. A third series shows oxyphile granules in the fine rim of early differentiated cell substance and all the subsequent grades from this to the fully-developed granular cell. This lymphoid or primitive cell is the most important of the white cells, and constitutes about 50 to 60 per cent. of all the myelocytes. An examination of marrow taken from a new-born infant shows over 70 per cent. of these cells, and in fetal marrow the proportion is very much greater.

(c) *The coarse oxyphile granular cells* have pale basophile staining nuclei resembling the primitive cell, surrounded by a varying amount of reticular cell substance in the meshes of which are coarse oxyphile staining granules. Apart from variations in size of the nucleus and in the number of the granules, these cells are the most constant and typical cells found in the marrow. They constitute about 5 per cent. of all the white cells. Clinically they are known as eosinophile myelocytes. It may be noted that these cells appear very

early in embryonic life; and they were found to be very constantly present in the marrow of all the various animals referred to above.

(d) *The fine oxyphile or neutrophile granular cells* differ from the preceding by the much greater size of their nuclei, which are often notched or reniform or tending to divide, and by the much smaller size of their granules. These granules vary extremely in their number; occasionally no granules can be detected, the reticular protoplasm of the cell substance is stained faintly neutrophile, and the cell is only recognizable by its general similarity in size and nuclear characters to those cells that possess granules. This variety constitutes about 30 to 40 per cent. of all the white cells. It is the "myelocyte" of the clinicians. It is regarded as the prototype of the polymorphonuclear leucocyte of the blood. In man the granules take a neutrophile stain.

(e) *The amphiphile granular cells* contain two kinds of granules; some that stain oxyphile and others that stain basophile. These granules are coarser than those of the fine oxyphile cells, but finer than those of the coarse granular cells. This variety has not been met with in human marrow examined by this method.

(f) *The coarse basophile cells* resemble the coarse oxyphile cells, but the granules, which vary greatly in size in the same cell, take a basophile stain. This variety has not been met with in human marrow examined by this method.

(g) *The fine basophile granular cells* resemble the coarsely-granular oxyphile cells in the character of their nuclei, but the granules are of the fine variety, and stain deeply basophile. This variety constitutes about 1.5 per cent. of all the white cells.

(h) *The polymorphonuclear leucocytes* are mostly brought to the marrow by the blood, and should not be regarded as true myelocytes. They comprise the neutrophile and eosinophile leucocytes of the blood, and need no description here.

It is to be noted that the large mononuclear hyaline cells of the blood have never been observed in any specimen of marrow examined by this method. The same remark also applies to the mast cell.

Results of Differential Counts.

The subjoined table exhibits examples of differential percentage counts of the cells of red bone marrow, prepared and classified according to the method that has been detailed above.

Source.	Giant Cells.	Lymphoid Cells.	Coarse Oxyphile Cells.	Fine Oxyphile or Neutrophile Cells.	Amphiphile Cells.	Coarse Basophile Cells.	Fine Basophile Cells.	Polymorphonuclear Cells.	Nucleated Red Cells.
Man,* healthy ...	0.2	50.2	3.3	39.3	0	0	0.2	6.7	8.2
Newborn infant ..	1.5	70.7	6.7	10.2	0	0	0	10.7	57.25
Rabbit, healthy ...	0.3	55.6	0	32.8	0.9	0.6	2.1	7.2	10.0
Dog, healthy ...	0.16	45.8	2.0	22.6	0.16	0.16	0.66	29.9	22.0

* This man died from internal hæmorrhage, the result of a carriage accident.

NOTE.

A short description of the above method was given, and microscopical specimens were shown, at a laboratory meeting of the Pathological Society of London, held at Guy's Hospital, in May, 1904.

CLINICAL METHODS OF ENUMERATING LEUCOCYTES.

By EDWARD TURTON, M.D., B.Sc., M.R.C.P.,
Hull.

THE methods of estimating the number of leucocytes in the blood, which are described in nearly all the medical textbooks and even in the excellent treatises on the blood, are generally very lengthy and tedious, and in most cases involve the use of complicated calculations. Now that the custom of making careful blood counts is becoming more common I am convinced, after a lengthy study of the procedures usually described, that some of the simpler methods which are available for quickly and easily performing a leucocyte count are

deserving of more universal practice. Rapidity, simplicity, absence of involved calculation, combined with sufficient accuracy, are the essentials of good bedside clinical examinations.

The first practical point to consider is, whether it is advisable to count the red corpuscles and leucocytes in the same preparation, or to make two slides and count the two varieties of cells separately. Haematologists seem about equally divided on the subject. It is certainly a great convenience, as well as a great saving of time, to require only one drop of blood, and to have to clean only one mixing pipette and slide. But with blood in which the ratio of red cells to leucocytes approaches to 500 to 1 or so, it is almost impossible by using one slide to avoid overcrowding of red cells on the one hand or a big margin of error, due to too few leucocytes, on the other. Cabot¹ says there should be at least a hundred leucocytes actually counted, to use as a basis for calculating the number present in unit volume of blood. Employing the method I shall describe later, I prefer to count the two varieties of cells in different slides, using the "white" Thoma-Zeiss pipette for the leucocytes with a dilution of 1 in 20, and the other pipette for the red cells with a dilution of 1 in 200. In any case where the leucocytes are increased to 25,000 or more per cubic millimetre, and especially when the red cells are also reduced, it is easy to count both varieties of cell in one preparation, using a dilution of 1 in 100. Sherrington² in his investigations used a dilution of 1 in 50, and emphasizes the importance of counting the two kinds of cells together.

With regard to the diluting fluid, when counting both red cells and leucocytes on one slide, a 3 per cent. solution of common salt deeply coloured with gentian violet is usually quite sufficient, though Toisson's or Sherrington's fluids may be used. The stained nuclei of the leucocytes render counting easy, and help is given by the suggestion of Gowers³ of raising the objective slightly out of focus when the white cells are distinguished by their greater refracting power.

When counting leucocytes alone, with the "white" pipette, I use a fluid containing 1 c.cm. of glacial acetic acid, or 3 c.cm. of the acidum acetum B.P. in 100 c.cm. of water, which gives a solution containing 0.99 per cent. of hydrogen acetate. This is stronger than Thoma's fluid, which is 0.3 to 0.5 per cent., but it serves to dissolve the red cells in half a minute, and leaves the leucocytes deeply stained by the gentian violet. Stronger solutions are liable to affect the white as well as the red cells. It is possible with this fluid to distinguish roughly the relative proportions of mononuclear and polynuclear cells, though for a proper differential count fixed and stained blood slides only are of any use. I have been unable to obtain properly stained specimens using Ehrlich's triacid or Leishman's Romanowski stain with fresh blood films, nor do I consider the enumeration of fresh eosinophile cells as done by Zappert⁴ or Elzholz⁵ a satisfactory clinical procedure.

The Thoma-Abbé Zeiss instrument, or one of the many modifications of it, as those of Zappert and Türck, is that which is now almost universally employed in this country as well as in America and on the Continent, and I shall in consequence confine my remarks to this apparatus. The procedures adopted in counting leucocytes may be divided into two classes—(1) the method of "squares," and (2) the method of "fields."

1. The Method of "Squares."

On the Thoma-Zeiss instrument, as is well known, an area of 1 sq. mm. is divided into 400 small squares, each with a side of $\frac{1}{20}$ mm. and an area of $\frac{1}{400}$ sq. mm. These small cells are arranged by additional lines into 16 groups each of 16 small cells. By means of a cover glass a chamber of $\frac{1}{10}$ mm. is formed. The cubic content of the space above each square is thus $\frac{1}{4000}$ of a c.mm. If, then, the leucocytes be counted in a number of cells and the average found, it is necessary to multiply the number thus obtained by 4,000 and by the dilution (whatever this may be); this gives the total number per c.mm. To facilitate matters it is better to count the cells in groups of sixteen squares, but even so this is a long and tedious method, and involves some eyestrain.

In the Prague clinic it is customary to count the leucocytes in the whole 400 squares; this, though taking a good deal of time, involves no calculation, as with a 1 in 10 dilution you multiply by 100 by simply adding two terminal cyphers to the number obtained.

Ewing⁶ counts both the red and white corpuscles in one specimen, using Toisson's fluid with a dilution of 1 in 100 and a modified Zappert chamber, which is ruled so that the cells in an area of 9 sq. mm. can be counted.

Preitetschensky⁷ describes a modified Zeiss slide, on which