may be seen in Fig. 3. Hyperplasia of the connective-tissue elements and of the submucous glandular structure ensues. Increased functional activity at first accompanies the latter, and is evidenced by a greater flow of mucus on to the surface of the cords, and by more frequent efforts to clear the voice. The hyperplasia may in itself become a cause for

Fig. 3.—Photomicrograph of vertical section through the posterior end of a vocal cord, showing proliferation of the blood vessels and hyperplasia of the subepithelial connective tissue.

hyperaemia, tissues having some power of regulating their blood supply. The formation of new connective tissue is progressive, and the strands of fibrous tissue interlacing in amongst the glandular structure compress and gradually

Fig. 4.—Photomicrograph of vertical section through a vocal cord, showing the increase of fibrous tissue compressing and obliterating the glandular structure.

obliterate it (Fig. 4). In consequence of this the supply of mucus to the cords is diminished. The muscle fibres themselves become surrounded by connective tissue, isolated areas are cut off and degenerate, a greater effort is required to produce the singing voice, and this increased effort in itself tends to renew the progressive nature of the morbid process (Fig. 5). The epithelium undergoes, *pari passu*, a hyperplasia, in time followed by a metaplasia, with, perhaps, the development of warty excrescences about the vocal processes and a diffuse pachydermia laryngis.

In even quite the earliest stages of the morbid process I have described, the singing function must become seriously impaired. The intrinsic muscles, highly differentiated and specially trained, only too readily forget their office. The changes in the epithelium materially affect the vibratory power of the cords and the quality of the notes produced.

The treatment indicated by a study of the pathogenesis is rest—absolute rest. Local applications, local remedies of all kinds, can only aggravate and irritate the processes it is sought to arrest. As regards prevention, the natural conclusion, and the one of first importance, is to ascertain the capabilities of any individual vocal organ, and then to develop

Fig. 5.—Photomicrograph of vertical section, showing hyperplasia of the epithelium and papilliform digitations. The strands of fibrous tissue in the subepithelial layer are extending amongst the intrinsic muscles of the vocal cords.

rather than to train them, and above all things to let the voice be natural.

In conclusion, it affords me pleasure to express my gratitude to the Council of the British Medical Association, under whose auspices my investigations into the pathology of the larynx have been conducted.

PRELIMINARY NOTE ON RENAL ACTIVITY DURING ANAESTHESIA.

By W. H. THOMPSON, M.D., Sc.D.,

Professor of Institutes of Medicine, Trinity College, Dublin.

RECENT observations which I have carried out in this College have disclosed an unexpected effect on the secretion of urine produced by ether and chloroform when inhaled for the purpose of inducing anaesthesia.

Two varieties of experiments on this question had been in progress for some time past. In the first the urine was collected by means of cannulae placed in the ureters, the animals being narcotized with morphine, supplemented, when necessary, by a short inhalation of a mixture of ether and chloroform. In the second class the urine was drawn off from the bladder of the animal (female) by catheter, no previous narcotic being required.

In both sets of experiments observations were made for the following periods, the duration of which varied from one to two hours:

- (a) Normal secretion before administration of the anaesthetic;
- (b) During anaesthesia; and
- (c) For one or more periods afterwards.

Two opposite effects were noted in the case of each anaesthetic. During the administration, some of the experiments showed a pronounced increase in the secretion of urine, others a marked depression. Both types of influence were exemplified in each class of experiment, but the increase effect was more frequent and pronounced in the ureter experiments, the decrease in the catheter experiments.

The following are examples of both kinds of result, the figures representing cubic centimetres of urine per period. During period (b) the anaesthetic was given:

I.—Ureter Experiments.			
(a)	(b)	(c)	(d)
12.6	25.0	32.75	24.0 (chloroform)
6.7	31.7	31.7	24.1 (ether)
II.—Catheter Experiments.			
23.5	3.6	40.0	114.0 (chloroform)
38.0	1.0	13.5	48.5 (ether)

Still more recent experiments have shown how these two opposite sets of results may be reconciled. From these it appears that the early effect of both anaesthetics, lasting up to the period of complete insensibility, is ordinarily an increase of the secretion of urine. This, as a rule, is more pronounced in the case of ether than of chloroform. But during full anaesthesia, however prolonged—and this is the remarkable result—the secretion of urine is almost completely arrested. By full anaesthesia I mean a degree just beyond the loss of conjunctival reflex—namely, that required in surgical operations. The depressing effect during this period appears likewise to be more pronounced with ether than with chloroform.

Upon removal of the anaesthetic the kidneys rapidly recover, and, at all events in dogs, are actively secreting once more within five or ten minutes, this activity passing on to a condition of hyperactivity which lasts for some hours.

The value of the foregoing results (obtained with dogs) has been greatly enhanced by corroboration from the clinical side. Observations on renal activity during surgical anaesthesia, kindly carried out in Mercer's Hospital by Mr. Maunsell and Mr. Seton Pringle, together with Dr. Harold Pringle, Senior Demonstrator of Physiology in this department, have yielded results in the case of ether entirely in accord with my own. These will be published *in extenso* later, but meanwhile I am permitted to say that with the onset of insensibility in ether narcosis a condition of depression of urinary secretion sets in and continues which amounts in many cases to almost complete suppression. This is preceded in some instances by a marked increase of secretion during the induction of the anaesthesia. The following are the figures from one observation:

Rates of Secretion of Urine.

- (a) For twenty-four hours ending at 9 o'clock on morning of operation=42 c.cm. per hour.
- (b) For two hours previous to ether inhalation=104 c.cm. per hour.
- (c) During the induction of anaesthesia (time 16 mins., urine 65 c.cm.)=210 c.cm. per hour.
- (d) During anaesthesia, first half-hour, 3.5 c.cm.; second half-hour, 1.7 c.cm.

It would seem, therefore, that this somewhat surprising effect is one of ordinary occurrence. That in the vast majority of cases it appears to do practically no harm is almost more remarkable than the fact itself. It is hardly necessary to say that our investigations extend also to alterations, both qualitative and quantitative, in the excretion of urinary constituents under anaesthesia. The full results will be published at an early period.

TRANSILLUMINATION OF THE ANTRUM OF HIGHMORE.

[WITH SPECIAL PLATE.]

By A. BROWN KELLY, M.D., D.Sc.,

Surgeon for Diseases of the Throat and Nose, Victoria Infirmary, Glasgow.

THE term "transillumination" (*Durchleuchtung*) was introduced by Czermak in 1858 to designate a method of examining the larynx devised by him. This is carried out in a darkened room by concentrating a light upon the front of the patient's neck, and observing in a laryngeal mirror the various parts of the larynx and trachea as they are successively transilluminated by the light applied to the corresponding region externally.

While experimenting with this method, Voltolini found that the bones of the face could be transilluminated if in a dark room a small incandescent lamp were held in the closed mouth. Although he failed to attach the proper significance to the phenomena he described, the credit of having first transilluminated the antrum (1888), and of having indicated that this procedure would be of value as a diagnostic test, belongs to him.

In the following year Heryng placed transillumination on a practical basis. He showed that when the light traverses a healthy antrum, part of the cheek and the lower eyelid become red (for convenience we shall term the bright area the *antral tache*), while in empyema these regions remain dark owing to the altered conditions in the cavity preventing the passage of light. On the other hand, he proved that when a serous cyst occupies the antrum transillumination is unaffected, while if it distend the cavity the area of brightness is increased.

Vohsen, in 1890, by using a stronger light, showed that the eye also may be transilluminated, the pupil then appearing as a red disc.

In 1892 the writer pointed out that during transillumination a person with closed eyes sees a dull red glow if the antra be normal, but if the *antral tache* be absent the subjective light sensation will also be absent.

Successful transillumination yields the three signs referred to—namely, the *antral tache*, the bright pupil, and the subjective light sensation. As to their importance the most diverse opinions have been expressed. I have, therefore, endeavoured to analyse the appearances observed during transillumination and to assign to these, individually and collectively, their proper value.

For the purposes of transillumination I have found a lamp, as depicted in Fig. 1, convenient. To the end of a suitably curved handle an incandescent lamp is fitted, over which various end pieces slip. One of these is scoop-shaped, and rests on the tongue during ordinary or palatal transillumination; another completely encloses the lamp, and has an opening on one side, which can be applied to the posterior surface of the superior maxilla inside the cheek for Escat's retro-maxillary transillumination; the third is funnel-shaped and open at the end for transillumination of the frontal sinus.

Fig. 1.

Transillumination should be carried out in a perfectly dark room. If the patient wears a dental plate, this must be removed. The lamp should be placed in the middle of the tongue and depressed sufficiently to keep it from burning the palate. The current should be gradually increased by means of a rheostat until the proper intensity is attained. Slight differences between the two sides may not be observed if too strong a light is used; while, on the other hand, a comparatively powerful light may be required to penetrate the tissues when thick.

TRANSILLUMINATION OF THE NORMAL ANTRUM.

Under ordinary circumstances the parts of the face that transilluminate are those around the lips, the cheeks, the bridge of the nose, the lower eyelids, the pupils, and the eyeballs to the inner side of the cornea. Only those bright areas or *taches* the intensity and extent of which may be influenced by the condition of the antrum will be considered.

Crescentic Tache.

It is very commonly supposed that in transillumination the light passes through the anterior wall of the antrum and produces redness of the corresponding part of the face. This is not what usually occurs.

Fig. 2.

In the great majority of persons it will be found that the illuminated area has a crescentic shape and lies above the lower margin of the bony orbit, which is indicated in Fig. 2 by a dotted line. This is at once evident if the test be practised on a cadaver with sunken eyeballs, when only the lower eyelids and tissues situated above the infraorbital margins are transilluminated. The light that produces the crescentic *tache* passes through the roof of the antrum; this may be demonstrated by drawing forward the lower eyelid, when the conjunctival sac is found to be rendered specially bright by the stream of light from below.

Infraorbital Tache.

In certain individuals, however, the light traverses the facial wall of the antrum and produces a bright area, which may be termed the *infraorbital tache*. The persons in whom this *tache* is observed are those that transilluminate well—for example, young adults, spare individuals, and those with good-sized, thin-walled antra. In short, to obtain the *infraorbital tache* the conditions for transillumination must be