noted that the vowel "Ah" must not be used until the voice is thoroughly under control. When this is so, there is no better vowel for practice, for it helps to give the open throat so very necessary for good voice production, but if used too early in the training, it tends to force the vocal cords apart.

No attention is paid to breathing, the whole of the mind is concentrated at first on getting a loud, sustained, metallic ringing tone. When the habit of contracting quickly and fully has so to speak been impressed upon the adductor muscles of the cords, gradations of tone can be carried out with safety and delightful ease, and it will be found that whatever amount of breath is required for phrasing, etc., can now be taken in and scientifically dealt with without having to worry about breath control. What is thought a normal breath now would not perhaps be considered one in, say, three months' time, for as control is obtained of the adductor muscles of the cords the muscles of expiration will also come under the influence of the will and inspiration unconsciously becomes deeper; but by no stretch of the imagination can this increased chest capacity be said to be abnormal. All the deep breathing exercises in the world cannot prevent the undue escape of breath during phonation, if the aperture of exit is too large, which is the case in ninety-nine out of every hundred persons. Unconsciously, from the first the pupil has been learning abdominal breathing, as it is impossible to say "Pat-a-wat" without an abdominal upheaval.

Now on attempting to sing with a good tone, the utmost use must be made of the mouth and pharynx to add all the resonance possible. The *voce di petto* of the Italians is simply and solely the result of duly approximated cords and an open pharynx and mouth. The chest plays only a very minor part in its production; in fact if any attempt is made to add resonance by filling it out unduly, it will be found most difficult to control this extra amount of air. Moreover, it is hard to understand how a chest full of compressed air can make any difference to its resounding qualities, and under the circumstances singing becomes more a feat of strength than a natural and pleasurable exercise.

No reference has been made to such terms as registers, chest note, head note, forward tone, etc., for in the properly-produced voice there is only one register or mechanism, and as long as the mind is capable of conceiving the pitch of a note, the cords will relax or stretch to produce it; but the size of the aperture between the cords remains the same throughout the whole of the compass. Every vibration partakes of the whole resonance of the cavities into which it can enter, and there is only one place of origin and emission of both the highest and lowest notes whatever the sensation may be.

A system of instruction such as the one sketched is actually in existence, and it has produced and is still producing amazing results in an incredibly short time.

But, to be a successful singer or speaker the possession of a voice is not enough; mental ability of an artistic kind is also needed, with, of course, a thorough knowledge of elocution and music.

SOME OBSERVATIONS ON THE PATHOGENESIS AND PREVENTION OF LOSS OF VOICE IN SINGERS.

By JOBSON HORNE, M.D., B.C.CANTAB., M.R.C.P.LOND.,
Surgeon, Metropolitan Ear, Nose, and Throat Hospital; late Ernest Hart
Memorial Research Scholar of the British Medical Association.

THE often-heard lament of the present age over its poverty in fine voices, and the want of lasting power in singing voices generally, has led to various explanations and to the formulation of extravagant theories. Some have attributed the decline in singing to the rise in pitch, which has been gradual, yet very considerable, during the past century; others have held composers responsible for not having sufficiently considered the voice, with the result that the vocalist has had to disregard his capabilities and put his vocal cords at everything—singable or unsingable. It is not the purpose of this paper to consider whether these explanations are or are not sufficient, or to offer any new theory in place of them; in passing, however, one might point out that neither the British language nor the British climate lend themselves so naturally to the old Italian method of voice production as a language teeming with vowels, and particularly with final vowels, and spoken in an atmosphere which is not conducive to chronic head colds.

The purpose of the paper is to summarize briefly the sequence of changes that are induced in the vocal cords by excessive or unnatural use, which may be represented in modern language by the word "strain"—an excellent term, unscientific, comprehensive, and truly descriptive of the "begin-all" and the "end-all" of faulty voice production.

The following facts have been arrived at partly through clinical observation and partly through *post-mortem* investigations, and it is hoped that the recording of them will serve further to emphasize the interdependence of art and science which has recently been brought under our notice.

The laryngoscope unaided affords practically no information about the finer movements and vibrations of the vocal cords, which are so essential to the production of the singing voice. To observe these it is necessary to supplement the laryngoscope with the stroboscope—an instrument known to scientists for nearly a century, and to children as a toy, in its modified form of the zoetrope, for almost the same length of time. To Dr. Musehold, of Berlin, we are indebted for knowledge about the singing voice that has been gained by the means of this instrument; in his clinic in Berlin, in 1897, I was afforded the opportunity of witnessing the more delicate functions of the cords, unobservable with the unaided laryngoscope.

Now to pass to the physiological changes which take place in the vocal cords during the production of the singing voice, changes which, when exaggerated, become the pathological factors to render the cords unfit for performing the finer movements required in singing. Of these changes hyperaemia is *facile princeps*.

Increased vascularity is the natural concomitant of increased function. The determination of blood to the vocal cords is in direct proportion to the effort made to produce the voice. It is possible to verify this by transillumination of the larynx. If in a darkened room an electric lamp—such as that used for illuminating the frontal sinus—be placed against the thyroid cartilage and the image of the cords be observed in a mirror in the pharynx, it will be noticed that the greater the effort used in the production of notes the more opaque the cords become; the increased opacity is due partly to increased muscular tension, but mainly to increased congestion of the vessels of the cords and to general vascularity. This hyperaemia may be further verified by observing the surface of the cords in the usual way. The increase both of muscular tension and of vascularity is illustrated in the accompanying photographs (Figs. 1 and 2).

Fig. 1.—Vocal cords in "falsetto register," *ad libit.*

Fig. 2.—Vocal cords in "chest register," *ad libit.*

taken by Dr. Musehold in 1887. The two photographs were given to me by Dr. Musehold to show the condition of the vocal cords during the production of the singing voice in the "falsetto" and "chest" registers respectively. One conclusion arrived at by Dr. Musehold from his investigations being that in the "falsetto" register the glottis is only widened and narrowed, whereas in the "chest" register the glottis is opened and closed, and the cords themselves become more rounded or sausage-shaped, in contrast with their condition in the "falsetto" register.

At the present time I am not concerned with registers, nor can Dr. Musehold be held responsible for my deductions from his physiological observations to explain my starting-point of a definite morbid process. I desire only to draw attention to the condition of the cords during the excessive or unnatural use of a register with a view of considering the pathological changes which result when the abuse is persisted in. The superficial vessels, owing to the determination of blood to the surface of the cords, become tortuous to a degree almost amounting to varicosity. In fact, submucous haemorrhages have been recorded as occurring after a straining of the singing voice. Excessive vascularity if often repeated acts as an irritant, and the capillaries and vascular elements proliferate, as

may be seen in Fig. 3. Hyperplasia of the connective-tissue elements and of the submucous glandular structure ensues. Increased functional activity at first accompanies the latter, and is evidenced by a greater flow of mucus on to the surface of the cords, and by more frequent efforts to clear the voice. The hyperplasia may in itself become a cause for

Fig. 3.—Photomicrograph of vertical section through the posterior end of a vocal cord, showing proliferation of the blood vessels and hyperplasia of the subepithelial connective tissue.

hyperaemia, tissues having some power of regulating their blood supply. The formation of new connective tissue is progressive, and the strands of fibrous tissue interlacing in amongst the glandular structure compress and gradually

Fig. 4.—Photomicrograph of vertical section through a vocal cord, showing the increase of fibrous tissue compressing and obliterating the glandular structure.

obliterate it (Fig. 4). In consequence of this the supply of mucus to the cords is diminished. The muscle fibres themselves become surrounded by connective tissue, isolated areas are cut off and degenerate, a greater effort is required to produce the singing voice, and this increased effort in itself tends to renew the progressive nature of the morbid process (Fig. 5). The epithelium undergoes, *pari passu*, a hyperplasia, in time followed by a metaplasia, with, perhaps, the development of warty excrescences about the vocal processes and a diffuse pachydermia laryngis.

In even quite the earliest stages of the morbid process I have described, the singing function must become seriously impaired. The intrinsic muscles, highly differentiated and specially trained, only too readily forget their office. The changes in the epithelium materially affect the vibratory power of the cords and the quality of the notes produced.

The treatment indicated by a study of the pathogenesis is rest—absolute rest. Local applications, local remedies of all kinds, can only aggravate and irritate the processes it is sought to arrest. As regards prevention, the natural conclusion, and the one of first importance, is to ascertain the capabilities of any individual vocal organ, and then to develop

Fig. 5.—Photomicrograph of vertical section, showing hyperplasia of the epithelium and papilliform digitations. The strands of fibrous tissue in the subepithelial layer are extending amongst the intrinsic muscles of the vocal cords.

rather than to train them, and above all things to let the voice be natural.

In conclusion, it affords me pleasure to express my gratitude to the Council of the British Medical Association, under whose auspices my investigations into the pathology of the larynx have been conducted.

PRELIMINARY NOTE ON RENAL ACTIVITY DURING ANAESTHESIA.

By W. H. THOMPSON, M.D., Sc.D.,

Professor of Institutes of Medicine, Trinity College, Dublin.

RECENT observations which I have carried out in this College have disclosed an unexpected effect on the secretion of urine produced by ether and chloroform when inhaled for the purpose of inducing anaesthesia.

Two varieties of experiments on this question had been in progress for some time past. In the first the urine was collected by means of cannulae placed in the ureters, the animals being narcotized with morphine, supplemented, when necessary, by a short inhalation of a mixture of ether and chloroform. In the second class the urine was drawn off from the bladder of the animal (female) by catheter, no previous narcotic being required.

In both sets of experiments observations were made for the following periods, the duration of which varied from one to two hours:

- Normal secretion before administration of the anaesthetic;
- During anaesthesia; and
- For one or more periods afterwards.

Two opposite effects were noted in the case of each anaesthetic. During the administration, some of the experiments showed a pronounced increase in the secretion of urine, others a marked depression. Both types of influence were exemplified in each class of experiment, but the increase effect was more frequent and pronounced in the ureter experiments, the decrease in the catheter experiments.

The following are examples of both kinds of result, the figures representing cubic centimetres of urine per period. During period (b) the anaesthetic was given:

I.—Ureter Experiments.				
(a)	(b)	(c)	(d)	
12.6	25.0	32.75	24.0	(chloroform)
6.7	31.7	31.7	24.1	(ether)
II.—Catheter Experiments.				
23.5	3.6	40.0	114.0	(chloroform)
38.0	1.0	13.5	48.5	(ether)