inflammation. Babies are born with adenoids, and there are small children who have never had a touch of acute tonsillitis whose tonsils yet meet in the middle line; neither are they to be called tumours in any strict sense of the word; neither are they evidence of any definite constitutional disease—such as rickets, or syphilis, or the predisposition to tuberculosis. We call them a local hypertrophy, but that name explains nothing. We know that the stores of lymphoid tissue throughout the body are proportionally greater in the child than in the adult; but that is a physiological affair, and enlarged tonsils and adenoids are pathological. Surely it is not unreasonable to suppose that the lymphoid structures in the pharynx are under the influence of that natural tendency toward defective development which we see in the teeth and in the upper air passages. Crowded, irregular, and carious teeth; crowded and irregular air passages lined with a mucous membrane which is liable to general atrophy at puberty; crowded stores of lymphoid tissue which is liable to an irregular sort of hypertrophy—surely these three conditions may have a common origin in a gradual change of type. It is certain that our teeth in the course of ages are become weak and ill-developed, and in a word, degenerate, and it may be that this degeneracy is at work also on the neighbouring and allied structures of the nose and the pharynx. For we know that the tonsils, like other organs which are in course of involution, are variable. Many people have no tonsils; that is to say, you can just discover, between the pillars of the fauces, a mere vestige of lymphoid tissue, which is, so far as we know, useless. Others have an excess of lymphoid tissue which is worse than useless. Surely this irregularity in size, and this apparent uselessness in function, suggest degeneracy.

THE OPENING OF PERITONSILLAR ABSCESES.

By STCLAIR THOMSON, M.D., F.R.C.P., F.R.C.S.,

Physician for Diseases of the Throat, King's College Hospital; formerly Physician to the Throat Hospital, Golden Square, and Surgeon to the Royal Ear Hospital.

THERE are few operations in the throat which are simpler, safer, and more satisfactory than the evacuation of a peritonsillar abscess, yet there appears to be considerable dissatisfaction both in the public and the profession in regard to "puncturing the tonsil," as it is termed. In his *Manual of Medical Treatment* Dr. Burney Yeo writes that "patients constantly complain of being uselessly put to pain by these punctures, which appear to give no relief."¹ Some medical men are equally frank in their confession of dissatisfaction. Thus, an American *confrère* writes: "I have seen a good many cases of quinsy, and hate the sight of them, because there seems so little that we can do for them" (P. F. Barbour)²; while another, in still more forcible and picturesque language, exclaims, "I would rather go against a penny-in-the-slot machine, or try to play for four aces than attempt to get pus out of one of these abscesses" (M. F. Coomes).³

The anxiety connected with attempts to relieve a quinsy is not limited to the fact that such efforts are frequently fruitless, but extends to a fear of plunging the bistoury into some large artery. Bosworth,⁴ in spite of his long experience, says that in no case has he incised a quinsy without a certain degree of nervousness on account of the proximity of the blood vessels.

The usual sequence of events in this disease is somewhat as follows: Attempts at relieving the pent-up pus are willingly postponed both by patient and practitioner as long as possible—by the former, to escape increased and possibly useless pain; and by the latter, to avoid adding to previous failures. Finally, towards the seventh day or later, the patient begins to get worn out, the medical attendant thinks he will surely find that "matter has formed," and hopes that he may be more fortunate this time in striking it. Armed with a narrow, sharp-pointed knife, and with a vivid mental picture before him of the proximity of the carotid artery, he has much difficulty in securing any but the most restricted view of the field of operation. If successful in seeing the tonsillar region he probably thinks it best to thrust the knife into the most inflamed and prominent part which presents, though, for choice, he prefers to insert it into the tonsil itself, as safer and offering more chance of reaching a "tonsillar abscess." If no pus appears, and the patient is sufficiently tolerant, he may make two or more stabs into the

tonsil, and the failure to find pus is glossed over by the encouraging statement that the bleeding will do good. This is a wiser comfort than the assertion that the abscess is not "ripe," for at the next visit one may be met with the disconcerting announcement that it "burst in the night."

It seems to me that the uncertainty of relieving a quinsy is due to a want of appreciation of its usual location, and that the anxiety connected with opening the purulent collection can be entirely avoided by not using a knife and by adopting a safe and equally effective instrument.

A quinsy, or an acute collection of pus in the faucial region, never forms in the tonsil itself. Although small, discrete foci of suppuration may occur within the lymphoid follicles of the tonsil, this takes place as part of the process of an ordinary tonsillitis, and does not give rise to the acute collection recognized clinically as "a quinsy." The latter is always peritonsillar, and forms in the areolar tissue between the tonsil and the pharyngeal aponeurosis. In very rare cases it collects externally to the tonsil. More frequently it is met with posteriorly—in the prolongation of the posterior faucial pillar. But in the large majority of cases—over 90 per cent., in fact—the abscess forms above and in front of the tonsil. It is only the latter condition I propose at present to consider.

Fig. 1.—Acute peritonsillar abscess in a female patient, aged 26. The tonsils had been incompletely removed at the age of 15, and since then the patient had suffered from repeated attacks of tonsillitis. These attacks had doubtless caused the adhesions between the tonsil stumps and the anterior faucial pillars, and these adhesions tended to cause and aggravate recurring attacks of quinsy. Both tonsils show lacunar tonsillitis. The inflammation of the right tonsil stump bulges forward the faucial pillar, to which it is adherent. Above this is the peritonsillar abscess, pushing the median raphe of the palate to the left and causing oedematous inflammation of the uvula. As the quinsy is situated above the tonsil it distends the soft palate, and brings it much nearer to the front teeth (compare Fig. 2). The abscess in this case burst spontaneously through the supratonsillar fossa.

The clinical symptoms of this affection are so well known that it is only necessary to point out that, although the tonsil itself is never the site of the abscess, it frequently shows some signs of lacunar tonsillitis, and may be raised from its bed by the underlying suppuration. This occurs in the region of the supratonsillar fossa, above and external to the tonsil, and between the posterior and anterior pillars of the fauces. The latter is bulged forward and upward, and distended over the swelling; while the posterior pillar, the margin of the soft palate, and the uvula are frequently acutely oedematous (Fig. 1).

Left to itself the peritonsillar abscess generally bursts spontaneously into the mouth. The site of rupture often escapes detection, as in a large number of instances—probably the majority—it takes place through the supratonsillar fossa, and the pus trickles down over or behind the tonsil. But sometimes a quinsy will rupture through the soft palate, and the opening is apt to pass unnoticed since it may only be large enough to admit a small probe and allow of a slow

oozing out of pus (Fig. 2). This is the route we should follow when relieving the pent up-pus.

To reach the abscess the soft palate itself must be traversed, and the "site of election" is a point which can readily be located by taking the following bearings: If an imaginary horizontal line is drawn across the base of the uvula, and another vertically along the anterior faucial pillar, they will intersect at a point overlying the supratonsillar fossa. Just external to this is the best point for opening a quinsy (Fig. 3). Another guide is that this point will be found on an imaginary line extending from the base of the uvula to the upper wisdom tooth on the same side.

To reach the abscess cavity it is sufficient to provide oneself with a pair of Lister's sinus forceps, or one similar to that figured on page 662. In most cases the patient can only open the mouth to a very limited extent, and inspection and precision are greatly facilitated by the use of a frontal mirror or searchlight. The field of operation is cleansed and sprayed

2.—Shows site of spontaneous opening of an acute peritonsillar abscess on the left side; about ten days after rupture. From a man, aged 30.

with a 5 per cent. solution of cocaine, and the patient's head steadied by an assistant. A general anaesthetic should never be given. If the surgeon is uncertain as to the presence or exact situation of the pus, it is easy to feel the region with the end of the forceps. When this encounters a resistant cartilage-like tissue he should "tap" onwards until the sensation of a boggy surface is reached—in the area above indicated. It is then sufficient to push the forceps backwards, with a slight inclination outwards, in order to reach the pus. In cases which have existed for a week or so the collection may be near the surface, but with a shorter history we may have to go to a depth of two or three centimetres. As the forceps are withdrawn the blades should be opened widely, as in Hilton's method, so as to produce a vertical opening through which the pus pours into the mouth.

There are several advantages in avoiding the use of a naked knife. The operator need have no anxiety as to a possible risk of wounding a large vessel, and the patient is much more likely to permit the proceeding. If a mistake in diagnosis should have been made, or the operator fail to open the abscess, the patient is not so apt to resent a manœuvre which, after all, he frequently regards as only a manipulation of examination.

As a matter of fact, I can only remember one instance in which I failed to open a quinsy, although I have now employed these forceps in many dozens of cases. The method has also the advantage of considerably curtailing the duration of the illness, as pus may frequently be found as early as the third day, and sometimes at the end of forty-eight hours from the commencement of the attack.

As the procedure is safe and simple, there need be no hesita-

tion in adopting it in an affection which is so painful and debilitating, with risks of pneumonia, albuminuria, haemorrhage, and even death from suffocation, if the abscess is left to open spontaneously.

There are only two other methods which appear to me to be equally safe. Thus, J. Killian reaches the abscess by introducing a curved, blunt probe through the supratonsillar fossa.⁵ I have not been successful in my trials with this plan, except in cases of some duration, and I do not find that patients readily lend themselves to it. The other method is that suggested by Moure, of cutting through, at the "site of election" already described, with a galvano-cautery knife.⁶ The principal objections to this method are that the treatment of quinsy is generally in the hands of the family physician, and the necessary armamentarium is not always at hand. Besides, patients are frequently so worn out and nervous, that the idea of treatment with "a battery" would be rejected in many cases. A great advantage of the plan I

Fig. 3.—Semi-schematic drawing to indicate the site for opening a peritonsillar abscess.

have described is that the instrument and preparations are so little alarming that the patient's consent to the exploration can generally be obtained.

REFERENCES.

- ¹ Vol. i, 1902, p. 17. ² *American Practitioner and News*, July 15th, 1902. ³ *Ibid.* ⁴ *Diseases of the Nose and Throat*, 3rd edition, 1897, p. 428. ⁵ *Munch. med. Woch.*, No. 31, August 5th, 1897, p. 866. ⁶ *Revue hebdomadaire de Laryngologie*, XXII, 11, 1901, No. 36, p. 273.

SOFT FIBROMA OF THE LARYNX AND NECK.

By F. T. PAUL, F.R.C.S.,
Liverpool.

IN THE BRITISH MEDICAL JOURNAL of January 7th, 1905, Sir Felix Semon relates the case of a patient suffering from the above rare condition. In his valuable comments on the case he states:

It is, in fact, the first case, so far as I know, in which a pure soft fibroma has ever been observed to have involved simultaneously the larynx and neck.

The specimen removed by Sir Felix Semon was sent to Mr. Shatcock for pathological examination. He considered it probably a neuro-fibroma, and states: "There is, as yet, no record of any similarly disposed laryngeal fibroma."

In 1893 I showed at the Liverpool Medical Institution a case pathologically similar; though being further advanced and more rapidly growing, it was not alike in clinical details. I described it then as an extrinsic spindle-celled sarcoma of the larynx, with marked fibrillation of the cells, and as "of