or a previous illness of some kind or other, is a bar to insurance, and it is noteworthy how almost invariably the reply is in the negative. My advice to the agent would be, if in doubt about having a case examined, "Do," and he will often be agreeably surprised.

REFERENCES.

¹ Dr. Pye-Smith, paper on Insurance, read before Alliance Insurance Company. ² BRITISH MEDICAL JOURNAL, 1907, vol. ii.

TWO CASES OF OSTEITIS DEFORMANS (PAGET'S DISEASE).

By R. MANWARING-WHITE, M.D. EDIN.,
HONORARY SURGEON, VICTORIA INFIRMARY, NORTHWICH.

OSTEITIS DEFORMANS is a sufficiently rare disease to make it worth while to report two cases in the same family which have come under my notice:

CASE I.

A man aged 56 first came under observation in April, 1907, complaining of headache and "rheumatic" pains in the limbs.

FAMILY HISTORY.

He had five children living; one died of convulsions and bronchitis in infancy. His father died at the age of 64 from a cause not known to the patient. His mother died of enlarged liver. The family history, so far as is ascertainable, is otherwise quite good, and presents no rheumatic tendencies. A brother who is suffering from the same disease in an incipient stage constitutes Case II.

PERSONAL HISTORY.

The patient, a timber sawyer by occupation, had not indulged to excess in tobacco or alcohol, and had never had syphilis or gonorrhoea. He had always enjoyed good health until the age of 40, that is, fifteen years earlier, when he had a sudden attack of acute pain in the left hip. Since then he has never been really free from pain, especially during the last ten years.

About nine years ago he was confined to bed for one month with pain in both legs; he then returned to his work, though he frequently had to sit down because of the pain. He stopped work about four years ago because of an accident, the fingers of his right hand being caught in the saw; eleven weeks elapsed before healing took place, and the first and second fingers have never been much use to him since, on account of chronic periostitis. About five or six years ago he first noticed the right leg just below the knee to be swollen, and three or four years ago he noticed the same in the left leg.

PRESENT STATE.

His general health is fairly good; he eats well and is quite intelligent, but at times very depressed. He always has pain in his legs and head. In the head the pain is usually "aching," mostly in the frontal region, but occasionally occipital. In the legs it is usually "aching" also, but at times acute, somewhat like cramp. Fleeting pains occur in other parts of the body; he cannot remain still for any length of time, as rest seems to increase the pains.

He maintains that he is worse on Sundays, as on that day he seldom goes out; on other days he takes walks in the morning and afternoon. His sleep is very disturbed, as he is frequently awakened by pains, which cause him to get out of bed and walk about the room. He has not slept for more than two hours continuously for some years. Usually he gets up at 5 a.m., as he says he cannot remain in bed any longer.

His gait is peculiar. The feet are placed far apart, the knees cannot be approximated, and he "waddles" along, but there is no real unsteadiness in the gait.

Osseous System.

His head is large; as a young man his hat was size 6½, now size 7½ is too small, and he finds difficulty in getting a hat to fit him.

Head.—The measurements are: occipito-mental, chin to the most distant part of occiput, 9 in.; occipito-frontal, point of union of the supraorbital ridges to the most distant part of occiput, 8½ in.; biparietal, 6½ in.; bitemporal, 5½ in.; glabella to occipital protuberance, 16 in.; circumference at a horizontal level immediately above the supraorbital ridges measures 24½ in.; junction of pinna and scalp over the vertex to a similar point on

Measurements of the patient's head compared to those of a man wearing a hat of size originally worn by the patient.

	Patient.	Comparative Case.
	Inches.	Inches.
Occipito-mental diameter...	9	9½
Occipito-frontal diameter...	8½	7½
Suboccipito-bregmatic diameter	7	7
Biparietal diameter...	6½	6
Bitemporal diameter	5½	5
Glabella to occipital protuberance	16	13½
Circumference	24½	22
Pinna to pinna	12½	12½

the opposite side, 12½ in.; suboccipito-bregmatic, from the neck immediately below the occipital protuberance to the junction of the parietals and frontal bones, 7 in.; suboccipito-frontal, from

the neck immediately below the occipital protuberance to the mid point of the junction of the frontals, 7½ in.

He has a peculiar visage, owing, first, to a slight degree of acne of the nose, and, secondly, to enlargement of the upper jaw, which is not symmetrical, being most marked on the left side; the left malar bone is also involved, giving the appearance at first sight of antral disease.

The Spine.—The curves are those of the normal spine, but greatly exaggerated. This appears to be due in part to the excessive weight of the head; the neck

is very short and thick, and there is a kyphosis of the dorsal region, being apparently compensatory to the cervical lordosis. The vertebrae themselves do not appear to be affected by the disease. Thirty years ago his height was 5 ft. 10½ in. At the present time it is 5 ft. 5½ in. This shows the extent of the spinal curves.

Upper Limbs.—The right clavicle is bowed anteriorly, and at its centre measures 1 in.; the left clavicle measures ¾ in. The surface is rough and nodular. The arms appear to be quite normal, with the exception of the right ulnar (see x-ray photograph, p. 14) and the injured fingers. The right elbow-joint appears to be a little stiff. It does not seem possible in any of the affected bones to say exactly what amount affected, the normal passing imperceptibly into diseased bone,

Right leg.

Left leg.

Right ulnar.

but in the case of the right ulnar it appears to be 3 in. or 4 in. The part affected is enlarged and nodular, as demonstrated by the x-ray photograph.

Chest.—The ribs do not show individually any alteration from the normal, but the chest in cross section (instead of being an ellipse, broader from side to side than from before backwards, in the proportion of 7 to 5) is a circle, measuring in circumference at the nipple line 37 in. The antero-posterior diameter at the nipple line is 10 in. The lateral axillary diameter is 10 in. These measurements were taken during forced expiration. The subcostal angle is more obtuse than in the normal chest.

The pelvic bones are apparently normal, but there is an increased breadth of the hips, particularly on the left side. The left trochanter stands out prominently, due to change of angle of the shaft and neck of the left femur, the neck joining the shaft at a right angle.

Lower Limbs.—Both are markedly diseased. The distance from the anterior superior iliac spine to the tip of the external malleolus is on the right 36 in. and on the left 35½ in. The left limb is more affected than the right. The fact that the neck of the left femur joins the shaft at a right angle accounts for this difference in length. The joints are apparently free from disease, but a few nodes similar to those seen in rheumatoid arthritis are found on the external surfaces of the condyles. The right femur is comparatively free, except that about 3 in. above the condyles it has the same circumference as the left at a similar level. Both legs show pigmentation of the skin around the external malleoli and extending up the leg. There is slight oedema of both ankles, especially at night or after walking. The veins of both legs are varicose. In both, as seen in the x-ray photographs, the tibiae are extensively diseased, whereas the fibulae are comparatively normal. The internal surfaces of the tibiae are very rough and nodular. In the x-ray photographs a central portion of the bone suggests the original size of the healthy bone, and the abnormal growth appears to be due to the deposition of lamellae of bone outside the compact tissue under the periosteum. The internal surface of the right tibia measures 2½ in., of the left 3½ in.

Nervous System.

Intelligence and memory good. Cranial nerves normal. Sensibility to touch, weight, temperature, pain, and muscular sense appear normal.

Reflexes.—Superficial abdominal and cremasteric reflexes, knee-jerk, ankle-jerk, plantar reflex flexion are present, and organic reflexes are normal.

Blood.

Erythrocytes number 4,900,000. No abnormalities. White blood corpuscles, 7,500. Differential count normal.

Thinking there might be some change in the amount of calcium present in the blood, I calculated the calcium index and calcium content by means of Blair Bell's calcimeter.

The calculation was made on several occasions, at varying hours. The results varied according to the hour at which the blood was taken, the index being high in the morning before breakfast and low after the midday meal. The average calcium index was 0.6, and the calcium content 1 in 9,000. The two highest counts recorded, strange to say, occurred at 5 p.m.—an hour at which the count should be low. The index in these cases was 1.2, calcium content being 1 in 5,000. These two counts, made at intervals of a month, occurred on days during which the patient suffered from severe headache and severe pains in the limbs.

According to Blair Bell, in health the calcium index may vary between 1.2 and 0.4, and from a calcium content of 1 in 5,000 to 1 in 15,000, according to the hour at which the blood is taken, being high in the morning and low at night. Judging by these figures, the patient never showed real excess or diminution in the calcium present in the blood.

Urine.

The amount passed in twenty-four hours was almost identical each day, the average being about 32 oz. On occasions it contained urates. Dr Scott of Liverpool kindly estimated for me the calcium, phosphorus, and nitrogen output in the urine. The patient was given a special diet, all food taken being weighed, the amount of nitrogen contained in such articles of diet being previously estimated. The fixed diet was given for three days previous to taking the first sample of urine, which was from a twenty-four hours' collection. In 32 oz. of urine, which was the average excretion,

The output of nitrogen was 20.7 grams.

The output of phosphorus was 2.20 to 1.53 grams.

The output of calcium was 0.0468 to 0.0108 grams.

Both the calcium and phosphorus figures are about the normal amount:

Ca ...	...	0.00425 to 0.00141 per cent.
CaO ...	...	0.00595 to 0.00191 "
P ₂ O ₅ ...	...	0.2536 to 0.1775 "
N ...	...	3.53 to 1.3 "

The calcium output shows no relation to the phosphorus output, one being high when the other is low. The figures approach the normal, except as regards nitrogen, which was mostly high—generally over 2.3 per cent.

TREATMENT.

Various drugs have been tried without benefit in osteitis deformans. In this case I first tried a course of calcium lac-

tate, with the idea that if the deposition of bone which takes place was Nature's method of strengthening the bones already weakened by disease, excess of calcium might materially aid the process. The patient continued to take calcium lactate at intervals for quite a month without the slightest benefit.

I next gave him citric acid as a natural antithesis to calcium, but without benefit. On the assumption that the blood of the patient was lacking in some internal glandular secretion, I gave him a course of Parke Davis haeminal, with the idea of supplying the missing secretion, but although treatment was continued for a month it was unsuccessful in removing pain or making the patient more comfortable. On the suggestion of Dr. Blair Bell I gave him thyroid extract, hoping that the missing secretion was that of the thyroid. This treatment had more effect than anything previously tried. After the dose had been increased to 15 grains a day the patient slept much better; a further increase of the dose up to 40 grains a day caused no diminution in the pains in the limbs or head. The dose was then gradually reduced, and finally treatment discontinued. Two months later sleep was about the same, and a great deal better than it had been previous to the administration of the thyroid extract.

CASE II.

A man, aged 62, brother of Case I, first came under observation by the patient coming to consult me as to the condition of his brother.

Personal History.

He had had rheumatic fever when a boy, which apparently left no after-effects. He had been a steady hard-working man, and had not indulged to excess in alcohol or tobacco. He had never had syphilis or gonorrhoea. For many years he had suffered from varicose veins of both legs, necessitating continual wearing of elastic bandages.

Present State.

In appearance he resembles his brother to a remarkable degree, most noticeable in the large size and shape of the head. There is slight kyphosis, and the head is held in a position similar to the attitude in his brother, but not so marked. His height as a young man was 5 ft. 8 in.; it is now 5 ft. 6½ in. When 23 years of age his hat was size 6½; he now wears a hat size 8½. He complains of frequent pains in his shin bones, particularly the right leg. On examination of the osseous system, there is very little change, except as regards the increase in the size of the skull, but there is a distinct nodule on the inner side of the right tibia, about 2 in. below the tubercle. There is also a thickening of the left tibia above the ankle-joint.

This case seems to me to be particularly interesting, from the fact that there appears to be previously no mention of this disease at such an early stage.

Macnamara considers osteitis deformans to be due to chronic rheumatic osteitis; all the patients, he says, complain of rheumatic pains; rheumatism affects aponeurotic and other dense fibrous structures and their prolongations into the bones, and he argues that rarefying osteitis must naturally follow, producing extensive areas of softened bone tissue, and this accounts for the bending under pressure.

All authorities agree that the disease is not associated with syphilis, and Paget states that the pain complained of is increased by administration of potassium iodide.

Gilles¹ recites two cases with autopsies. He concludes that the bony alterations are due to trophic disturbances depending on medullary lesions. In the first autopsy there was oedema of the cerebrum and cord, and on hardening a pale yellow coloration of the posterior cords, contrasting with the rest of the white substance, was observed. Sections stained with copper acetate made these differences visible to the naked eye. He acknowledges that these alterations do not appear to be a true sclerosis, but considers that they establish connections with other diseases of the nervous system, such as tabes. In the second autopsy he found paleness of the column of Flechsig, especially in the middle and inferior dorsal region; this he considered to be due to rarefaction of nerve fibres. In the superior and middle lumbar regions there were triangles of healthy substance separated by bands of altered tissue. As regards the significance of these lesions, he does not consider them due to fixing and staining, because the characteristics noted invade the posterior cords everywhere in the dorsal region, being either much less marked or absent in other regions. He states that in neither case were there any nervous symptoms during life, and explains this by the fact that previous to death when the cases were examined there was considerable general oedema. It may be asked whether this might not be the cause of the *post-mortem* condition of the cord which he described. I should suggest that it was most marked in the dorsal and lumbar regions owing to posture of the body, *ante mortem* and

post mortem. In any case I should imagine that such lesions, if present, would have caused some symptoms during life. In Case 1 I have failed to find any such nervous symptoms on careful examination.

Butlin regards osteitis deformans as a chronic inflammatory condition. On microscopic examination sections of the bones showed the Haversian canals wide, with the walls eaten out and rough, the so-called Howship's lacunae, which he considers characteristic of bone inflammation; the canals also showed lesions analogous to chronic inflammatory disease.

Lancereaux suggests that the new bone formation is an attempt at reinforcement of the long bones which have been weakened by absorption. Its distribution around the entire circumference, however, differs from other instances of compensatory growth, as seen in rickets, where the reinforcing bone is developed on the concave side of the shaft.

The association of cancer with osteitis deformans is interesting. Of Paget's 5 cases, 3 died from cancer: in two of osseous growths, in the third an epithelioma of the dura mater. Butlin states that sarcomata occurring may be accounted for as follows: The condition of the bones is attended by the formation of a large quantity of material rich in cells which is prone to sarcomatous degeneration, especially when, as in osteitis deformans, this material remains cellular and stored within the bones.

Blair Bell believes that the amount of calcium salts present in the blood is regulated by some internal secretion, and suggests the thyroid. Is it possible that the deposition of such salts is regulated in a like manner, and the derangement of that mechanism causes the trophic changes in the osseous system?

REFERENCE.

¹ *Nouvelle Iconographie de la Salpêtrière.*

A. DANGER ARISING FROM THE USE OF PLATED INSTRUMENTS IN OPHTHALMIC OPERATIONS.

By W. C. ROCKLIFFE, M.A., M.D., M.R.C.S.,
OPHTHALMIC SURGEON, HULL ROYAL INFIRMARY.

HAVING recently had the misfortune to lose an eye from acute panophthalmitis after extraction from a complication I am inclined to think is unique, I trust this report may be of value as a warning to others.

A. B., aged 73, was admitted to the Hull Royal Infirmary on February 15th, 1909, with ordinary senile cataract. He was a perfectly healthy old man, with no trace of catarrh or mucocoele. The eye, as is my custom, was cleaned and covered with a carbolic pad the night previous to the operation, every usual aseptic treatment was carried out during and after the operation, which was performed on February 19th, and was uncomplicated throughout.

On the following morning (February 20th) the lids were somewhat puffy, but not more so than is frequently seen as the result of pressure of the speculum; nevertheless, there was a slight mucous discharge, and the cicatrix did not look perfectly healthy.

On February 21st the eye was in a state of acute panophthalmitis. He was ordered leeches and perchloride irrigation every two hours; on February 22nd the galvano-cautery was freely applied to the cicatrix; nevertheless the pain and distension of the eye with pus became intense during the day, and the house-surgeon did paracentesis during the night, with great relief to the patient. On February 23rd the eye was excised.

Early on the morning of February 23rd, before the eye was excised, a bright foreign body was noticed during an irrigation floating away with the lotion, and was fortunately secured by the nurse.

This turned out to be a piece of plating measuring about 1.5 mm. by 1 mm., and clearly came from some instrument. I therefore examined the forceps and de Wecker's scissors, and found a corresponding deficiency of the plating near the hinge of the scissors, and a second larger strip almost detached. On inspecting my other plated instruments, some also showed signs of the plate peeling off.

I assume that during the iridectomy the fragment had dropped unseen into the anterior chamber or cicatrix and caused the panophthalmitis.

The question naturally arose how this septic condition

occurred, everything having been thoroughly sterilized and every aseptic precaution having been carried out, especially as it has been stated that a copper foreign body in the anterior chamber never causes suppuration of the vitreous. I might add that all instruments after cleansing are placed in pure alcohol.

Could the carbonate of soda in the boiling sterilizer, the 20 per cent. carbolic in the instrument dish, or the alcohol, have had anything to do with the cause of the separation of the plating?

Being by chance at the Royal London Ophthalmic Hospital, among others I mentioned the case to Mr. Bishop Harman. He suggested that a fault in the copper-nickel plating in the presence of a saline solution might set up galvanic action; this would explain the separation of the plating and also the subsequent inflammation produced in the eye. He therefore kindly undertook to submit the fragment to an expert Cambridge physicist, who reported "that he thought that pieces of double metal like this would very much strengthen the electrolytic action if they got into the saline fluids of the eye, and can quite believe that it would produce much greater inflammation than either metal singly. The strength of the action would depend somewhat upon the electro-magnetic force of a copper-nickel couple in the salt solution, but even if this were small the fact that the two metals are in actual contact would, of course, produce a comparatively strong current."

Since then Mr. Bishop Harman has obtained more such scales of plating and submitted them to Dr. Crocker, of the South-Western Polytechnic Institute, Chelsea, London, who reports:

The electro-magnetic force of nickel and copper electrodes in normal saline solution is 0.02 volt. When the electrodes are placed in perchloride of mercury solution, and the circuit closed, calomel deposits pretty readily on the nickel and mercury on the copper. It seems probable that the effect was primarily due to the action between the couple itself and the organic fluids of the eye, judging by the ease with which the couple corrodes near the surface of the liquid, in air.

These reports appear somewhat conclusive as to the cause of the peeling off of the plating, and the subsequent acute panophthalmitis, and I am much indebted to these gentlemen for them.

That the peeling off of the plating on steel instruments (although, as far as I am aware, not hitherto commented upon) is not so uncommon as it would appear, is proved by the findings of Mr. Johnson Taylor (Norwich) and my colleague, Mr. Roe. When I mentioned this case to them they inspected their instruments, and in each case parts of the plating on pairs of de Wecker scissors were found either partly or almost completely detached.

In conclusion, as a warning, I will suggest to all using plated ophthalmic or other instruments to examine them thoroughly prior to use, and I much doubt the advisability of ever using such instruments again.

Probably most ophthalmic surgeons of experience can recall cases of acute panophthalmitis following extraction, which they have been absolutely unable to explain. This case may suggest a possible cause of, what would have been in my case, a painful ignorance, had not my nurse detected the foreign body.

The spot in the square is the exact size of the detached piece of plating.

SEVERAL hospitals benefit largely by the will of the late Mrs. Charlotte Sarah Greenhill, of Norfolk Square, W. The Royal Ophthalmic Hospital, City Road, and the Children's Hospital, Paddington Green, are each given one-sixth of the residuary estate, and the Royal Free Hospital two-sixths. As the total residue estate appears to be over £100,000, the amounts which these institutions are likely to receive are very considerable. In addition, the Royal Free Hospital receives a direct bequest of £2,000 and the Children's Hospital one of £1,000. Like sums also go to the Middlesex Hospital and the West London Hospital.