photograph in Fig. 7 and in Figs. 10 and 14. Compare the fourth dorsal nerve distribution in Fig. 19.

Figs. 36 and 37 are rodent ulcers on points comparable to scleroderma in Fig. 14, leucoderma in Fig. 17. See also Fig. 19.

In Fig. 38 the rodent ulcer occupies the region of the ilio-inguinal nerve's distribution on the skin. See Fig. 19 and compare with the indistinct distribution of scleroderma on the right side in Fig. 13.

The observations contained in this paper strengthen the opinion which I have already expressed, namely, that the incidence of cancer may be due to the direct or indirect nervous influence of the area in which it occurs: as we now believe is the case in leucoderma and scleroderma.

Further, if I am right in believing that the rodent ulcers on the face and body coincide in incidence with scleroderma and leucoderma, and that these points of incidence are due to direct or indirect nervous influences, then I am right in saying that the tenability of part or all of the cleft-inclusion theory is threatened. In support of this I point out that the points of incidence of rodent ulcer on the trunk are not usually on the position of clefts as we know them.

As I have already said, I have never seen a rodent ulcer appearing on two or more points at which the same nerve root becomes cutaneous on the trunk, but only a rodent ulcer on one of the points. Hence the nervous influence, if present, which affects its incidence is probably either peripheral or a peripheral modification of central influence and not only segmental. The spread of rodent ulcer on the trunk, as it does on the head and face, appears on the whole to resemble the spread of scleroderma and leucoderma, and hence the spread as well as the incidence of rodent ulcer on the body is under a nervous influence which, if present, is peripheral.

Can this investigation throw any light upon the vexed question of the germ theory of cancer?

I have already pointed out that there are areas, around the mouth for instance, which are common to the spread of disease in processes known to be infective and of scleroderma, leucoderma and rodent ulcer. But there are others which are not common to them all, but only to leucoderma, scleroderma, and rodent ulcer.

Moreover, I have shown previously cases in which the spread of rodent ulcers have ceased in areas which had become denervated. And, finally, I cannot obtain sufficient evidence to support the idea that processes known to be purely infective are more liable to select those points of the skin at which nerves become cutaneous. However, on the whole time is not ripe for a definite statement on this point, although the little evidence I have just given appears to me to oppose rather than support the theory that cancer may be entirely due to the effects of a micro-organism.

Finally, I have to thank Sir Victor Horsley, Dr. Sequeira, Captain Pinch of the London Polyclinic, Dr. Pusey of Chicago, Dr. Norman Walker, and Mr. Hall-Edwards for many pictures of rodent ulcers.

REFERENCE.

1 BRITISH MEDICAL JOURNAL, April 18th, 1903, and December 12th, 1903.

CINNAMIC SALTS IN THE TREATMENT OF CANCER.

An Address Delivered before the Beckenham and Penge Medical Society.

By LOVELL DRAGE, M.D. OXON.,
Hatfield.

At the outset of this discourse it is necessary to apologize for venturing to believe that there is a possibility of the treatment of cancer becoming more generally recognized than it is at the present time by the members of our profession. The mere attempt at a treatment of the disease by any other method than that by the knife brings down on him who makes it a certain amount of discredit—to put the matter in the least objectionable manner.

In addressing a society of gentlemen engaged in the general practice of the profession, I have much less fear of such an attitude of mind than I should have in addressing many assemblages. We general practitioners tread out the corn after surgery has had its fill, and we know only too well what difficulties must be faced when the much-vaunted knife has had its swing. What matters it to us when pursuing our calling, and using our best endeavours, if we leave no remedy untried which may bring palliation to those who may be

numbered amongst the failures of surgery. What man amongst us can foretell if we who are cutting new furrows may not lay a foundation upon which the successful treatment of cancer, without the aid of surgery, can be made?

Sodium Cinnamate or Orthocoumarate.

It is my purpose to attempt to explain my position with reference to the treatment of cancer by the subcutaneous injection of a 10 per cent. solution of sodium cinnamate, or of a 22 per cent. solution of the orthocoumarate of the same base. Perhaps it will be well to state at once that the dose of the former is 30 minims, administered once or twice a week, and that of the latter 25 minims administered at the same interval.

For many years the problem presented itself to me, as it must have done to each of you—this problem of the treatment of cancer. It cannot be asserted that surgery, great as its triumphs undoubtedly have been, can be looked upon as providing the most scientific method.

The resort to surgery is to my mind nothing less than a confession of defeat. Here is a disease consisting of degeneration of cells, or, if it be preferred, a retrogression to the genetic type of cell. If this is accepted, it must occur to most minds that if the environment of the cells affected can be profoundly changed in that direction which leads to an arrest of this degeneration or retrogression, a successful result is not only possible, but certain. It is common knowledge at the present moment that it is perfectly easy to profoundly change the environment of cells by the artificial alteration of the composition of the blood. Thus, the experiments of Riether and Spiro, confirmed by Batty Shaw, prove that, at all events in cats, a very considerable leucocytosis can be produced by the intravenous injection of sodium cinnamate. This leucocytosis has been proved to be not the result of depriving the blood of a certain amount of liquid, but is the result of stimulating the simpler lymphocytes to transformation into the polymorphonuclear cells and intermediate forms.

So far as my observation is concerned—and it is supported by Dr. Randell, of Beckenham—both the coumarate and the cinnamate of sodium produce in the human subject a leucocytosis of this class.

The working hypothesis, then, upon which the whole treatment is based is this: If it is possible to make, artificially, a large increase in the most organized cells—I mean those most endowed with the properties of living cells—it is probable that such an alteration will be made in the environments of cells which have reverted to the genetic type that an arrest in the reversion is possible and probable.

Having arrived at this point, the first difficulty which presents itself is the provision of clinical material, surgery claiming all cases in early stages, but here is a point which I desire to especially emphasize: Do something more than removal by the knife in early cases of the disease when operation has been decided upon. Inject either of these drugs into the circulation before operation and for three months afterwards.

One of my cases, which made a great impression upon my mind, was that of a woman whose breast I removed for scirrhus. She was treated with the oil of cinnamon for a fortnight before operation, and for three months afterwards. That woman made a recovery from the operation of simple removal of the breasts, which impressed me very much, was at work in a short space of time, and has remained well after four years have nearly elapsed, although well past middle life. More than this, some eighteen months after the operation she showed me an old sebaceous cyst on her back which presented all the signs of scirrhus cancer in a state of atrophy, and this cyst has given no trouble, and has healed up under a scab.

Since this case a forward step has been taken, following upon the discovery of the therapeutic action of cinnamic acid. Cinnamon oil is practically a mixture of cinnamic aldehyde and cinnamic acid, and the aldehyde is rapidly oxidized into cinnamic acid when introduced into the circulation.

It may here be stated that the most active chemical substance at present investigated is believed to be the cinnamic aldehyde. Its administration, though less painful than the other chemical substances investigated, is exceedingly painful within twenty-four hours of administration, and has been given up for the present. My friend Dr. Morgan, of the Royal College of Science, South Kensington, first obtained for me the glycerine solution of sodium cinnamate, and then the sodium salts of the three isomeric coumaric acids.

Results.

Two cases of cancer in the region of the pharynx and larynx have been treated. In both relief from pain and distressing symptoms was experienced, and not only so, the discharges were free from the fetor usually associated with the disease when found in these regions. The patient died when defeated by the disease, and painlessly passed away in sleep.

In one case of lymphatic recurrence in a woman whose breast my friend, Mr. Lockwood, removed in St. Bartholomew's Hospital, the gland disappeared rapidly after treatment, and the elderly patient remains well two years at least after operation.

In a second case of growth in the breast, the patient, a nurse in the middle period of life, refused operation absolutely, but she clamoured for treatment because she suffered so much pain during the exertion of moving patients in bed that the fear of losing her work was always before her. I am forced to confess that I only undertook the treatment after pressing operation upon her as forcibly as is in my power, and refused to take any responsibility upon myself in case of disaster. That woman has been continuously employed in her work for two years past, and although the tumour cannot be said to have disappeared, it has markedly altered so far as the sense of touch is concerned, and now gives no pain. So far as my judgement goes it was a scirrhus tumour, but in the absence of microscopical evidence it is not my intention to assert its nature. The case is given for what it is worth.

A third case of breast tumour was more successfully treated, because the tumour disappeared altogether. The woman in whose breast this tumour was found had been previously treated for cancer in the abdominal cavity; the signs and symptoms presented certainly suggested to me that the patient was suffering from cancer, either of the pyloric end of the stomach or the upper end of the duodenum. All symptoms, such as pain and vomiting, slowly subsided after treatment, and the patient, after many months of comfort, died of heart disease, from which she had suffered for many years. It was only during an examination of the heart after its final breakdown that I discovered the lump in the breast; the patient had been afraid to tell me of this lump, which she had noticed three months previously. In advanced cases of cancer of the breast failures of surgery, no failure to alleviate has been observed except in two cases.

Abdominal Cases.

Of cases of cancer which have been tested by the microscope one of spindle-celled sarcoma of the spleen excited great interest. The unfortunate man had had one testicle removed, and had been under one operation for the removal of glands. When the patient came under observation a very large tumour of the spleen presented itself. The man, under treatment, lived for many months without pain, and died peacefully when the disease was victorious. It may be noted that when the treatment was interrupted, owing to inability to come to me, pain was complained of, and the treatment was then carried out, under my direction, by Dr. Berry of Grantham.

At the same time that this patient was under treatment, a man afflicted with a large growth of liver, which had been investigated by Mr. Lockwood by means of exploratory incision, came under my notice; the result was the same.

Several cases of cancer of the bowel have been placed under treatment, and the points noted have been relief of symptoms of obstruction and of pain. In one case there was a very marked cessation of collection of ascitic fluid. This patient had a considerable ascites when he first presented himself. After one tapping there was no further collection.

It appears to me that enough has been stated in order to demonstrate the fact that there is a remarkable reaction in patients when the drugs in question have been exhibited. The view taken of their therapeutic action is this: If the case is advanced it appears improbable from the nature of the disease to suppose that any treatment can remove it, but there is no doubt in my mind that the administration of the treatment suggested inhibits the activity of the organisms which are connected with putrefactive changes. The relief from pain is in all probability due to this fact, and so is the prolongation of life.

When this treatment was not in my hands it was my experience that patients died in a very large percentage of cases from the results of sepsis. In no case treated by my method have I seen a patient so die.

Patients, so far as my observation goes, die from simple exhaustion, and not with the old horrors connected with the

disease. The relief of obstruction in the bowels, however, rather inclines me to believe that even in old cases the treatment is capable of profoundly influencing the character of the growth, and certainly the treatment of very early cases leads in the direction of the belief that the disease is more capable of satisfactory treatment than is usually supposed.

Early diagnosis, as in all cases of disease, is here of the most supreme importance, and I believe that the treatment in the hands of those who have cases of difficulty connected with the abdominal cavity in which there is reasonable suspicion that the disease is cancerous in its nature will lead to surprise. I must confess to have experienced these feelings of surprise myself.

In conclusion, gentlemen, I express a hope that some of you, at all events, will be sufficiently interested to carry on the work further; and I may add that my friend, Dr. Morgan, of the Royal College of Science, has already furnished me with other allied chemical substances of which I have hopes.

A CASE OF CANCEROUS DISEASE OF THE STOMACH,

WITH SECONDARY INFECTION OF THE LIVER, OF THE THORACIC DUCT, AND OF THE CERVICAL LYMPHATIC GLANDS.

By W. MITCHELL STEVENS, M.D. LOND., M.R.C.P. LOND.,

Fellow of University College, London; Assistant Physician and Pathologist to the Cardiff Infirmary.

THE case which forms the subject of this communication is of especial clinical and pathological interest, being one of "latent" carcinoma of the stomach in which I was able to make an early and correct diagnosis by discovering the presence of glandular enlargement in the left supraclavicular region and in which a *post-mortem* examination showed extensive cancerous affection of the stomach and liver associated with marked cancerous disease of the thoracic duct and localized infection of the above-mentioned glands.

REPORT OF CASE.

J. D., a labourer, aged 46, came under my care as an out-patient at the Cardiff Infirmary on September 30th, 1904, complaining of swelling of the legs and genitals of about a week's duration.

History.

Except for an attack of ague in India in 1890, he had enjoyed excellent health until about twelve months ago, when he became pale and somewhat weak, and he states that he was then supposed to be suffering from "anaemia."

Since that time until the advent of the dropsy he had been able to do his work, though he never felt really well but had noticed that he was getting gradually weaker and rather short of breath. His appetite had been good; he states that he had not lost any flesh, he had no gastric symptoms, and, indeed, there had been no localizing symptoms of any kind.

Condition on September 30th.

The patient is somewhat sallow and exhibits a moderate degree of pallor, and there is slight wasting. While lying down he is perfectly comfortable, but when walking he is obviously weak and short of breath. There is no jaundice. There is extreme oedema of both legs and of the scrotum and penis, and slight puffiness of the lower eyelids.

The pulse is 84 per minute, of low tension, and quite regular. The heart and arteries present nothing abnormal except a very faint haemic murmur over the pulmonary area.

The lungs are perfectly normal. The tongue is slightly furred, the appetite is good, the bowels are regular, and there is an entire absence of any gastro-intestinal symptoms.

There is no abdominal swelling or tenderness, the spleen is not palpable, nor is the area of splenic dullness increased, and the stomach is neither tender, nor enlarged, nor can any physical signs be discovered except an enlargement of the liver, the lower border of which can be felt extending to two fingerbreadths below the costal margin. The hepatic enlargement is perfectly uniform, and there is no associated pain nor tenderness.

The urine is normal in every respect. A blood examination presented nothing except a moderate degree of diminution in the red corpuscles.

In the left supraclavicular region there is an enlarged gland about the size of a hazel nut which is situated just under the posterior border of the sterno-mastoid muscle, and is fairly movable, but very hard.