For the next ten days the urine contained blood in considerable quantity, showing the patency of the ureter; a few days after the blood disappeared pus occurred in the urine and continued for about three weeks—that is to say, she had pyelitis. This cleared up at the end of the three weeks.

The success of this plastic operation on the ureter is shown by the fact that there has been no recurrence of the hydro-nephrosis, and no other symptoms except occasional discomfort in the loin. It is doubtful, however, whether the kidney is doing much work.

CASE III.

A boy, aged 21, was admitted to hospital on September 19th, 1903, with a history that ten days previously he had suddenly been attacked with violent pain in the left lumbar region. This pain continued intermittently until admission.

On examination, there was marked tenderness to bimanual palpation of the left lumbar region, and muscular resistance prevented a satisfactory examination. The urine contained pus in small quantity, and microscopically a few red cells were also seen. He was examined by the x ray, but the radiograph showed no renal calculus.

His pain continued after admission, and on September 26th the left kidney was explored. The renal pelvis was distended and contained about 8 oz. of pus; no calculus was present. A careful examination of the pelvis and ureter showed that where the former passed into the latter was a very narrow constriction, above the level of the superior isthmus. This admitted of dilatation with a fine pair of dressing forceps pushed into the stricture from inside the pelvis. The pelvis was freely irrigated with water and the incision into its wall was closed; the wound in the soft parts was closed with drainage.

The after-history is interesting: For five days pus and blood were present in the urine in diminishing quantities, but on the sixth both had disappeared. The total amount of urine passed became markedly increased; from a mean of 30 oz. before operation it rose to an average of over 70 oz. for the twenty days following. The wound healed satisfactorily. I have seen this patient recently. On August 19th he had some pain in the left loin again for the first time since operation. I kept him under observation for a month, during which time he had no return of the pain or discomfort and no abnormal constituent in the urine.

REMARKS.

I wish to remark on the influence of renal stasis as illustrated in this third and other cases as a factor in the production of pyelitis and pyonephrosis. Cases of pyelitis (excluding the tuberculous form) are not uncommon where the affection is primarily of the renal pelvis and where no very definite cause for the condition is discovered. Fagge is content to dismiss such cases as probably due to renal calculi which have disappeared. It is known that the kidney is a regular channel of escape for bacteria during apparent health. It has been shown in dogs that with experimental bacteriuria a unilateral pyelitis can be produced by ligation of one ureter, the opposite kidney remaining apparently normal. It appears very probable that a certain number of the cases of primary pyelitis of obscure origin are dependent on renal stasis due to some defect in the mechanical conditions under which urine is expelled from the renal pelvis.

Pyonephrosis, as distinct from pyelitis, is no evidence that there was any primary obstruction to the urinary discharge, for we know that a pyelitis which is due to ascending infection from the bladder not infrequently becomes pyonephrosis from the obstruction caused by swelling of the mucous membrane and the impaction of shreds and detritus.

In this third case I think it may be assumed that the ureteral stricture was a determining factor in the pyelitis; what we know of the kidney as a portal of escape for the bacteria of the body, and what experimental work has shown to be the effect of ureteral ligation in the presence of definite bacteriuria, justifies the supposition that interference with the mechanics of renal discharge may play an important part in the etiology of similar cases of primary pyelitis.

The plastic procedure adopted in the second case is that which is usually spoken of as Mynter's method. It is suitable for those cases in which the stricture is situated at the junction of the renal pelvis and ureter or close below. For strictures below the level of the lower pole of the sac some other method has to be adopted. For these Küster's procedure is to be recommended, or the stricture may be excised and the

ends of the divided ureter brought together by ureteral anastomosis, as I have recently done in such a case.

The table on p. 1324 shows a high percentage of recoveries after plastic operations for ureteral constriction. In addition to the cases tabulated there are others quoted by other writers to which I have not been able to obtain access. They are as follows: Israel, 2 cases in his *Chirurgische Klinik der Nierenkrankheiten*, quoted by Moynihan, *BRITISH MEDICAL JOURNAL*, 1904, vol. 1, p. 1010; Gerster, 1 case, quoted in *Bryant's Operative Surgery*; Cramer, Elliot, and Albarran, quoted by Grosset, *Gaz. hebdom.*, 1900, p. 373; and a case, quoted by Young, in which Meyer found a stricture 6 in. below the kidney, complicated by calculus in the lower part of the ureter (*Medical News*, September 22nd, 1900).

The 33 cases given in the table include 11 which are quoted by Morris. In 5 no plastic operation was done on the ureter; of the remaining 28, in 4 the stricture was situated close to the bladder, in 24 at or close to the junction of ureter with renal pelvis. The latter were treated by three methods: (1) Fenger's operation, which resembles the Heinecke-Mikulicz pyloroplasty; that is to say, longitudinal incision and transverse suture; (2) Mynter's operation, as in Case II, described in this paper, which resembles the Finney operation for pyloric stenosis; (3) Küster's operation, section of the ureter below the stricture, and implantation into the lowest part of the renal pelvis. Fenger's operation was done in 12 cases, Mynter's operation in 3, Küster's operation in 11. In 2 of the cases Fenger's procedure was first done, and Küster's operation later owing to recurrence of the hydro-nephrosis; in the remaining 10 cases the operation was successful. The 3 cases treated by Mynter's method were all successful. Küster's method does not show so good a record—6 successes and 5 failures—but the operation should not be credited with some of the latter. In 1 case (Morris) nephrectomy was done on the table for the reason that the success of the plastic implantation seemed doubtful; in the remaining 4 cases the causes of failure were as follows: Ileus (Trendelenburg) after transperitoneal operation; necrosis of kidney (Helferich), second stricture lower down (Van Hook), infection (Bazy). In no case could the non-success be ascribed to a defect in the plastic operation itself, and it is to be noted that the 2 cases in which implantation was undertaken after the failure of Fenger's operation were successful.

Mynter's operation is particularly suitable for the cases in which the ureter is inserted high and obliquely into the renal pelvis; it gives a long slit-like opening, and when the incisions are extended sufficiently it has all the advantages which are afforded by implantation into the lowest point of the pelvis. The latter presents more technical difficulties; it is the method of choice when the stricture is below the level of the lowest point of the renal pelvis and involves so much of the channel that the longitudinal incision and transverse suture of Fenger is not applicable. Even where Fenger's operation is practicable, implantation is theoretically preferable, especially where the pelvis is markedly dilated, as it affords a more perfect drainage channel.

THE VALUE OF THE USE OF A SHADOWGRAPH URETERIC BOUGIE IN THE PRECISE SURGERY OF RENAL CALCULUS.

[EXTRACT FROM A POST-GRADUATE DEMONSTRATION.]

By E. HURRY FENWICK, F.R.C.S.,

Surgeon to the London Hospital.

[WITH SPECIAL PLATE.]

BEFORE the routine employment of expert radiography in cases of renal pain, the diagnosis of stone in the kidney and ureter was merely speculative. The clinician was rarely certain as to whether a stone was present or not, or if he was assured in his own mind that a calculus did exist, he could not say whether it was imprisoned in the kidney or whether it had passed into and had become arrested in the ureter; so similar are the symptoms evoked by stone in either position. The surgeon was forced, therefore, to explore, first the kidney and then the ureter before he could declare the patient free from stone.

Expert radiography* has, however, entirely altered the

* I do not allude to amateur, but to expert work. The statistics relate to Mr. Harnack's routine in our x-ray department.

character of the diagnosis and has modified the operative procedure. A clear shadowgraph affords the surgeon who can correctly interpret shadows a fair idea as to whether an operation is needed or not, it indicates the difficulties of the operation, affording a clue as to whether the surgeon will have to open the pelvis or cut through the kidney cortex or even remove the kidney. But it does more than this—it is able, by detecting a stone in the ureter, to draw the attention of the surgeon away from the kidney altogether, and to concentrate his efforts upon freeing that canal.

One of the most distressing of failures in urinary surgery is to cut into a kidney and mutilate it, perhaps irreparably, in order to find a stone which has long ago left that organ and travelled down the ureter; and yet this must happen in 22 per cent. at least of all cases of renal stone if the x-rays expert is not called upon to help in the diagnosis.

That the expert radiographist can guide the urinary surgeon with a precision unattainable by any other means is without cavil, and I assert this on an experience of 500 operations upon the kidney. In fact, I know of no obligation so great as that which the expert radiographist places the surgeon and patient under when he accurately defines a stone in the lower ureter.

But the interpretation of radiographic shadows is still in its infancy. There can be no doubt about such ureteric shadows as these I place before you. Here are half a dozen taken in the last three months. All these are stones, confirmed by operation (Figs. 1 and 2). You notice that they are in the line of the ureter, their outlines are sharp, and that their shapes are more or less oval. These are the three great features of ureteric stone shadows.

But how are we to be certain in cases when the shape of the shadow is ill-defined, when its size is small or when it is apparently away from the line of the ureter? It cannot be to the patient's benefit or the surgeon's credit to make a huge extraperitoneal flap, and search up and down the ureter for a stone which does not exist. Personally speaking, when I am in doubt I pass along the ureter a bougie which casts a shadow with the x ray. The patient is then radiographed. As the bougie just fills the canal of the ureter, I can tell at once by tracing the ureteric shadow of the bougie as to whether a stone is in the ureter or not.

Now, it may well happen that this expedient has been resorted to by many others. It is so simple and easy in skilled hands, that I can hardly believe it has not been utilized in cases of doubtful shadows. I have not seen any publication, but this is no matter; the idea is practical, and the procedure will save much unnecessary mutilation.

Let me illustrate. Dr. Bertrand Dawson brought me a nurse suffering from severe right renal colic. He showed me radiographs, taken separately by Mackenzie Davidson and Mr. Harnack, showing two large shadows visible in the true pelvis in the track of the right ureter (Fig. 3), and he suggested they were due to calculi in the right ureter, and that these were the cause of the renal colic. I demurred, because the shadows were large and irregular. To prove to him they were away from the region of the ureter, I passed a catheter with a lead styilet up both ureters, and Mackenzie Davidson radiographed that patient again with these hard-cored catheters in position. Here is the radiograph (Fig. 4) You notice that the ureteric bougie shadow is far away from the supposed stones, thus proving uncontestedly that these shadows are not stones in the ureter. I may as well tell you the sequel. Finding the passage of my right ureteric bougie obstructed 3 in. below the kidney—a point, of course, well above the pelvic shadows—I cut on to its point and found a small oxalate of lime calculus in the ureter at that spot. This is what had caused the severe right renal colic. I turned to the radiograph and found the shadow of that stone, which I had overlooked—I suppose because I was so intent in deciphering the larger shadows below the true brim of the pelvis. To make the clinical aspects of the case complete I made a small peritoneal opening in my original wound and drew up from below the pelvic brim the packet of calcified mesenteric glands which had cast the large shadows. They were then excised.

The next pair of radiographs (Figs. 5 and 6) were taken by Mackenzie Davidson from a lady who was sent to me with bilateral renal suffering by Dr. Athel Saw, from Perth, Australia. They afford a very excellent illustration of the extraordinary precision of the method.

Mackenzie Davidson drew my attention to a shadow, suspicious in shape, position, and size, of a small stone lodged in the right ureter (marked by arrow, Fig. 5). But the

pelvic contents were very dense and dark, and had to be lightened up—and I want you to take this valuable hint how to clear up the radiograph of the lower pelvic contents. You see the white half moon, and on it—or in it, rather—the little supposed stone and the ureteric bougie *in situ* (Fig. 6). The clearing of the picture is entirely due to filling the bladder with air. Always fill your bladder with air in radiography if you are in doubt as to a stone in the lower ureter, whether you use a ureteric bougie or not. The air is introduced by merely passing a catheter and elevating the pelvis.

But to return. This little shadow (Fig. 6) is no stone. You see it is outside the ureter, for the ureter is filled by the bougie. Now the sequel. I had to perform laparotomy for another reason, and took the opportunity of finding out the cause of that little shadow. It was a patch of atheroma at the branching of large artery. This is a clinical fact of which I made certain—that an atheromatous calcareous patch can cast a shadow.

I propose now to take you to my operating theatre and demonstrate the technique of the procedure in the case of a girl with right renal pain and calcified mesenteric glands. The radiograph shows shadows (Fig. 7, marked by arrows) which, as she has had severe pain in the region of the right kidney and ureter, have been diagnosed as stones in the kidney and ureter—I assure you they are calcified glands and not stones.

Fig. 9.—Ureteric bougie when passed from above sagging down ureter just at its bend into the bladder.

[The patient was then anaesthetized, placed in the Trendelenburg position, the ureteric catheter was passed by Kelly's method (time 10 seconds) and radiographed by Mr. Harnack (time 5 minutes). The plate was then developed and shown at the conclusion of the demonstration (Fig. 8).]

The radiographic bougie stands out plainly. Its deep shadow is due to its being a very excellent shadowgraph bougie, which was specially designed and made for me by Mr. Marshall*; it is solid, aseptic, easily passed, and proves very dense to x-rays. It is much better in every way than my original hard-cored catheter. It will be noticed that the point has been bent down into a lower calyx, and its entire track is far away from the shadows. Case was confirmed by exploration, calcified glands being found in mesentery.

As a last and very beautiful illustration of the value of this method I now draw your attention to a series of radiographs, in which little round shadows appeared in the area of the bony pelvis, shadows which lie three or four together apparently along the true brim, or one at the ischial spine, or along

* It can be obtained at Croyden s. 55, Wigmore Street. It is scheduled as the Marshall shadowgraph ureteric bougie.

Fig. 1.—Arrested ureteric stone (shadow arrowed), severe renal symptoms for 25 years; operation; cure. (Taken by Dr. Mackenzie Davidson.)

Fig. 2.—Descending ureteric stone (shadow arrowed) subsequently passed; two years' history. (Taken by Mr. Harnack, London Hospital.)

Fig. 3.—Radiograph of bony pelvis, shadows (arrowed) in track of right ureter, supposed to be calculi in ureter.
(Taken by Dr. Mackenzie Davidson.)

Fig. 4.—Radiograph of bony pelvis of same case; both ureters occupied by shadowgraph bougie, showing shadows to be outside ureters proved operatively to be calcareous glands in mesentery. (Taken by Dr. Mackenzie Davidson.)

Fig. 5.—A shadow (arrowed) supposed to represent small calculus in right ureter, low down. (Taken by Dr. Mackenzie Davidson)

Fig. 6.—Same patient; small shadow (arrowed) shown to lie outside ureter, which is filled with a shadowgraph bougie; shadow proved to be a calcareous patch in an atheromatous artery; notice semilunar clear space due to inflation of bladder. (Taken by Dr. Mackenzie Davidson.)

Fig. 7.—Shadows (arrowed) supposed to represent stone in ureter and kidney. (Taken by Mr. Harnack, London Hospital.)

Fig. 8.—Same patient ; ureter occupied by shadowgraph bougie which turns down into lower calyx ; shadows shown, therefore, to lie outside ureter and kidney ; proved operatively to be calcareous glands in mesentery. (Taken by Mr. Harnack, London Hospital.)

the track of the ureter. These are not stones, they are calcified glands and are negated by the ureteric bougie.

There is but little difficulty in the male in passing these bougies; they are introduced through a Caspar ureter cystoscope or through an ordinary prostatescope or open urethroscope tube, the pelvis being elevated. In the female, a Kelly tube is used. There is decided abdominal discomfort after the bougie has been passed, just as there is after ureteric catheterization, but this is relievable by a little opium.

If you have not the x ray to help you, of course the kidney must be explored first, and the ureter opened and a tube passed down it, but remember these two objections: The ureteric catheter passed from above can easily glide over the mucus-covered stone—for all arrested ureteric stones are embedded in thick jelly and the surgeon may be unaware that he has touched a stone. Secondly, the bougie may never reach the bladder at all, but press the ureter down in front of it just at that spot where the tube bends upwards to enter the bladder (Fig. 9).

I do not claim any credit or originality for the shadowgraph ureteric bougie. If any is due it would fall to Kelly or Nitze. I only urge it as a valuable method, and one likely to prevent needless operation in the lower ureter.

ONE HUNDRED CONSECUTIVE LAPAROTOMIES FOR INTUSSUSCEPTION IN CHILDREN.

By CHARLES P. B. CLUBBE, M.R.C.S.ENG.,

Honorary Surgeon, Royal Prince Alfred Hospital; Honorary Surgeon, Royal Alexandra Hospital for Children, Sydney; Joint Lecturer in Clinical Surgery at the Sydney University.

In this part of the world, as elsewhere, we now get more intussusceptions in hospital than in former years, because the condition is more readily recognized and the children are sent to us more promptly than heretofore. I find that up to November, 1904, I had performed 100 laparotomies for intussusception in children, with sixty-three recoveries and thirty-seven deaths.

The first 50 cases occurred from August, 1893, to April, 1901 (a period of seven years and eight months); of these, 25 lived and 25 died. The second 50 cases occurred from April, 1901, to November, 1904 (three years and six months); of these, 38 lived and 12 died. The great difference in the mortality in the second 50 cases is probably due to the fact that the children were sent into the hospital earlier.

	Cases that Lived.	Cases that Died.
Average time from onset of symptoms to operation	23 hours.	68 hours
First 50	23 "	48 "
Second 50		

The varieties were as follows:

	Cases.	Cured.	Died.
Ileo-caecal	64	43	21
Ileo-colic	12	7	5
Double	20	11	9
Colic	3	2	1
Enteric	1	—	1
	100	63	37

In my cases the relative frequency of the various forms differs considerably from those collected by Weiss, who gives as the following statistics: Ileo-caecal, 43; ileac, 23; ileo-colic, 14; colic, 26.

It appears that I have met with singularly few enteric and colic varieties. Cases I have classified as "double" were primarily enteric, generally starting a few inches from the caecum. This intussusception reaching the caecum has caused the secondary infolding of the colon. I have noticed that in the ileo-caecal, the common variety, the starting point is in the large majority of cases in the caput caeci and not at the valve. This free end of the caecum is often a little difficult to unfold. The main part of the intussusception has been reduced, the lumen is free, but this small, cup-shaped depression, with its thick, hard wall, persists, and has often to be moulded into a convexity, and held in that position for

a minute or two. The arrangement of the muscular fibres at this part of the caecum seems, in some way or other, to tend to the production of this small cup or dimple, since it appears to be so frequently the starting point in what is the commonest variety of intussusception. In intussusceptions beginning in the ileum the first unfolding seems frequently to start at the side of the bowel and always opposite the mesenteric attachment.

DIAGNOSIS.

This is not a matter of great difficulty. The signs and symptoms are now so well known that, even with an ordinary amount of care, the condition should not be overlooked. Indeed, the Sydney graduate is now nearly always able to "spot" the case at once. If there is any doubt, and the mass cannot be felt, chloroform should always be administered for the purposes of diagnosis. The condition of the child is apt to mislead the unwary, because in a large number of what may be called early cases, that is, cases seen within eighteen or twenty hours from the onset, the condition of the child is remarkably good. The pulse rate may not be raised, and they do not look ill. It is difficult to believe that they are suffering from acute intestinal obstruction. I think this is the reason why men sometimes miss these cases, if they are not on the watch for them, and have not paid attention to the story of the onset of the attack told by the mother. In some cases the children are collapsed from the very onset, and in later cases it is nearly always easy to see that the child is profoundly ill.

IRRIGATION.

In spite of the fact that nearly all surgeons now condemn irrigation as useless and unscientific, I always inject before operation. The patient is anaesthetized, everything is prepared for operation, and then oil or warm saline is injected by means of an ordinary Higginson's syringe. Why is it useless? It may complete the reduction. It always reduces the mass to a certain extent, sometimes very largely, and in a very gentle manner. Frequently a mass that was in the rectum or descending colon can, after the injection, be felt in the caecal region, and can, therefore, when the abdomen is open, be now got at and reduced with much less handling, and therefore much less risk, than before. But irrigation alone must never be considered a complete form of treatment, and when we irrigate we must always be prepared to operate. My experience has taught me that it will only do away with the necessity for operation in 10 per cent. of the cases. It is, therefore, always necessary where the mass disappears entirely, as the result of irrigation, to keep the child under careful observation. If these facts are recognized I fail to see how irrigation can be considered dangerous or unscientific.

INCISION.

In cases where the mass has been reduced to the region of the caecum, I usually make incision at the outer border of the rectus. The mass can then be easily delivered and reduced without any prolapse of the intestines and therefore with but little shock to the child. The advantages of the side incision are certainly less immediate shock, but it is more likely to be followed later on by hernia than the central incision. Where the mass is in the transverse or descending colon the incision should be in the middle line, the umbilicus being the centre, on one or other side, of the cut. I always try to keep the intestines in, but rather than waste much time I let them prolapse, and after the intussusception has been reduced I have warm saline solution poured over them as they are being replaced. If the serous coat of the intussusciens is torn I stitch it with fine catgut. I notice that the appendix is nearly always discoloured, but unless it seems to be very much damaged I rarely remove it. In this series of cases I have been obliged to resect the bowel seven times, for various reasons, with but one recovery—a case I have already recorded.

To close the abdomen I nearly always use through-and-through silkworm gut sutures with a Reverdin's needle. It saves much time. With a side incision, if there is not any great necessity for haste, I sometimes sew up in layers with catgut.

AFTER-TREATMENT.

They are always given morphine, $\frac{1}{10}$ to $\frac{3}{10}$ gr.; this is repeated if necessary. Strychnine is given hypodermically till the immediate shock passes; sometimes salines. Within twelve hours the colon is washed out with a weak solution of permanganate of potash. This nearly always produces a motion. If there is any difficulty with the bowels small doses of calomel or castor oil are given.