the raw sterile ascitic fluid were added. At the end of a week at 37° C. acid reactions had appeared in the case of glucose and galactose, but in the case of saccharose and maltose the reaction of the medium was unchanged. All three colonies responded in the same way to these carbohydrate tests.

6. *A Gram-negative Coccus from the Normal Urethra*.—This micro-organism was isolated from mucus present in a sample of normal urine and was found to grow readily on ordinary agar at 37° C., and on ordinary gelatine at 20° C. When examined in the carbohydrate tests the coccus rapidly produced an acid reaction in all four, namely, in glucose, galactose, maltose, and also in saccharose.

The Chief Characters of the Six Gram-negative Cocci.

Number.	Source.	Indifferent Specimens Examined.	Different Colonies Tested.	Growth on Naggar at 37° C.	Growth on Gelatine at 20° C.	Carbohydrates. Production of an Acid Reaction in			
						Glucose.	Galactose.	Maltose.	Saccharose.
1	(<i>M. catarrhalis</i> .) Nasal and pharyngeal secretion in 22 cases of the Hertford outbreak. Sputum chronic bronchitis	23	65	Opaque granular	Positive	—	—	—	—
2	Nasal secretion of 2 Hertford cases in both of which <i>B. diphtheriae</i> also present	2	4	Opaque granular	Negative	+	+	+	+
3	Nasal secretion of 1 Hertford case in which <i>Coccus</i> No. 1 also present	1	4	Clear, smooth, later becomes yellowish	Negative at first, positive later	+	—	+	—
4	(<i>Meningococcus</i>) Lumbar puncture 2 specimens; meningitis post-mortem 1 specimen	3	9	Clear, smooth	Negative	+	+	+	—
5	(<i>Gonococcus</i> .) Urethral pus	1	3	No growth unless blood added to medium	Negative	+	+	—	—
6	Mucus from normal urethra	1	2	Opaque, somewhat granular at centre but smooth at edge	Positive	+	+	+	+

CONCLUSION.

It follows from these observations that while the virulence for mice and guinea-pigs of the various Gram-negative cocci examined appears to be of little, if any, value for differentiating them, the effect of these cocci upon glucose, galactose, maltose, and saccharose, respectively, is of much value for this purpose. Owing to the special pathogenic significance of the meningococcus, gonococcus, and possibly also of *M. catarrhalis* and other individuals of the group, it is advisable in future, when examining a Gram-negative coccus, to test it, amongst other things, in all the four carbohydrate media before considering that its identity has been established. If mistakes are to be avoided, this procedure would seem to be particularly desirable—I may even say necessary—when examining nasal or pharyngeal secretion for the meningococcus, or when examining discharge from the urogenital tract, or other material, for the gonococcus.

REFERENCES.

- ¹ Paper read before the Epidemiological Society on January 20th, 1899.
² Memorandum by Dr. Farrar to Local Government Board, September, 1903.
³ *Flugge, Micro-organismen*, Band II, 1896, pp. 154-5. ⁴ Loc. cit.
⁵ *Zeitschr. f. klin. Med.*, Band XLIV, 1901. ⁶ *Centralbl. f. Bakt. Originale*, Band XXXIV, 1903, p. 641. ⁷ *Bull. Soc. Méd. Hôp. de Paris*, March 2nd and 10th, 1905.

OLD AGE.—The London Vegetarian Association has arranged a meeting at which the question what period of life should be termed old age will be discussed by a number of gentlemen between the ages of eighty and ninety, who are still engaged in literary, medical, agricultural, and other work. The chair will be taken by Mr. C. P. Newcombe, who is an octogenarian, and other speakers will be Professor Mayor, Mr. Joseph Wallace, and others. The meeting will be held in the Memorial Hall, Farringdon Street, on October 13th.

RATS IN RELATION TO PLAGUE.

By BRUCE SKINNER,
Lieutenant-Colonel R.A.M.C.

SINCE the publication in the BRITISH MEDICAL JOURNAL of May 6th, p. 994, of a paper on Plague and the Geographical Distribution of Rats, and, indeed, since the following remarks were drafted, I have become indebted to Lieutenant-Colonel Leishman for a reference to a paper by Dr. Rudolf Abel relating to Rats and Plague.¹ This paper contains a record of research into the ancient accounts of plague epidemics extending back to that described in 1 Sam., chap. v and vi. These records associate rodents and many other animals with plague. Dr. Abel comments on the want of reference to rats in relation to the plague epidemics which have occurred in Europe since 1730, and regards this as pointing to the fact that improved sanitation is probably the reason why Europe has since that date been less liable to plague epidemics. He suggests that perhaps *M. rattus* was less predisposed to plague than *M. decumanus*, and that as *M. decumanus* "belongs to the species that now in India and China suffers so much from plague, the rats in South and East Europe ought to have played a much more important part in plague epidemics, which is not the case." "Also," he says, "we are not justified in assuming that *M. rattus* shows less predisposition for plague than *M. decumanus*, as the matter has not been settled by experiment." He concludes that his historical survey shows that the most serious plague epidemics known which ravaged Europe in olden times without special participation of the rats and mice warn us that these rodents must not be regarded as the first cause in the epidemiology of plague.

Dr. Abel has not conceived the possibility that *M. decumanus* may be responsible for the comparative freedom of Europe from plague, as previously explained, evidently because he is under the impression that this species is the rat of the areas of endemic plague. We know, however, that *M. rattus* is the indigenous species of the plague endemic areas; and we are justified in inferring that if the latter species causes or spreads plague, he does not do so in Europe because he is now a very subordinate occupant of that area.

Some further observations on the habits of the different species of rats are, however, required, especially in their relation to plague.

The following extract from a classical writer is not without interest in its bearing on the part played by rats in its dissemination:

Neither are they (the Iberians) the only sufferers in being tormented with vast swarms of mice, from which pestilential diseases have frequently ensued. This occurred to the Romans in Cantabria, so that they caused it to be proclaimed that whoever would catch the mice should receive rewards according to the number taken, and even with this they were scarcely preserved, as they were suffering besides from want of corn and other necessities.²

Although the word mice is used we may accept it in its broadest sense as a family name, as the differentiation of the family into genera and species seems to be modern. It does not necessarily mean, as I have seen suggested, that the Romans were unaware of the existence of rats.

One point in the above quotation which especially merits attention is that the Romans employed the modern unscientific method of dealing with the mouse family when plague was prevalent.

We in the present day are aware that long-tailed rats suffer from plague. But we are not aware that they initiate plague. Analogy rather points to the certainty that these rats contract plague from some extraneous source, as animals cannot be said to initiate any specific fever.

That source has not yet been discovered. But inasmuch as it must be something outside the rat, it would appear more scientific to adapt our defensive measures to our immediate surroundings rather than to destroy those animals whose misfortune it is to suffer in common with ourselves from pestilence. Further, in our country at least, rats in towns are valuable scavengers.

There are certain seasons when rats betake themselves to houses. For instance, in England, the short-tailed rat, when he has adopted a country life, lives entirely out-of-doors from spring to autumn; in winter he resorts to corn-ricks, bean-stacks, barns, and houses.

The town rat lives usually in sewers. Where there is a *rus in urbe*, some of the town rats may be classed as to habits with the country rat. The great plague epidemics of England raged during the period when the rat lived out-of-doors.

The foreign country which perhaps most interests us, so far as plague is concerned, is India. Inasmuch as India is a country of such vast extent that climatic changes are neither equal nor synchronous in different localities, the habits of rats proper to one locality will be found to vary as to time when compared with those of other localities. Broadly speaking, the wet season, or monsoon, is the great climatic change which most profoundly influences India.

There are but few localities which are so unfortunately placed as to escape the downpour of rain which commences in May or June, or even later, according to the distance of the locality from the south-west source of the vapour-laden winds. And with the advent of the monsoon the rats of India take shelter. "In dry weather most of these gentry live out-of-doors, but the first heavy, steady, soaking rain is the signal for a general invasion" of houses.³

The Bombay rats, therefore, should take to the houses in May or June, and the Punjab rats at the end of June or beginning of July. It is possible that in the Punjab the rats may also in the cold weather take shelter, as the cold from November to the end of January is sharp. Perhaps naturalists in India may be able to furnish exact information on these points from various parts. If it be borne out by investigation that rats throughout India live out-of-doors, as in England, during the warm season, and only invade houses during the monsoon, or during the cold months, the fact should cause us to review our ideas as to the part taken by them in the propagation of plague. A glance at Appendix I in Professor Simpson's *Treatise on Plague* will show that in 1904 the plague was at its highest when the rats should have been out in the field, and at its minimum when they took shelter during the rains. See also Diagram B on page 148, in connexion with the fact noted above. [Note.—In Mauritius, while every month shows some rain, the rainy season is from the end of December to the end of April.]

The report of the Sanitary Commissioner with the Government of India for 1903 bears out the same conclusion (see Appendix B to Section VI, Table IV, for the plague incidence, and Appendix to Section I, Table II, for the rainfall). Here it at first sight appears that Madras forms an exception, for the plague incidence in that Presidency was most severe in January, during which month the Madras meteorological station showed 4.53 in. of rain, and dry weather in March and April, when the plague deaths were falling. But on more detailed investigation the apparent exception demonstrates the necessity for local comparisons of the rainfall and plague incidence. Some of the Madras districts (see p. 94) did not suffer from plague, while the rainfall was widely different for different stations. Bellary district had the most severe plague incidence in the Madras Presidency, while Bellary station had no rainfall until May. There is not, in the tables relating to Madras, sufficient differentiation of data to argue for or against the condition which has been pointed out above as holding in other areas where the climatic conditions, as well as the plague outbreaks, are more uniform than in the Madras Presidency. The Bengal Presidency also contains some localities where the rainfall is out of proportion to others as to amount and duration, and therefore generalized tables lose much of their value, though even as they stand they show a great fall in plague deaths during the wet months when rats should be in houses. In these cases especially, but in all cases for purposes of exactness, the actual meteorological conditions for each locality should be tabulated with the local plague incidence and with the dates of migrations of rats from houses to the fields and of their return. Broadly speaking, the Punjab presents fairly uniform rainfall, and the following table shows the plague-deaths in conjunction therewith in 1903:

	Jan.	Feb.	March.	April.	May.	June.	July.	August.	Sept.	Oct.	Nov.	Dec.
LAHORE.												
Plague deaths	8,906	16,435	36,789	62,907	40,426	14,270	163	65	151	1,961	4,267	5,728
Rainfall	0.52	0.5	0.64	0.02	0.82	0.33	4.36	5.31	2.25	0.10	0.0	0.31
BOMBAY.*												
Plague deaths { Presidency	38,310	38,476	32,188	17,571	3,931	3,268	8,033	17,454	30,489	37,819	29,179	24,453
City	1,693	3,486	5,858	5,721	1,580	657	347	335	322	262	220	270
Rainfall	0.0	0.0	0.17	0.0	7.79	18.64	24.2	18.76	9.02	5.91	0.0	0.0
CALCUTTA.												
Plague deaths	245	1,058	3,526	2,596	320	74	52	55	76	71	81	68
Rainfall	0.21	0.64	0.77	1.71	1.53	10.7	6.35	10.17	14.02	8.02	0.02	0.0

* The same seasonal relation is shown by the charts in the Report of the Plague Commissioners, vol. I, p. 513, and vol. III, p. 647.

As was formerly the case during the great plague epidemics of Europe, plague in India is most acute during the period when the rats are out in the fields. The plague diminishes rapidly after the wheat harvest, almost disappears in the rains, and begins again with the millet harvest. It is not here suggested that the rains have anything to do directly with the plague—the tables show that plague has commenced to fall before the rains begin.

But what I do suggest as borne out by the examination of the habits of the rat, so far as they are known, is that though he suffers from plague he is not the source of the disease. Were he the source, the epidemic should be most acute when the rat is in most intimate contact with the human population. The agriculturist does well to destroy rats on account of the enormous damage they work in the farms; but such a measure carried out as a preventive of plague cannot be regarded as scientific, while it must be classed as waste of energy when the work done is measured against the object aimed at, namely, the reduction of plague incidence. There is nothing to show that destruction of rats reduces plague incidence. It is true now, as formerly, that "even with this they were scarcely preserved, as they were suffering besides from want of corn." Perhaps it may be that the want of food caused by the depredations of rats, made the people more liable to plague. Such depredations are also suggested in the Biblical account of the epidemic among the Philistines: "And in the villages and fields in the midst of that country there came forth a multitude of mice, and there was the confusion of a great mortality in the city."⁴ Further on we read: "the mice that have destroyed the land."⁵ We shall see later that the same occurred in India.

But inasmuch as rats suffer from plague, which some have suggested⁶ is conveyed from them to man by the flea peculiar⁷ to *M. rattus*, many may consider it politic to exclude rats from dwellings. The feet of rats are covered with an extremely delicate integument. Rats, therefore, avoid lime, for it burns their feet. The free use of unslaked lime placed in rat holes, along the runs they make to water supplies, on the floors of cellars, and generally about the approaches to houses, will keep rats away. *M. decumanus* is an extremely thirsty animal. Perhaps *M. rattus* also possesses an equal love of water. By cutting off the water supply by some such means as that mentioned above the rat will be caused to remove to another locality. Should it be the case that the plague-stricken rat comes to the house to die, the use of unslaked lime in the house as here indicated may suffice to ensure his remaining elsewhere. If unslaked lime is scattered over the mud floors of Indian huts, and subsequently watered, insects will be destroyed, and the suspected plague carrier may possibly be thus removed. But if it be true that plague deaths are fewer when rats swarm in the houses, it would appear desirable not to exclude these scavengers, at least during the rainy season, but rather to ascertain if there is any fact to be learnt from their habits which may give a clue to the source whence plague is derived.

The intelligence, assisted by the keenness of the sense of smell possessed by *M. decumanus*, and presumably also by *M. rattus*, is so keen, that it is more than likely that rats of either species would not return even in the rains to plague-infected dwellings if the cause of that disease were inherent in the dwellings. The rat, however, does return at that season and remains throughout the rains, so it may be allowed that he cannot be responsible for the recrudescence of plague which occurs after the rains, and which must be due to some other cause apart from rats coming into play after a period of quiescence. In the Punjab the rat returns to the house not later than the beginning of July. The plague begins to increase at the end of September, when the rat is returning to the fields after the rains, and when the cultivator

is harvesting his millet crops. It continues throughout the cold weather, becomes virulent during the hot weather, and rapidly drops just before the rains, after the wheat crop has been gathered.

It is thus evident that during this period, when both rats and mankind are out in the fields, they are liable to attack by some agent common to both as causing plague, and that this cause decreases to a minimum during the period when rats and men are most closely associated during the rains.

This common cause is one which has not been in constant universal action throughout India, though it must have been of fairly frequent recurrence in Kumaon and Garhwal. The Indian epidemic of 1896 did not spread from the latter districts, but appears to have originated in Bombay. Plague was discovered in Bombay city in 1896. Possibly it existed in the Presidency before that year, and was only recognized when it had attained epidemic proportions.

What this common cause may be forms a subject for further consideration.

REFERENCES.

¹ Geschichtliches über Ratten und Pest, by Dr. Rudolph Abel, *Zeitschr. für Hygiene*, Band xxxvi, p. 89. ² Strabo's *Geography*, B. iii, civ, para. 18; translated by H. C. Hamilton; Bonn's Libraries. ³ *The Tribes on My Frontier*, by E. H. A., ed. 1887, p. 12. ⁴ Douay Bible, 1 Kings, v. 6. ⁵ *Ibid.*, vi. 5. ⁶ Simond, *La Propagation de la Peste*, *Ann. de l'Inst. Pasteur*, Oct., 1898, pp. 669 et seq. ⁷ Capt. Liston, I.M.S., *Plague, Rats, and Fleas*, *Bombay Nat. Hist. Society Proceedings*, Nov., 1904.

SEVENTY-THIRD ANNUAL MEETING

OF THE

British Medical Association.

Held at Leicester, July 24th, 25th, 26th, 27th, and 28th, 1905.

PROCEEDINGS OF SECTIONS.

SECTION OF OPHTHALMOLOGY.

G. A. BERRY, M.B., F.R.C.S.Ed., President.

PRESIDENT'S INTRODUCTORY REMARKS.

FIFTY years ago such rapid strides were made in the science of ophthalmology and in ocular therapeutics that every year brought its record of important discoveries in diagnosis, of new methods of examination, of far-reaching generalizations, often of real advances in ophthalmic surgery. At that time, indeed, a few years sufficed to revolutionize the whole science of this department of medicine, until ophthalmology attained to a degree of perfection which led von Helmholtz to say of it that it was "the most strictly scientific department of medicine, the one which most nearly approached to the exact sciences."

That ophthalmology can hardly, with the same justice, be said to occupy so prominent a position at the present day is due to the rapid advances which have been made more recently in other departments, and more particularly in other departments of surgery.

Some few advances in our special science are no doubt of comparatively recent date, for example, the introduction of antiseptic and aseptic measures in connexion with injuries and operations; the use of cocaine as a local anaesthetic; the appreciation of the rôle played by microbes and the toxins they engender, in connexion with various inflammations of the eye; some few points in pathology, notably with reference to tuberculous manifestations; the use of the electro-magnet for the removal of particles of iron and steel, the application of the Roentgen rays for their detection and localization, etc. From the therapeutical point of view, however, there has been otherwise but little advance to record, while the knowledge of the connexion between diseases of the eye and diseases of other parts of the body is hardly less crude than it was fifty years ago, except perhaps as regards ocular manifestations in certain nervous disorders. Latterly, indeed, instead of leading the way the ophthalmic surgeon has mostly had to follow, and adapt to the requirements of his own speciality the advances, so far as they are at all applicable, which have extended the sphere of general surgery.

This comparatively slow advance of recent years is not due to there being nothing left to find out. It is not due to the interest in the special subject of ophthalmology being for the time diverted to branches which seem to offer more scope for research. There is, at least, no dearth of writings on specially

ophthalmological topics. It is not due to the absence of experimentation with new therapeutic agents. New drugs are constantly appearing which, if they do not succeed in doing a tithe of what is claimed for them, at least raise one's hopes for the time being. Not a few books devoted exclusively to a consideration of new remedies have appeared quite recently. Even in connexion with eye diseases, however, for which it should be possible to estimate the results of treatment, in many cases at least, more fully than for other less accessible organs, there is so often shown the same absence of critical insight into the difficulties of doing so which seems to characterize the majority of writers on therapeutics. Notwithstanding valuable discoveries in pathology, we make, it must be admitted, but slow progress in treatment.

The problems which, for us, await solutions are in point of fact numerous enough. But they are for the most part such as seem only likely to be solved when our knowledge of general pathology and of medicine have reached a stage or two further than they have at present. We do not even know yet what it is that determines the selection of certain organs and tissues for the manifestation of certain definite pathological processes. We are consequently and necessarily empirics as regards the manner in which we may try to supplement purely local by more general means of treatment.

But if we cannot to-day check the progress of an optic atrophy, or treat, on anything but empirical lines, a choroiditis; if we cannot prevent the development of a cataract or of a detachment of the retina; if we cannot even cure a conjunctivitis, we may surely hope that some day it will be possible to do all these things, and more.

Because, gentlemen, if our progress is slow it is none the less real. We may have need to be critical—even sceptical—in the present, in view especially of the flood of literature, from which we must separate the chaff from the wheat. But we can afford to be—we have reason to be—optimistic as to the future.

My own predilections have always been for surgery. There is so much pleasurable interest to be got from watching the effect of surgical interference of any kind, so little satisfaction in cases for which one has to rely upon medical treatment alone, that one is apt to forget that the laborious, patient investigations of recent times may, in the perhaps not distant future, turn the tables on surgery. We may some day have to admit that many conditions for which surgical measures seem the only suitable ones at present will allow of rational, successful treatment on medical lines alone. Much of our treatment at the present time may even in the light of future knowledge be of a nature which, if it were then adopted, would be little less than criminal. I may illustrate this by a true story from the past. About 100 years ago there lived a king of one of the most enlightened European States who was in the habit of being periodically bled. One day his physician finding him suffering more than usual from the discomfort of a too generous diet, whipped out his lancet and relieved him of the superfluous quantity of his vital fluid. Next day the monarch's arm was painful and swollen. Phlebitis followed, which in a few days proved fatal. The physician's attentions were rewarded by a pension of £800 a year, to be continued to his son and his grandson. The grandson is still living in enjoyment of that pension.

There are many topics which, though they do not present the interest of novelty, may be discussed with more or less profit at a meeting like this. Formal discussions, with the necessary limitations imposed upon the time allowed for each member to speak, do not often, it is true, lead to much. Yet they sometimes help to lift one out of the groove of self-satisfaction. Those of us who have had an extensive personal experience are apt to think that the only "sensible" men are those who hold the same views as ourselves. But even in our little branch of science it is surely our duty "to prove all things and hold fast to that which is good."

The subjects for discussion at this meeting are ones on which we must all have some ideas, and, as we are not overloaded with papers on other matters, we may hope that others than those who have already intimated their intention of doing so may take part in them. You will see from the programme for the day that we have departed from the usual custom of beginning with the regular discussions. This was necessary, as the gentlemen who are to open the discussions are unable to do so before to-morrow and the day following. The papers have therefore to be taken first, but before calling for the first paper on the list, I have, as your chairman, to offer, in your name and in my own, a special welcome to the distinguished foreign colleague whom we are happy to see