THE EMPLOYMENT OF SILVER WIRE TO BRIDGE THE GAP AFTER RESECTION OF A PORTION OF THE LOWER JAW.

By SINCLAIR WHITE, M.Ch., F.R.C.S.Eng.,
SENIOR SURGEON TO THE ROYAL INFIRMARY, SHEFFIELD.

THE very distressing deformity, the complete loss of masticatory power, and the constant dribbling of saliva which follow the removal of a considerable segment of the lower jaw, are only too well known, and constitute a serious objection to this operation. The desire to overcome these disabilities prompted me in the following case to bridge the hiatus with silver wire, and the gratifying result obtained must be my excuse for sending a report of it for publication.

H. H., aged 9 years, was admitted to the Sheffield Royal Infirmary for a tumour of the lower jaw which had been growing for nearly two years. The growth involved the alveolar border and upper half of the left side of the body of the mandible from the site of the first molar tooth backwards for about 2 in. It had obviously started in the interior of the bone and grown upwards. Two enlarged glands could be felt in the submaxillary triangle. A small portion of the tumour was removed and examined by Professor Beattie, who reported it to be a small round-celled sarcoma.

In view of the pathological finding it became necessary to

Fig. 1.—Tumour and portion of lower jaw resected.

resect widely the involved area of bone, and to remove the lymphatic glands in the submaxillary triangle.

Free exposure was secured by carrying an incision from the left angle of the mouth downwards and backwards to the neck. The bone was divided in front at the level of the left canine tooth, and behind just above the junction of the body with the left ramus, and the tumour together with 2 in. of the body of the jaw, as shown in Fig. 1, removed. The periosteum covering

Fig. 2.—Diagram to show position of wire. A = Silver wire.

the lower border of the excised segment was carefully preserved. The resected surfaces of the lower jaw were pierced with a drill to the depth of $\frac{3}{8}$ in. The drill hole in the body was horizontal, and placed near its lower margin, so as to miss the teeth roots; that in the ramus was vertical and somewhat posterior to the mandibular foramen. The ends of a suitable length of stout silver wire were jammed tightly into the drill holes, and the wire completely covered by suturing together the mucous membranes of the cheek and the floor of the mouth over it. The diagram (Fig. 2) indicates the position and curve of the wire.

A small drain tube was placed in the neck end of the wound and retained for forty-eight hours, and the mouth was rinsed frequently with hydrogen peroxide solution. A little pus formed in the track of the tube, but the wound in the mouth healed quite kindly.

At the present time, except for the skin scar, there is, as the photograph (Fig. 3) shows, absolutely no external deformity. He can open his mouth almost to the full, and

Fig. 3.—Patient after recovery from the operation.

when the jaws are closed the teeth on the right side meet accurately those in the corresponding side of the upper jaw. He is able to bite soft things, and has to be restrained from attempting greater masticatory feats.

I think it is probable that new bone may form around the wire from the preserved periosteum and still further strengthen the mandibular arch.

Unfortunately the majority of cases in which wide resection of the lower jaw is called for are not likely to benefit by the adoption of this procedure. To be successful the wire should be completely buried in healthy tissue, and the wound should heal kindly. These conditions are seldom, if ever, attainable in operations for epithelioma, which is much the most common lesion for which resection is practised.

A CASE OF "GIANT FEET."

By HOWARD STEVENSON, B.A., M.B., Ch.B.R.U.I.,
F.R.C.S.I.,

SURGEON, CHILDREN'S DEPARTMENT, ULSTER HOSPITAL, BELFAST;
ASSISTANT TO THE PROFESSOR OF SURGERY, QUEEN'S
UNIVERSITY, BELFAST.

THE subject of these illustrations is a boy of 4 years of age, who came under my notice early this year.

The history, as given by the mother, was as follows: At birth he was apparently a normal child, with the exception of the deformity of the toes and some enlargement of the ball of the left foot. At the age of 6 months the child was vaccinated, and about the same time the lumps in the leg and groin appeared. She first noticed the swelling in the left groin, and thought the child was ruptured. Gradually the enlargement extended towards the armpit, the arms at the same time getting thinner.

Present condition: The two inner toes of the right foot are normal in development, the three outer very much enlarged and resembling small tubers; they are acutely dorsiflexed, and arise from a much thickened sole pad. The left foot is even more deformed, only the inner toe

escaping, and the characteristics described on the right side are more exaggerated. The left leg, thigh, groin, and infrascapular regions are sites of large fatty tumours, while a single large one is also present on the right buttock. When the child is standing there is well-marked lordosis, due apparently to the weight of the tumour in the groin, and possibly to some intra abdominal growths.

Notwithstanding the condition of its feet, the child is able to run about, play football and other games when shod in a pair of boots resembling horse boots.

The skin is everywhere normal in appearance, and a section from one of the tumours had the ordinary characteristics of a lipoma, and presented no evidence of angioma structure. Dr. Rankin kindly took an x-ray photograph, and this showed that there was no bony malformation.

THE third edition of the Hungarian *Pharmacopoeia* will come into force on January 1st, 1910.

THE SCIENCE COMMITTEE OF THE British Medical Association.

REPORT CXVIII.

SOME OBSERVATIONS ON A TWENTY-FOUR HOURS' WALKING RACE.

BY

F. COOK, B.Sc.; E. G. SCHLEISINGER, B.Sc.;

AND

A. H. TODD, B.Sc.

(From the Physiological Department, Guy's Hospital.)

A TWENTY-FOUR HOURS' WALKING RACE, promoted by the Blackheath Harriers, took place at the Stadium, Shepherd's Bush, on September 17th and 18th, 1909. By the courtesy of the secretary and of the club's medical officer, Dr. Munro, we were enabled to make some observations on the physical condition of the competitors. Inquiries were also made concerning their age, mode of training, diet, costume, and the time and nature of any discomfort experienced during the race. Circumstances prevented our making a complete examination of every man before and after the contest, so that the results lack the exactitude of a laboratory experiment; it is interesting, however, to compare them with the many data that have been published within recent years concerning more strenuous and less prolonged feats of endurance, such as running, cycling, and rowing races. For these reasons the observations made are here recorded.

1902

CONDITIONS.

The men were allowed to bring with them attendants, by whom they were massaged, fed, and cared for generally. The race began at 5 p.m., and continued until the same time next day. The weather conditions were as follows:

Temperature.

	Dry Bulb.	Wet Bulb.
September 17th, 6 p.m. ...	60.3°	57.0°
September 18th, 7 a.m. ...	56.4°	54.8°
" " 1 p.m. ...	62.3°	55.1°
" " 6 p.m. ...	59.2°	54.3°

On the second day there were five hours of sunshine. There was little wind throughout the race, and the cinder track was considered by the competitors to be in perfect condition.

The greatest distance covered was 127 miles; 24 of the 50 starters walked over 100 miles.

OBSERVATIONS.

1. Clothing.

Special leather boots were worn in the majority of cases in preference to shoes; besides shorts, long pants or stockings were generally worn. Most of the men had on vests and sweaters, and some wore hats, especially during the day. Others were more scantily clothed. The weight of clothing carried was therefore considerably less than that of everyday attire.

The advantages of boots were apparently: (a) The fixation of the ankle-joint, thus obviating the strain on the flexors and extensors of the foot, which are of no great assistance in walking; and (b) the avoidance of excessive friction and consequent blisters. All the men who were asked said that pants were worn for warmth only, not for support.

2. Cardiac Condition.

(1) *Pulse.*—The average pulse-rate of competitors before the race was from 70 to 80; two were of low frequency (47, 54) and two of high (112, 128), the latter being apparently due to excitement. At the end of the race the pulses were, as a rule, below 100. One competitor, however, whose pulse was 128 at the start, showed a frequency of 160 at the finish. In another the rate was only 52.