had had a good night. The tongue was clean; the temperature was 97.4°, the pulse 100, and the respirations 16.

On February 6th the pulse was 84, the temperature 98.6°, and the respirations 16; she had had a good night.

On February 7th the pulse was 74 and the temperature 97°; she had again had a good night and was feeling better.

From this time on her recovery was uninterrupted and uneventful. I regret to say, however, that the serum has not fulfilled expectations as far as preventing necrosis is concerned. There is complete ectropion of the upper eyelid, so that later on some plastic operation will be necessary; all the tissues superficial to the cartilage seem to be gone in the upper eyelid; but that is comparatively a small consideration, and the patient is to be congratulated on being alive at all. As to what influence Sclavo's serum had in bringing about recovery, it is impossible to say. I was unable to note anything that could be called a marked reaction after its use. Cases occasionally got well spontaneously in the old days, and it is just possible that this case got well simply *post hoc sed non propter hoc*. On the other hand, when we take into account the favourable results obtained in Italy with the serum, and the few cases recorded in this country, I think that we are justified in hoping that a remedy for this dread disease has at last been found.

Photograph eight weeks after commencement of disease showing amount of ectropion after complete cicatrization: A, upper ciliary margin; B, swollen conjunctiva, ocular; C, everted inner surface of upper eyelid; D, outer end of cavity left by separation of slough; E, area from which necrosis of lower eyelid separated, showing oedematous lower eyelid above it.

Mr. Bowlby and Dr. Andrewes of St. Bartholomew's Hospital, in their report of a case in the BRITISH MEDICAL JOURNAL of February 11th, state that the oedema increased after the use of Sclavo's serum in both the cases treated by them, and they attribute this to the liberation of an intercellular toxin from the disintegrated bodies of the bacilli. These dead bacilli they found chiefly enclosed in leucocytes. In my case there was also more swelling after the first dose of serum. Compare this with the claim, No. 5, of Dr. Legge that it quickly arrests the extension of the oedema. Mr. Bowlby and Dr. Andrewes also state, in connexion with their second case, that nineteen hours after an injection of 40 c.cm. of Sclavo's serum, fluid taken from the vesicles was sterile as far as anthrax bacilli was concerned. This is not in accordance with my observations, as I have shown that fluid from the vesicles after the administration of 70 c.cm. in all, and three days after the initial dose of 20 c.cm., proved virulent when tested physiologically.

In conclusion, I would mention that large initial doses of 50 to 60 c.cm. afford the best results, but whether the new treatment will stand the test of time or not remains to be seen. In the present state of our knowledge, it does not seem justifi-

able to treat a case of cutaneous anthrax with serum alone and without excision, provided that the pustule is anatomically situated in such a way as to render excision possible. Better to wait for cases such as this, where excision was impossible or practically so, than to willingly omit any procedure which is calculated to save life.

REFERENCE.

¹ See Dr. Legge's Milroy Lectures, BRITISH MEDICAL JOURNAL, 1905, for fuller particulars.

A SPOON-ELEVATOR FOR RAISING THE PELVIC FLOOR.

By J. LYNN THOMAS, C.B., F.R.C.S.,
Cardiff.

DR. GARSON, in 1878, made an important communication to the *Edinburgh Medical Journal* upon the "displacement of bladder and peritoneum by distension of the rectum." Soon afterwards the suprapubic operation for removal of vesical calculi was resuscitated, and for a time Petersen's bag enjoyed great popularity for lifting the base of the bladder by distending the rectum; it was also used in operations for removal of vesical tumours, but before long it was discovered that its use was not devoid of danger, and that it also caused increased congestion of the bladder by its pressure on all the blood vessels of the rectum.

Owing to the recent successful introduction of the operation of removal of the prostate by the suprapubic route, the surgeon has often wished he could supplement fingers in the rectum by an instrument which would answer the same purpose, although Dr. Eugene Fuller is able to perform his prostatectomies with marked success by simply pushing his closed fist against the perineum. Nevertheless, many surgeons feel that it is necessary to raise the prostate by fingers in the rectum, either the operator's or those of a skilled assistant, in order that improved conditions may be obtained for rapid enucleation of the gland; on one occasion I had the privilege of seeing Dr. Proust assist Professor Tuffier under such circumstances, but such skilled assistance is rarely available.

Messrs. Weiss have made for me a spoon which answers this purpose admirably. The illustration shows its chief mechanical points. The spoon is of the sixteenth-seventeenth century shape, and is 4 cm. across at its widest part: the spoon and its straight handle which are inserted into the rectum are made to imitate the length and the breadth of two long and broad fingers inserted into the rectum. The crank handle is made for the purpose of having the instrument conveniently at hand for manipulation in front of the patient's thigh. The curved processes, one on each side of the handle, are for the purpose of preventing the instrument rolling between the patient's thighs when the surgeon is not using it.

As soon as the bladder is opened the spoon is passed into the rectum and the patient's thighs are approximated; under these conditions it will remain *in situ* and quite out of the way. The surgeon can perform all the necessary manipulations without infecting his hands, and the instrument answers the purpose better, I think, than the fingers of an assistant in the rectum, because the operator acts directly on the spoon: he can push it up or draw it down to the exact position at which he requires the rectal counter-pressure; it can also be tilted to either side so that elevation of any particular area of the base and posterior wall of the bladder can be regulated to a nicety, for every movement of the handle is transmitted directly to the spoon, as it is specially made in one solid piece for that purpose. I have used it in three prostatectomies and in one case of evacuation of blood clots completely filling the bladder. In the latter case the clot was cleared out through a suprapubic wound with great ease and rapidity by means of the spoon in the rectum and suprapubic massage of the abdomen.

In a case of prostatectomy upon a friend of Dr. W. T. Edwards, J.P., I was able to remove forty-seven calculi from a large post-prostatic pouch which I could not otherwise reach with my fingers.

In one of the other cases there was a calculus weighing 2 oz., and its delivery was facilitated by using the spoon in the rectum.

In operations upon the female pelvis I introduce the spoon into the vagina and let it remain *in situ* until required for lifting either the uterus, Douglas's pouch, or bladder. The particular point at which lifting of the pelvic floor is required is easily obtained.

In a case of coeliotomy for soft, friable, and separate malignant growths of both ovaries secondary to carcinoma of the stomach, I found it very useful as the tumours would not withstand any traction upon them for delivery.

The face of a hysterectomy or of an ovarian stump can be easily manipulated so as to give a better view for its inspection, by means of the spoon in the vagina.

The reason that I record my short experience of the uses of the spoon is that Messrs. Weiss informed me that they had sold several of them after I referred in a P.S. to a paper on prostatectomy to the fact that I had used such an instrument. The instrument which I had used at that time, and which has been sold, is only three centimetres wide, but in practice I soon found that it was not broad enough for cases of enlarged cervix uteri, or of a very big prostate to hold them firmly during lifting of the pelvic floor, as they had a tendency to slip over to one side. The instrument which is described above is four centimetres wide and is more satisfactory on account of its increased breadth.

AN EXPERIMENTAL INQUIRY INTO THE INFECTION OF OPERATIVE WOUNDS, FROM THE SKIN, THE BREATH, AND THE AIR.*

By J. RUPERT COLLINS, M.D., B.Ch. (UNIV. DUB.).

As a natural prelude to the investigation as to the sterilization of the skin, was the determination of the germicidal power of the chemicals to be used. There exists a considerable literature upon this point, but, as the published results differ somewhat widely, I thought it preferable to simply accepting without question the observations of others to make some observations myself, if only for my own enlightenment. The method adopted was that of Drs. Rideal and Walker¹ with two modifications.

1. It was not feasible to maintain the temperature of the laboratory here at a constant.

2. One minute in place of half a minute elapsed between the inoculation of the various tubes.

The following was found to be the strength required of the chemicals named in order to have a sterilizing action upon a twenty-four hours culture of *staphylococcus pyogenes aureus* in broth incubated at blood heat, and taken from an agar-agar slant culture obtained from Král of Prag:

Antiseptic.	Period of Action.		Carbolic Acid Coefficient.
	One Minute.	Five Minutes.	
Carbolic acid	1 in 50	1 in 100	1
Lysol	1 in 100	1 in 100	1
Cyllin (medical)	1 in 600	1 in 1,250	12.5
Perchloride of mercury ...	1 in 2,500	—	50
Mercuric potassium iodide	1 in 4,000	1 in 8,000	80
Formalin	—	1 in 30	0.3

In attaining these results over 250 experiments were made.

It is now established that the resistance of the same micro-organism varies considerably in different cultures, but that the ratio between the different antiseptics as to germicidal action remains almost a constant in each culture. Recognition of this fact by bacteriologists has led to the general adoption of a carbolic acid coefficient as the unit of measurement of germicidal power.

In my own experiments the mercuric potassium iodide was

found to vary considerably in germicidal power. This may have been due to some technical error, but I cannot say where such arose. Fresh tabloids, supplied by Messrs. Burroughs Wellcome, were used and the dilutions were carefully made.

Klein² has clearly shown that reasoning from the actions of a germicide in a broth culture to the same action in the tissues and body secretions is fallacious. I therefore repeated the above experiments upon a milk culture of the same strain of *staphylococcus pyogenes aureus*, selecting milk as a medium most nearly approximating to the conditions present in a wound made through the skin and subcutaneous tissues. Bovine milk was that used.

Antiseptic.	Period of Action.		Carbolic Acid Coefficient.
	One Minute.	Five Minutes.	
Carbolic acid	1 in 20	1 in 60	1
Cyllin (medical)	1 in 250	1 in 800	13.3
Perchloride of mercury ...	1 in 1,000	—	50
Potassium mercuric iodide	—	1 in 500	8.3

For the purpose of getting rid of the natural grease of the skin many different chemicals—alcohol, ether, turpentine, benzine, xylol, potash soap, and soda soap—are employed by surgeons, and each chemical has its advocates.

Suppose, now, we take some lanoline as approximating fairly close to the human skin grease, and place a piece the size of a pea in a test tube containing one of these chemicals, what happens?

Alcohol (cold) ...	No perceptible effect.
" (hot) ...	Partial solubility.
Ether ...	Solubility takes place gradually.
Xylol ...	Ditto.
Benzine ...	Ditto.
Turpentine ...	Very little effect.
Potash soap ...	Scarcely any effect in the cold; in hot water complete emulsification is almost instantaneous.
Soda soap ...	Similar effects, but less rapid.

The second test, perhaps, conforms more nearly to the natural conditions. A thin layer of lanoline was smeared over the finger, which was then dabbled in one of these fat removers. Ether, ether and alcohol in equal parts, and the soaps in hot water in each case removed the fat in two minutes, so that no grease mark was left by the imprint of the fingers upon paper. This test was not applicable to xylol, benzine, and turpentine.

Now, although the lanoline in mass is different from the condition present in the skin, yet there is a sufficient analogy to justify us in drawing conclusions of practical value in reference to the relative activity which each respective chemical may be expected to exert upon the skin.

I now proceeded to apply these results to the disinfection of the hands, using solutions somewhat stronger than was found to be necessary as a germicide to *staphylococcus pyogenes aureus* in the test-tube, but weaker than I think is generally employed in surgery. The first experiment was a confirmation of that recorded in the close of my previous paper.³ After cleansing the hands with lysol, 1 in 100, for five minutes, and then in perchloride of mercury, 1 in 1,000, for one minute, the hands were dried with a sterilized cloth and then immersed in a litre of sterile nutrient peptone broth. The entire broth on being incubated remained sterile. A similar experiment, with the same result, was performed, using biniodide of mercury, 1 in 1,000, for five minutes, followed by carbolic acid, 1 in 40, for one minute.

It was pointed out to me, however, that though this test proved that the surface of the skin was sterile, yet it did not prove that it would remain so if subjected to friction such as takes place at an operation. At the annual meeting at Oxford last year many speakers enforced the same remark. To investigate this point I adopted the plan of using marble dust or sea sand, in order to obtain friction, as described by Dr. Engel⁴ and other German observers. The method pursued was as follows:

After cleansing and trimming the finger-nails, the hands were first thoroughly scrubbed with a nail-brush, soft soap and hot water containing some antiseptic for ten minutes, three changes of water being used. They were next soaked in a different antiseptic for two minutes. Finally, the hands were dried with a sterilized cloth and plunged forthwith into a jar containing a litre of nutrient peptone broth with a layer of marble dust or sea sand at the bottom. The hands were now vigorously scrubbed in the nutrient broth with the marble dust for five