In a case of prostatectomy upon a friend of Dr. W. T. Edwards, J.P., I was able to remove forty-seven calculi from a large post-prostatic pouch which I could not otherwise reach with my fingers.

In one of the other cases there was a calculus weighing 2 oz., and its delivery was facilitated by using the spoon in the rectum.

In operations upon the female pelvis I introduce the spoon into the vagina and let it remain *in situ* until required for lifting either the uterus, Douglas's pouch, or bladder. The particular point at which lifting of the pelvic floor is required is easily obtained.

In a case of coeliotomy for soft, friable, and separate malignant growths of both ovaries secondary to carcinoma of the stomach, I found it very useful as the tumours would not withstand any traction upon them for delivery.

The face of a hysterectomy or of an ovarian stump can be easily manipulated so as to give a better view for its inspection, by means of the spoon in the vagina.

The reason that I record my short experience of the uses of the spoon is that Messrs. Weiss informed me that they had sold several of them after I referred in a P.S. to a paper on prostatectomy to the fact that I had used such an instrument. The instrument which I had used at that time, and which has been sold, is only three centimetres wide, but in practice I soon found that it was not broad enough for cases of enlarged cervix uteri, or of a very big prostate to hold them firmly during lifting of the pelvic floor, as they had a tendency to slip over to one side. The instrument which is described above is four centimetres wide and is more satisfactory on account of its increased breadth.

AN EXPERIMENTAL INQUIRY INTO THE INFECTION OF OPERATIVE WOUNDS, FROM THE SKIN, THE BREATH, AND THE AIR.*

By J. RUPERT COLLINS, M.D., B.Ch. (UNIV. DUB.).

As a natural prelude to the investigation as to the sterilization of the skin, was the determination of the germicidal power of the chemicals to be used. There exists a considerable literature upon this point, but, as the published results differ somewhat widely, I thought it preferable to simply accepting without question the observations of others to make some observations myself, if only for my own enlightenment. The method adopted was that of Drs. Rideal and Walker¹ with two modifications.

1. It was not feasible to maintain the temperature of the laboratory here at a constant.

2. One minute in place of half a minute elapsed between the inoculation of the various tubes.

The following was found to be the strength required of the chemicals named in order to have a sterilizing action upon a twenty-four hours culture of *staphylococcus pyogenes aureus* in broth incubated at blood heat, and taken from an agar-agar slant culture obtained from Král of Prag:

Antiseptic.	Period of Action.		Carbolic Acid Coefficient.
	One Minute.	Five Minutes.	
Carbolic acid	1 in 50	1 in 100	1
Lysol	1 in 100	1 in 100	1
Cyllin (medical)	1 in 600	1 in 1,250	12.5
Perchloride of mercury ...	1 in 2,500	—	50
Mercuric potassium iodide	1 in 4,000	1 in 8,000	80
Formalin	—	1 in 30	0.3

In attaining these results over 250 experiments were made.

It is now established that the resistance of the same micro-organism varies considerably in different cultures, but that the ratio between the different antiseptics as to germicidal action remains almost a constant in each culture. Recognition of this fact by bacteriologists has led to the general adoption of a carbolic acid coefficient as the unit of measurement of germicidal power.

In my own experiments the mercuric potassium iodide was

found to vary considerably in germicidal power. This may have been due to some technical error, but I cannot say where such arose. Fresh tabloids, supplied by Messrs. Burroughs Wellcome, were used and the dilutions were carefully made.

Klein² has clearly shown that reasoning from the actions of a germicide in a broth culture to the same action in the tissues and body secretions is fallacious. I therefore repeated the above experiments upon a milk culture of the same strain of *staphylococcus pyogenes aureus*, selecting milk as a medium most nearly approximating to the conditions present in a wound made through the skin and subcutaneous tissues. Bovine milk was that used.

Antiseptic.	Period of Action.		Carbolic Acid Coefficient.
	One Minute.	Five Minutes.	
Carbolic acid	1 in 20	1 in 60	1
Cyllin (medical)	1 in 250	1 in 800	13.3
Perchloride of mercury ...	1 in 1,000	—	50
Potassium mercuric iodide	—	1 in 500	8.3

For the purpose of getting rid of the natural grease of the skin many different chemicals—alcohol, ether, turpentine, benzine, xylol, potash soap, and soda soap—are employed by surgeons, and each chemical has its advocates.

Suppose, now, we take some lanoline as approximating fairly close to the human skin grease, and place a piece the size of a pea in a test tube containing one of these chemicals, what happens?

Alcohol (cold) ...	No perceptible effect.
" (hot) ...	Partial solubility.
Ether ...	Solubility takes place gradually.
Xylol ...	Ditto.
Benzine ...	Ditto.
Turpentine ...	Very little effect.
Potash soap ...	Scarcely any effect in the cold; in hot water complete emulsification is almost instantaneous.
Soda soap ...	Similar effects, but less rapid.

The second test, perhaps, conforms more nearly to the natural conditions. A thin layer of lanoline was smeared over the finger, which was then dabbled in one of these fat removers. Ether, ether and alcohol in equal parts, and the soaps in hot water in each case removed the fat in two minutes, so that no grease mark was left by the imprint of the fingers upon paper. This test was not applicable to xylol, benzine, and turpentine.

Now, although the lanoline in mass is different from the condition present in the skin, yet there is a sufficient analogy to justify us in drawing conclusions of practical value in reference to the relative activity which each respective chemical may be expected to exert upon the skin.

I now proceeded to apply these results to the disinfection of the hands, using solutions somewhat stronger than was found to be necessary as a germicide to *staphylococcus pyogenes aureus* in the test-tube, but weaker than I think is generally employed in surgery. The first experiment was a confirmation of that recorded in the close of my previous paper.³ After cleansing the hands with lysol, 1 in 100, for five minutes, and then in perchloride of mercury, 1 in 1,000, for one minute, the hands were dried with a sterilized cloth and then immersed in a litre of sterile nutrient peptone broth. The entire broth on being incubated remained sterile. A similar experiment, with the same result, was performed, using biniodide of mercury, 1 in 1,000, for five minutes, followed by carbolic acid, 1 in 40, for one minute.

It was pointed out to me, however, that though this test proved that the surface of the skin was sterile, yet it did not prove that it would remain so if subjected to friction such as takes place at an operation. At the annual meeting at Oxford last year many speakers enforced the same remark. To investigate this point I adopted the plan of using marble dust or sea sand, in order to obtain friction, as described by Dr. Engel⁴ and other German observers. The method pursued was as follows:

After cleansing and trimming the finger-nails, the hands were first thoroughly scrubbed with a nail-brush, soft soap and hot water containing some antiseptic for ten minutes, three changes of water being used. They were next soaked in a different antiseptic for two minutes. Finally, the hands were dried with a sterilized cloth and plunged forthwith into a jar containing a litre of nutrient peptone broth with a layer of marble dust or sea sand at the bottom. The hands were now vigorously scrubbed in the nutrient broth with the marble dust for five

minutes, the latter being worked down into the interstices round the edges of the nails. At the end of five minutes the hands were taken out, the cover replaced upon the jar and the whole incubated. In preparing it, the entire jar which contained broth and sand was sterilized *in toto*. The risk of atmospheric infection during the period in which the hands were being washed in the broth was very greatly reduced by the following device. A cloth was prepared with two holes round the edges of which was hemmed an elastic band. The cloth fastened by string was stretched over the top of the jar underneath its glass cover, and was sterilized in position. In use, as an assistant removed the glass cover, the hands were quickly thrust through the holes in the cloth, the elastic edges of which clinging tightly to the arms prevented the entrance of infection while permitting free washing movements of the hands. When removing the hands, the cord securing the cloth to the top of the jar was cut and as the hands with cloth attached were quickly removed from the jar, the glass cover, which had been lying face downwards upon a sterilized cloth during the operation, was immediately replaced by the assistant. The jar with its contents was then incubated, and subsequently subcultures were also made in broth and on agar-agar. In the first of these experiments, as it was expected that a considerable infection might take place, the broth was stiffened by the addition of 10 per cent. gelatine in the manner described by Engel,⁵ in order that it might be possible to count the number of colonies present. As, however, this experiment proved sterile, and the presence of the gelatine prevented incubation in the warm oven, with the exception of Experiment VI it was not used again.

No. of Experiments.	Mode of Disinfecting the Hands.	Mode of Testing the Hands After Disinfection.	Result.
I.	10 minutes in hot cyllin lotion, 1 in 200, together with soft soap, nail-brush, and Schleich's marble soap; 5 minutes in perchloride of mercury, 1 in 500	Scrubbed with marble dust in peptone broth; the broth incubated	Sterile.
II.	10 minutes in hot cyllin lotion, 1 in 400, together with soft soap, nail-brush, and Schleich's marble soap; 2 minutes in perchloride of mercury, 1 in 1,000	Ditto	Septic.
III.	10 minutes in hot cyllin lotion, 1 in 200, together with soft soap, nail-brush, and Schleich's marble soap; 2 minutes in perchloride of mercury, 1 in 500	Ditto	Sterile.
IV.	10 minutes in hot biniodide of mercury lotion, 1 in 2,000, with soft soap, nail-brush, and Schleich's marble soap; 2 minutes in cyllin, 1 in 100	Ditto	Septic.
V.	10 minutes in hot biniodide of mercury lotion, 1 in 2,000, with soft soap, nail-brush, and Schleich's marble soap; 1 minute in methylated ether; 2 minutes in carbolic acid, 1 in 20	Ditto	Septic.
VI.	10 minutes in hot cyllin lotion, 1 in 400, with soft soap, nail-brush, and Schleich's marble soap; 2 minutes in perchloride of mercury, 1 in 1,000	Ditto (gelatine)	Sterile.
VII.	10 minutes in hot cyllin lotion, 1 in 600, with soft soap, nail-brush, and Schleich's marble soap; 2 minutes in biniodide of mercury, 1 in 2,000	Ditto	Septic.
VIII.	10 minutes in hot cyllin lotion, 1 in 600, with soft soap, nail-brush, and Schleich's marble soap; 2 minutes in carbolic, 1 in 40	Ditto	Septic.
IX.	10 minutes in hot cyllin lotion, 1 in 600, with soft soap, nail-brush, and Schleich's marble soap; 2 minutes in a 70 per cent. alcoholic solution of biniodide of mercury, 1 in 2,000	Ditto	Septic.
X.	10 minutes in hot cyllin lotion, 1 in 60, with soft soap, nail-brush, and Schleich's marble soap; 2 minutes in perchloride of mercury, 1 in 2,000	Ditto	Sterile.
XI.	Ditto	Ditto	Sterile.
XII.	10 minutes in hot cyllin lotion, 1 in 600, with soft soap, nail-brush, and Schleich's marble soap; sponged for half-minute in methylated ether and methylated alcohol equal parts; 2 minutes in perchloride of mercury, 1 in 2,000	Ditto	Sterile.
XIII.	Ditto	Ditto	Sterile.
XIV.	10 minutes in hot cyllin lotion, 1 in 600, with soft soap, nail-brush, and Schleich's marble soap; 2 minutes in carbolic acid, 1 in 40	Ditto	Sterile.
XV.	10 minutes in hot cyllin lotion, 1 in 600, with soft soap, nail-brush, and Schleich's marble soap; 1 minute in cyllin, 1 in 200	Ditto	Septic.
XVI.	10 minutes in hot cyllin lotion, 1 in 600, with soft soap, nail-brush, and Schleich's marble soap; 2 minutes in perchloride of mercury, 1 in 2,000	Ditto	Sterile.

Experiment XVI was not carried out on my own hands, but was performed by the sister of Ward VII, who, immediately previous to the experiment was engaged in the ordinary routine of ward work.

It now appeared to me certain that, with an adequate use of cyllin and perchloride of mercury, it was possible to obtain a degree of sterility of the skin which would stand the test of considerable maceration and friction. But there remained untouched the question of an ulterior infection of the skin surface by the secretion of the sweat and sebaceous glands. To determine this point it was clearly necessary to make these glands actively secrete into the nutrient medium; to this intent the following procedure was adopted:

At the end of the disinfecting process the hands were tested for five minutes, as before, in a jar containing a litre of broth with marble dust at the bottom. They were then transferred from this, one hand being inserted into another jar containing a small roll of lint enclosing some absorbent wool, inside which again was fused chloride of calcium.

In order to prevent evaporation of the secretions the top of the jar was covered with a piece of jaconette containing a hole edged with elastic. The arm passing through the hole was gripped tightly by it. In this way a serious error was avoided. Pilocarpine was then injected in equal doses into each arm. Each hand was kept in its respective jar for one hour. The jar containing the broth was incubated, and afforded evidence as to the septicity of the secretions. The other jar acted as a hygrometer, the amount of the secretion from the hand and wrist being estimated by weighing the roll containing the chloride of calcium at the beginning and at the end of the experiment.

In describing the results of these experiments, the first jar of broth containing marble dust, in which the hands were scrubbed, will be termed A. The second jar of broth, in which the hand remained for one hour after the injection of pilocarpine, will be termed B.

No. of Experiment.	Mode of Disinfecting the Hands.	Result of Jar A.	Dose of Pilocarpine.	Amount of Secretion.	Result of Jar B.
I.	10 minutes in cyllin, 1 in 600; 1 minute in methylated ether and alcohol; 2 minutes in perchloride of mercury, 1 in 2,000	Sterile	0.2 gr.	4.7 grams	Septic.
II.	10 minutes in cyllin, 1 in 600; 2 minutes in perchloride of mercury, 1 in 2,000	Sterile	0.3 gr.	5.2 grams	Septic.
III.	10 minutes in cyllin, 1 in 600; 1 minute in methylated ether and alcohol; 2 minutes in perchloride of mercury, 1 in 2,000	Sterile	0.2 gr.	3.4 grams	Septic.
IV.	20 minutes in cyllin, 1 in 600; 1 minute in methylated ether and alcohol; 2 minutes in perchloride of mercury, 1 in 2,000	Sterile	0.2 gr.	3.6 grams	Sterile.
V.	20 minutes in cyllin, 1 in 800; 2 minutes in perchloride of mercury, 1 in 2,000	Sterile	0.2 gr.	2.6 grams	Sterile.
VI.	20 minutes in cyllin, 1 in 800; 2 minutes in perchloride of mercury, 1 in 2,000	Sterile	0.2 gr.	3.8 grams	After four weeks a mould grew. Sterile.
VII.	20 minutes in cyllin, 1 in 800; 2 minutes in perchloride of mercury, 1 in 2,000	Sterile	0.2 gr.	5.1 grams	

At this point these experiments had to cease owing to the termination of my appointment to the hospital. Looking at the above results, it will be seen that only when the preliminary washing lasted twenty minutes did the hands prove to be absolutely sterile, also that they then proved sterile in every case except in Experiment VI. That this was no true exception the following facts, I think, show:—The jar remained apparently sterile for four weeks; the mould, penicillium glaucum, which then developed, produced a luxuriant growth in three days in a subculture; hence I think we may conclude that this was a case of accidental infection from some disturbance of the lid of the jar. I think that these experiments, though few, definitely prove that an absolute sterility of the skin is attainable, and to attain it a prolonged careful scrubbing is necessary in order to remove the outside cuticle. The notion that by increasing the strength of antiseptics the period required for disinfecting the hands can be shortened is erroneous.

To any one who may desire to repeat these experiments, I strongly recommend the hypodermic injection of

atropine 1/10 gr. with strychnine 1/10 gr. as alleviating the extremely disagreeable after-effects of the pilocarpine.

The reason why the combination of cyllin followed by perchloride of mercury gave better results than other antiseptics of the same germicidal power puzzled me for long, but I believe the explanation is given by the following experiment.

If one c.cm. of egg albumin be placed in each of a series of test tubes and to each 5 c.cm. of one of the following antiseptics be added the precipitated albuminates can be compared.

Again, if the contents of each tube be filtered and the filtrate tested by the albuminometre the complement of each of these values is obtained by thus determining the respective ratios of the soluble albuminates. If the precipitated albuminates be arranged in respect to their density the order is as follows:

1. Alcohol	70 per cent.
2. Perchloride of mercury	1 in 2,000
3. Carbolic acid	1 in 40
4. Bichloride of mercury	1 in 2,000
5. Cyllin	1 in 600

Naturally the best results would be obtained by the use of a primary antiseptic which disintegrated rather than condensed the albumin of the skin, followed by an antiseptic which formed a hard, firm albuminate upon the surface of the skin—qualities fulfilled by cyllin and perchloride of mercury respectively.

From the discussion at the Oxford meeting, and from the somewhat polemical writings on the same subject which subsequently appeared in the *BRITISH MEDICAL JOURNAL*,⁶ it would appear that the results obtained by bacteriologists in the laboratory did not quite accord with the experience of surgeons in the theatre, and my attention has been more particularly drawn to this point by Mr. Howell. The chief cause for this apparent discrepancy is, I believe, to be found in an insufficient appreciation of the necessity to remove the superficial part of the horny layer of the epidermis, which forms a porous coating to the skin, and of which the interstices are swarming with micro-organisms, as the first stage in any process of sterilization. The skin, after immersion in antiseptics with this layer unremoved, is fitly described as a whitened sepulchre. This was brought home to me in a very forcible manner in Nos. II and III of the last series of hand experiments. In each case Jar A. proved sterile, but Jar B. was not only septic but stank appallingly, the stench exceeding that of any necropsy, or, in fact, any other stench which it has been my lot to experience. I therefore believe that the important point now for determination is the best means of removing this porous layer rapidly and without irritation, and I am at present experimenting upon this question. Once this layer is removed the rest is easy, as we already possess many antiseptics reliable for surface sterilization.

Since the foregoing was written the following extract from the *Arch. per le Scienze Med.* appeared in the *EPITOME of the BRITISH MEDICAL JOURNAL* for January 21st, 1905. V. Tirelli, from the results of 409 horizontal and vertical sections of the human skin, concludes as follows:

(1) That in the skin of ordinary hospital in-patients and in those parts of it not subjected to traumatism or pressure, and only scantily supplied with hair, the germs are collected in the superficial strata of the corneous layer. Only exceptionally, as the result of accidental lesions, do they reach the deep strata. (2) That the epidermis and the hairs and cutaneous glands are proof against infection, which, however, can penetrate into the more peripheral part of the hair follicles. (3) That under such conditions the germs in the skin are accessible to disinfection, and the most important element in the success of the disinfection is the maceration and removal of the corneous strata of the skin. (4) That it is reasonable to hope that further study of the technique of disinfection will, in the near future, reveal a method which can be relied upon to give uniformly successful results in the disinfection of skin, at any rate in in-patients.

These conclusions, arrived at by an absolutely different method, almost exactly coincide with my own.

It may excite surprise that I have not employed more extensively alcohol, which is so strongly recommended by far the greater number of investigators. In the first place, alcohol not only causes severe damage to my skin, but affords me acute pain when using it. In the second place, it appears to me that alcohol can only act as a dehydrating agent and fix the surface tissues with their included bacteria. According to Mr. Leedham-Green, an ardent supporter of alcohol, it does not owe its value to its fat-removing, to its cleansing, nor even to its antiseptic powers so much as to its property of hardening and fixing the superficial cells of the epidermis.⁷ We might then expect that, as the skin becomes again hydrated and macerated during the course of an operation, the pro-

bability would be that the hands would again become septic; and this was found to be actually the case in Mr. Leedham-Green's instructive experiments. I entirely agree with him, however, that alcohol admirably prepares the way for the subsequent use of watery antiseptics, owing to its dehydrating action.

I now turn to the other possible causes of infection in operation wounds—namely, instruments, dressings, and the air of the operating theatre. Instruments and dressings can be satisfactorily sterilized by heat; there remains therefore for consideration aerial infection.

At the introduction of Listerism, the air as a source of contamination received a large amount of attention from surgeons, but with the disuse of the carbolic spray the atmosphere as a factor in the causation of sepsis came to be practically ignored. Recently interest has been roused in a particular form of aerial infection—that by the surgeon's breath.

I cannot depict more clearly the nature of aerial infection than by quoting the following passage out of Tyndall's book on *Floating Matter of the Air*.⁸

Reflecting on the whole of this, I conclude that the germs float through the atmosphere in groups or clouds, and that now and then a cloud specifically different from the prevalent ones is wafted through the air. The touching of a nutritive fluid by a bacterial cloud would naturally have a different effect from the touching of it by the sterile air between two clouds. But, as in the case of a mottled sky, the various portions of a landscape are successively visited by shade, so in the long run are the various tubes of our tray touched by the bacterial clouds; the final fertilization of them all being the consequence.

The variation in numbers of these bacterial clouds is so greatly affected by the numerous and varying determining factors present in different places, that I think it would be useless to try to determine the number of bacteria that might be expected to fall upon a wound in the course of an operation. I will merely mention that last year I performed one experiment for Dr. Ferguson on this point. During an operation performed in the hospital theatre, the number of bacteria falling upon an agar-agar plate worked out at 32 per square inch per hour.

It is difficult to conceive any plan by which the air of a room could be made and kept absolutely sterile throughout an operation. A very great deal can be done, however, by methods easily carried out to bring the air of a theatre much nearer to that desirable state than at present is usually the case. Consider for a moment what is the reason for the presence of organic particles floating in the air. This is due to the movements and concussions of large bodies against one another, combined with the action of air currents from various causes. These forces suffice to cause dust, which is mainly composed of organic matter, to rise in clouds in opposition to the force of gravity, eddying through space in response to the slightest disturbance of the temperature or movement of the atmosphere. Tyndall⁹ performed a very ingenious experiment to show this. A box was coated on its interior with glycerine. A concentrated beam of light was then passed through the box showing the air to be laden with floating matter. After three days of quiet the air was found to be optically pure, all floating particles having become adherent to the sides of the box. Now, I think that this experiment is a very important one to the surgeon.

I will cite one other experiment:

It has been shown by Carnally, Haldane, and Anderson¹⁰ that the disturbance of dust by stamping of the feet causes an enormous increase of organisms in the air, in one case from 11 to 150 per litre. They are of opinion, however, that the ordinary movements of even many persons in a room are not sufficient to produce any marked change in the number of organisms in the air. I submit, then, that the following details would form a rational method in order to arrive at a maximum of purity in the atmosphere of the operating theatre. The walls of the theatre should be kept moist, as bacteria are detached with difficulty from a moist surface. The theatre ought to contain the minimum of furniture. There ought to be as little running about as possible by the attendants during an operation. The air admitted to the theatre should be filtered through cotton-wool. Once an operation has begun no one should enter or leave the theatre until the completion of the operation. All spectators ought to be kept outside the theatre altogether, and observe the operation through a glass screen. They would thus be able to approach quite sufficiently close to the scene of operation without being subjected to the heat of the theatre, and with freedom to come and go without disturbing the operator. Finally, all those in the theatre—operator, assistants, anaesthetist, and nurses—should be clothed in a complete suit of sterilized material, including foot-gear.

It is extraordinary, considering the trouble an ordinary man will take to clothe himself appropriately before indulging in any game demanding even a moderate amount of physical exercise, that the surgeon will descend into a heated operating theatre and perform there an operation entailing extreme physical and mental effort without any change in his attire beyond the substitution of a linen overall for a coat. Surely the custom of wearing a sterilized linen suit would be not only more cleanly and comfortable, but also more hygienic for the surgeon himself.

I believe that the methods suggested would render the atmosphere materially purer, would exclude the worrying criticisms and suggestions of bystanders, and, by increasing the physical comfort of the surgeon, would insure the accomplishment of an arduous operation with the minimum expenditure of that most valuable commodity—energy.

Owing to Mikulicz of Breslau and Dr. Mendes de Leon, a great deal of attention has been centred on the question of the surgeon's breath as a source of wound infection. Now, in the first place, I may state that expired air is sterile, as proved by Tyndall¹¹ in 1869, and since confirmed by later observers. It is, therefore, only from solid particles discharged by a sputtering action that infection can occur from the breath. In order to ascertain the degree to which this is usually present, I made the following experiments upon twenty-five different people:

Two agar-agar plates were placed at 1 ft. and 2 ft. distance respectively from the individual's mouth, who then read aloud for ten minutes from a book at a distance. A third plate was exposed during the same period behind the reader to act as a control for the air of the room. The room was, as far as possible, kept absolutely still, and the person reading kept the head fixed in the same position. In the first place, I performed this experiment upon myself ten times, with the object of seeing whether the results were at all constant for the same individual.

No. of Experiment.	Control.	One Foot Distance.	Two Feet Distance.
I.	5	11	3
II.	10	13	11
III.	Overgrown	Overgrown	26
IV.	Overgrown	"	10
V.	Overgrown	13	4
VI.	"	15	Overgrown
VII.	4	15	"
VIII.	2	Overgrown	11
IX.	5	16	28
X.	Overgrown	Overgrown	31

On adding up each column some months after the experiments were completed I found the total to be:

Control.	One Foot Distance.	Two Feet Distance.
83 + 4 overgrown	83 + 4 overgrown	118 + 2 overgrown.

The identity of the totals for the first two columns was, of course, a mere coincidence and is of no import. But for Experiment IV there would have been a small increase in the total of the second column over that of the first. "Overgrown" in above and subsequent experiments does not imply that the plate was more than usually infected, but simply indicates that a rapidly-growing mould quickly overspread the surface of the agar-agar so as to render impossible a correct enumeration of the other colonies.

Of the twenty-five different individuals tested in the same way the results were as indicated in the table at the head of the next column.

By substituting for the cases in which the plates were overgrown the average number obtained from the remaining plates of the same series there are obtained the following numbers for comparison:

Comparative numbers: 289.04 : 588.75 : 238.23.

That is to say there was, on an average, a gain of 11.98 colonies on a plate $3\frac{1}{2}$ in. in diameter at 1 ft. distance as the result of reading for ten minutes in a loud voice, while at a distance of 2 ft. there was no increase. It was found that, as a rule, one to two thousand words were articulated during this time.

As the above individual experiments showed varying results, I thought it would be instructive to see what would be the effect of breathing into a space of sterile air, where the complication of bacterial clouds would not exist. Accordingly, with the aid of Mr. Lionel Colledge, I prepared a tin box with certain openings, but with its joinings, etc., airtight. In the

Individual Tested.	Control.	One Foot Distance.	Two Feet Distance.
M. D. ...	Overgrown	Overgrown	Overgrown
T. M. ...	"	13	"
S. A. ...	"	70	2
S. H. ...	"	6	"
S. B. ...	1	0	0
N. N. ...	7	Overgrown	Overgrown
A. C. ...	Overgrown	28	10
E. R. ...	8	16	Overgrown
A. C. ...	10	29	"
A. L. ...	Overgrown	Overgrown	11
L. C. ...	9	"	32
E. C. ...	12	14	7
M. D. ...	Overgrown	20	0
N. W. ...	0	31	12
K. F. ...	Overgrown	10	2
L. B. ...	4	16	4
L. W. ...	8	16	Overgrown
C. J. ...	Overgrown	3	5
N. G. ...	13	71	11
N. G. ...	25	3	7
N. W. ...	3	3	0
N. W. ...	53	01	31
G. M. ...	5	30	5
J. C. ...	8	8	9
M. S. ...	Overgrown	Overgrown	Overgrown
Total ...	185+9 overgrown	471+5 overgrown	162+8 overgrown

top of the box there was an opening, through which passed a glass tube packed with cotton-wool, so that filtered air only could enter the chamber. At one side there was an opening for the insertion and removal of a Petri dish, and on the opposite side a circular opening $1\frac{1}{4}$ in. in diameter for speaking into the box. When these two openings were closed the box was airtight, with the exception of the opening at the top, by which filtered air only could enter. A Petri dish, containing agar-agar, was then placed in the box, the two lower openings were closed, and the upper opening was plugged with cotton-wool and the whole box sterilized on three successive days for one hour. Before sterilization a fine piece of tape was attached to the lid of the Petri dish and passed up through the cotton-wool. By this means the lid could be removed and replaced when desired, while preserving the aseptic condition of the box. In use, a mouthpiece was fitted tightly by a rubber washer to the opening in the side, and the experimenter breathed, sang, or talked into the enclosed sterile space, measuring about 9 in. square. At the end of the experiment the lid was lowered down on to the dish again, the box opened and the dish removed and incubated. As the gaseous exchange was not active, four minutes' respiration with active vocalization into this small space was found to be as much as could be done with comfort.

The following were the results obtained:

Experimenter.	Mode of Experiment.	Result.
Self	Simple breathing	Sterile.
"	Singing loudly	1 Colony.
"	"	2 Colonies.
"	"Reciting"	1 Colony.
S. A.	Singing	Sterile.
"	Reciting	1 Colony.
M. S.	"	4 Colonies.
"	"	8 Colonies.
G. M.	"	Overgrown (lid was knocked off as the plate was being removed).

The three last experimenters were selected as having proved especially septic in the open series of experiments, yet the results obtained when an ordinary mouthpiece of a Clover's ether inhaler was worn pointed to the conclusion that the infection carried by the breath in ordinary speech is small.

It seemed to me, then, that a mask of the kind now to be described would suffice for those who require a mask at all.

An ordinary wooden chip pillbox is taken, the lid is detached, and a hole $\frac{5}{8}$ in. in diameter is drilled in the bottom. Two segments are now removed from the sides of the box to adapt it to the convexities of the upper and lower maxillae. The edges of the box are bound with lint to render it comfortable. The mask is now ready, and can be attached to the mouth by pieces of elastic passing from either side of the box to the ears. Such a mask is cool, comfortable, and, I believe,

efficient. The following experiments were performed with it, selecting again S. A. and M. S. as the operators.

Experimenter.	Control.	One Foot Distance.	Two Feet Distance.
S. A.	Without a mask 4 With a mask ... 8	25 12	Overgrown. "
"	Without a mask 5 With a mask ... overgrown	10 8	6 Overgrown.
"	Without a mask 2 With a mask ... 5	Overgrown "	5 10
M. S.	Without a mask overgrown With a mask ... 24	" "	25 4
"	Without a mask overgrown With a mask overgrown	" "	Overgrown. "
Self	Without a mask 1 With a mask ... 1	7 5	1 2
Total	Without a mask 12+2 overgrown With a mask 38+2 overgrown	42+3 overgrown 25+3 overgrown	37+2 overgrown. 16+3 overgrown.

The large number of plates which were overgrown somewhat mars the value of the results, yet it is seen that in the total of the cases without a mask there was a marked increase of contamination, while when the mask was worn there was no increase in the contamination of the plates read over beyond that of the control. The experiment without a mask was in each case performed first, hence probably the greater pollution of the controls when the mask was worn.

In regard to the use of gargles, the following was the result obtained by a strong gargle containing carbolic acid, chlorate of potash, and glycerine, on S. A.:

S. A.	Control.	One Foot Distance.	Two Feet Distance.
Before gargle	4	31	2
After gargle	5	28	3

In conclusion, I desire to express my obligations in the first place to Sister Hastie of Ward VII, without whose indefatigable and skilful help I could not have executed these experiments; to Mr. Lionel Colledge for his many valuable suggestions; to Professor O'Sullivan of Dublin University for his kindly aid with the bibliography of the subject; and, finally, to all those who, at considerable personal discomfort, took part in these experiments.

REFERENCES.

- ¹ Vide *Practitioner*, vol. lxxix, No. 413, p. 520. ² BRITISH MEDICAL JOURNAL, July 2nd, 1904. ³ Ibid., June 11th, 1904. ⁴ *Centralblatt für Bacteriologie*, xxiv, Bd. 1903, No. 1. ⁵ Ibid. ⁶ Loc. cit., October 8th, 15th, 20th, 1903. ⁷ *Sterilization of the Hands*, by Mr. Leedham-Green, pp. 65, 72. ⁸ *The Floating Matter of the Air*, by John Tyndall, second edition, p. 117. ⁹ Ibid., p. 51. ¹⁰ *Journal of Hygiene*, vol. 3, No. 4, p. 570. ¹¹ Loc. cit., p. 47.

KERATOSIS PALMARIS ET PLANTARIS IN FIVE GENERATIONS.

BY
F. H. JACOB, and ADAM FULTON,
M.D.Lond. M.B., B.Ch., B.A.O., R.U.I.

THERE lives in this town a man, J. W., who suffers from keratosis palmaris et plantaris in a marked degree.

In adjoining houses live three children and three grandchildren who have inherited the condition.

The youngest child affected is twenty months old, and the skin of the palms and soles, instead of being thin, pink, and shiny, is considerably thickened, white, and rough, like ground glass.

In the adults the horny epithelium is about $\frac{1}{8}$ in. thick and almost black. It presents, in lines corresponding to the normal folds of the skin, deep fissures which have at their base apparently normal epithelium and which, therefore, do not get sore, nor crack and bleed, as do those in acquired keratosis.

J. W. gives the following family tradition as to the origin

of the complaint. His great-grandmother, while returning on a troopship from India, was one day looking over the bulwarks and was startled by the unexpected jumping of a large fish out of the water. "A fish, a fish," she cried, and clapped her hands together. The child of which she was then pregnant was born with "fish scales" on its hands and feet, as were also the eleven other children to whom she subsequently gave birth. J. W. remembers his grandmother perfectly well, and vouches for the fact that both she and his mother suffered from this affection.

Unfortunately this hereditary condition is looked upon as a skeleton in the family cupboard, and our request for permission to take photographs of it was refused.

Appended is a full genealogy of the family:

TREATMENT OF SEROUS EFFUSIONS BY INJECTION OF ADRENALIN CHLORIDE.

BY
H. W. PLANT, and PATRICK STEELE,
B.A.Lond., M.R.C.S., L.R.C.P., M.B., Ch.B.Edin.,
House-Physician, House-Physician,
Queen's Hospital, Birmingham.

THIS method of treating serous effusions was first advocated by Barr, of Liverpool, in 1903.

In the first case, one of malignant disease of the pleura with haemorrhagic effusion, he used adrenalin injections empirically, and found the treatment entirely successful as regards the effusion. He then tried it on other cases, both pleural and peritoneal, with almost invariable success even in a case of pericarditis with effusion. In his pamphlet he states that the results obtained in cases of ascites from hepatic cirrhosis were not quite so good as in those resulting from tuberculous peritonitis and malignant growth. The only cases of ascites, with one exception, in which we have tried adrenalin have been those due to cirrhosis of the liver, and without exception the treatment has proved successful. In only one case have we had to inject the adrenalin more than twice, and in the majority of our cases one injection has proved sufficient.

In all cases in which we have tried the effects of the injection of adrenalin the following has been the method adopted:—As much as possible of the fluid was withdrawn by a two-way trochar and cannula, and through the cannula, still *in situ*, 1 drachm of adrenalin chloride (1 in 1,000), diluted to $\frac{1}{2}$ oz. with sterile water, was introduced by means of an exploring syringe. The cannula was then removed, the wound closed with wool and collodion, and, in abdominal cases, the abdomen was gently manipulated for five minutes and a binder firmly applied.

Appended is a summary of the cases in which we have tried this method:

Cirrhosis of Liver.

CASE I.—G. K., aged 54, wireworker. Admitted July 11th, 1904. Complaint, abdominal pain and swelling. Abdomen greatly distended and giving all the usual signs of fluid. There was an umbilical hernia, fairly easily reducible. Paracentesis abdominis was performed on the following dates:

July 14th. 20 pints drawn off: specific gravity 1009.