A Post-Graduate Lecture

ON

INTERMITTENT CLOSING OF CEREBRAL ARTERIES:

ITS RELATION TO TEMPORARY AND PERMANENT PARALYSIS.

BY WILLIAM RUSSELL, M.D., F.R.C.P. EDIN.,

PHYSICIAN AND LECTURER ON CLINICAL MEDICINE, ROYAL INFIRMARY, EDINBURGH.

GENTLEMEN,—I have been seeking a new word to express the kind of vessel action that I propose to speak to you about this afternoon, but I have failed in the search. Many of you must know the term "intermittent claudication," which has had a place in medical literature for about eighty years. It was introduced from veterinary medicine, where it was first used to describe recurring attacks of lameness or limping, with pain in the affected limbs, which occurred in a mare, and were due to obstruction to the arterial blood supply when the muscles were put in vigorous action. In human medicine the term has always carried with it the conception of loss of power as a result of the closing of arterial channels. A record of mere limping in a horse would be of no more interest to the veterinarian than to us; it was the *cause* of the limping that was of interest; it was this that led to the introduction of the term into human medicine. It is this association of effect and cause that makes the term of interest to us. I have tried to manufacture a new word from "claudere," to close, like "claudication," which is derived from "claudere," but have not succeeded, and so I use the simple word "closing."

I ask your attention to a corresponding loss of brain power due to intermittent closing of cerebral arteries, as it is seen in temporary paralysis, and as it takes a part in the production of permanent paralysis.

As illustrating the subject, I give you briefly, in mere outline, the salient points in the case of a patient who was in my male ward in the Royal Infirmary from March 5th to 31st.

Mr. M., aged 50, a farmer, was admitted for observation. The history was that during the months of May, June, and July, 1908, he had three attacks, characterized by tingling in the right arm and hand, a stiff and numb feeling in the right side of the face, and difficulty in articulating. In August he had an attack of complete loss of power on the right side, and stumbling speech. During this attack the symptoms passed off and again returned. Within ten days he had three similar attacks, but not so severe as the first. In November and December he had slight attacks. In the end of December he noticed that when reading he could not see the beginning and the end of a long word. This eye condition led him to consult Dr. Wilson Black of Inverness, who kindly sent me a report on the condition of the eyes, and I show you the perimeter chart of the field of vision at that time. There was a right homonymous hemianopia, but not of the entire right half of the field, only of its outer half.

On admission, the patient weighed 13 st., looked robust and well nourished. The pulse was 84, regular, and of moderate force, brachial pressure was 210, the vessel wall was thick, but not tortuous. The heart dullness extended to the nipple line. The amount of urine passed in twenty-four hours measured from 50 to 60 ounces; the specific gravity was 1020, and it contained a trace of albumen, but no casts were found. The blood was rich in quality. There was no trace of paresis anywhere, no alteration or loss of the ordinary or of any of the special senses. The reflexes were all normal. Speech was clear. The field of vision in both eyes was normal. Dr. Arthur Sinclair, who examined the eyes, found in the left fundus an area of slight local oedema.

In this patient the symptoms may be summarized thus: Extending over a period of three months there were recurring *premonitory* symptoms; in one month there were four attacks of hemiplegia or of hemiparesis, during which time he was medically treated; slight recurrences of premonitory symptoms; and, finally, a temporary and partial right hemianopia. From all these he had recovered by the time he was admitted to hospital, and during his sojourn there there was no return of the symptoms. He had slight digestive disturbance when admitted, which was readily rectified. The arterial pressure varied between 188 and 234. An exclusively milk diet for some

time, the administration of iodide of potassium in 10-grain doses thrice daily, and free action of the bowels made no permanent impression on the haemomanometer readings. On leaving hospital the diet was regulated, and when I last heard of him there had been no return of the special symptoms.

CAUSES OF SUDDEN HEMIPLEGIA.

Before discussing this case, let us look at the causes of sudden hemiplegia or hemiparesis. They are haemorrhage, embolism, thrombosis, and, I add, temporary closing of cerebral arteries. The effects of these are, in the case of haemorrhage, laceration of brain tissue; in the cases of embolism and thrombosis, necrosis and softening of brain tissue. In both laceration and softening there is no restoration of destroyed brain tissue. In temporary closing of arteries there is impairment of function in the part of the brain affected—a halting of brain function from closing of the channels which convey the blood, a condition comparable to intermittent claudication.

DIFFERENTIAL DIAGNOSIS.

Embolism.

Turning to the question of differential diagnosis, a diagnosis of embolism requires at least the coexistence of mitral or aortic disease, or disease of the aorta itself, in addition to sudden paralysis. Putting this aside, there are the other two causes, haemorrhage and softening; the latter as resulting from thrombosis, and much more common than embolism.

Haemorrhage and Thrombosis.

With regard to thrombosis—thrombus, once formed, does not break up and disappear in some mysterious way; it is a difficult thing to get rid of, and we cannot appeal to recurring formation and disappearance of thrombi to explain our clinical phenomena. That is not pathology. With regard to haemorrhage, there seems to me to have been a marked tendency in later years to regard all cases of sudden hemiplegia as caused by cerebral haemorrhage. However this teaching may have arisen, I do not think it is correct. Tooth agrees with Gowers in thinking that thrombosis is the commonest cause of hemiplegia, and that is my experience. The outstanding symptom which differentiates the two conditions is loss of consciousness. Trousseau, following Récamier, pointed this out long ago. He said that "complete loss of motor power, without accompanying coma, belongs more especially to softening"; and he repeated it thus: "Whenever hemiplegia complete and absolute occurs suddenly, without loss of consciousness, softening of the brain may be diagnosed." Tooth accepts Trousseau's position as the "broad rule" to which there are many exceptions. Liddell says that two-thirds or so of all cases of hemiplegia due to haemorrhage show loss of consciousness. If these estimates are even approximately correct, about 1 in 5 cases of hemiplegia without loss of consciousness is due to haemorrhage. This symptom is therefore of predominant importance in the differential diagnosis of these two conditions.

Temporary and Intermittent Closing of Cerebral Arteries.

In this condition the closing of the blood channels is initiated by the channels themselves, and is effected by means of their muscular coat. It is a local closing, and may be partial or total—that is to say, it may only lessen the amount of blood passing, or it may completely arrest the flow of blood to the part. As a result there is either impairment or complete suspension of function, until the vessel opens and permits the blood to resume its course.

That such local closing of arterial channels is possible is seen in the local syncope of "dead fingers," in the "local syncope" of Raynaud's disease, in some cases of migraine, in the vessels of the retina in quinine blindness, and in temporary partial blindness seen to be due to closing of a branch of the retinal artery. A corresponding explanation is required for the phenomena which were exhibited in Mr. M.'s case. His case is only one of many I could give you. They are known to all physicians and practitioners of experience, and, were it necessary, I could gather striking examples of such recurring phenomena from

medical literature. The complete recovery which occurred in Mr. M.'s case could not have occurred had there been either haemorrhage or softening in the motor regions affected, or in the part of the optic radiation which determined the hemianopia.

A WORKING HYPOTHESIS.

The explanation, moreover, provides a working hypothesis of great practical value, and this is the reason of my bringing the subject before you to-day. Take the phenomena in Mr. M.'s case. Extending over three months there were symptoms commonly regarded as premonitory, or as the threatenings of cerebral haemorrhage. But what is meant by *threatening* of cerebral haemorrhage? It is either haemorrhage or not; there is no half-way stage in haemorrhage, although the quantity varies greatly. But these threatenings also precede thrombosis and softening; they also precede the paralysis due merely to temporary arterial closing, as in Mr. M. The "paralytic stroke," due to any of three different pathological processes, has the same premonitory threatenings, and it is of no small practical importance that there should be a clear picture of the clinical pathology of this stage. These threatenings can only, I take it, be due to irregular vessel action, and so to irregular blood flow through the part. There must be vessel constriction, local in site, varying in degree and in extent, coming and going, intermittent.

EFFECTS OF INTERMITTENT CLOSING.

The effects of intermittent closing may be placed in four groups:

1. It leads to impairment or suspension of the function of the part affected, as manifested by sensory disturbances and motor faltering or paralysis.
2. It favours thrombosis in certain states of the vessel wall, or of the blood itself, or of both combined.
3. If the heart be weak the blood flow is more easily arrested, thrombosis is favoured, and the risk of softening is greater.
4. If the heart be strong and the vessels not sound, haemorrhage is liable to occur. The haemorrhage, however, does not take place in the closed portion of vessel; it must take place on the proximal side, where the pressure is raised because of the distal closing.

The significance of the premonitory warnings and threatenings is thus seen, and the aim of practical therapeutics is made clearer.

ETIOLOGY AND TREATMENT.

We have thus a picture of intermittent closing of arteries occurring in different parts of the brain, occurring in persons of very different types, and occurring, as in our case, at the early age of 50 years. If you ask about the etiology of this irregular vessel action we are at once brought face to face with fundamental problems. There are questions of constitution such as are embraced in the words "gouty" and "rheumatic"; there are questions in metabolism, such as the accumulation of the waste products of cell activity, the fatigue toxins, and the inefficient removal of such. There are questions of food and drink; there is the condition of the bowels, and the absorption of toxins from the intestinal tract. There are questions of skin and kidney action. There are questions of mental work and of mental repose; there is emotion and passion. In no department of our work is it so necessary to form a sound all-round estimate of the patient—habits, digestion, bowel action, kidney efficiency, the cardio-vascular system, have all to be reviewed and estimated aright, if we are to be guided aright. What is to be aimed at is the lowering or removal of the abnormal vessel irritability. In the robust plethoric subject the threatenings of which we have spoken have long been treated by active purgation, diminished food supply, physical and perhaps mental rest. All these measures are good, but the plethoric type does not supply the greater part of the cases we are now considering. In many cases active purgation, and all extreme measures, are to be avoided as positively hurtful. Rest, physical and mental, gentle bowel action, spare diet with no red flesh, are the general lines of treatment. The irritable tonicity of the vessel wall is in many cases wonderfully controlled by spiritus aetheris nitrosi. When symptoms appear to be more urgent I have much confidence in erythrol tetranitrate in $\frac{1}{4}$ to $\frac{1}{2}$ grain

doses, repeated every three, four, or six hours, as may be required. The action is temporary, but it allows time for the bowels to act fully and for the elimination of the excess of waste products in the tissues. Thereafter restricted diet and daily bowel action may prevent the recurrence of symptoms. In other cases the vessel irritability is markedly allayed by the administration of iodide of potassium. I would very strongly warn you against violent purgation with calomel and similar drugs in persons with a feeble circulation, for the lowering of such a circulation is the surest way to ensure softening, and even subsequent haemorrhage. The feeble circulation has to be recognized behind thickened and incompressible vessels; see to it that you are not deluded into thinking that there is a strong heart because there is a high haemomanometer reading. A tonic as nux vomica may be a valuable addition to spirit of nitrous ether or to iodide of potassium in cases of this kind.

By a true estimate of the patient as a whole, by the wise use of the simple methods indicated, the irritability of the arterial wall disappears and can be prevented returning to any alarming extent. In this way we can by foresight prevent such truly pathetic cases as must be known to most of you.

The Wightman Lecture

ON

CONGENITAL HEART AFFECTIONS,

ESPECIALLY IN RELATION TO THE DIAGNOSIS
OF THE VARIOUS MALFORMATIONS.

DELIVERED BEFORE THE ROYAL SOCIETY OF MEDICINE,
SECTION OF DISEASES OF CHILDREN.

BY GEORGE CARPENTER, M.D.,

PHYSICIAN TO THE QUEEN'S HOSPITAL FOR CHILDREN, LONDON.

GENTLEMEN,—In acknowledging the honour that you have conferred upon me by inviting me to deliver the Wightman Lecture, I must confess that the acceptance of this distinction proved far easier than the selection of a suitable subject for the occasion. Passing now to the subject of congenital affections of the heart, I will first draw your attention to the known causes which bring about the various departures from the normal, which are classed under the heading "Congenital Heart Affections."

The coincidence of congenital malformations of the heart with malformations in other regions of the body is a matter of everyday experience. Heart malformations are not always accompanied by other congenital bodily defects, but they are sometimes found in association with such diverse structural faults as hare-lip, cleft palate, ill-developed teeth, supernumerary auricles, and supernumerary nipples, polydactylism, syndactylism, webbed fingers, steeple-skull, defects in the abdominal wall, herniae, ill-developed and absent rib cartilages, undescended testis, atresia of the anus and rectum, post-anal dimple, congenital opacity of the cornea, coloboma iris, congenital defects of the orbits, congenital ptosis, not to mention many other congenital irregularities of development which are obvious to the naked eye—cretinism, for instance—which sometimes bear them company.

Since heart malformations have been shown to be a family complaint in some instances, it must be admitted that inherited as well as acquired defects in the sperm and germ cells of the parents are of some importance in the production of malformations of the heart, as in other regions, for erratic tendencies to produce deformities, when once acquired, are handed down from parents to children.

Inquiry has been directed to the state of the mother's health, both bodily and mental, during the pregnancy to serve as an explanation for the production of congenital cardiac defects. When it is realized that the heart is perfectly formed, though in miniature, in seven weeks' time from the date of conception, it is obvious that there must be some error in the various examples that have been recorded of congenital heart disease following upon frights and so on during the last two or three months