When seen at 2.30 p.m. there was less pain and the diarrhoea seemed to have ceased. The tongue had a whitish fur. After this the diarrhoea returned, and the eyes were noticed to become of a darker shade; later he became restless, with slight low delirium.

At 8.30 p.m., when I saw him again, the patient was collapsed, the aspect being that of abdominal disease. The respiration was jerky and somewhat laboured, the pulse small and rapid, and the abdomen was somewhat distended, hard, and tympanitic; the temperature was 100.5° F. Nothing abnormal could be detected in the chest.

The patient was brought to Her Majesty's Hospital (in connexion with Dr. Barnardo's Homes) and warmth, stimulants, and turpentine enemata applied. Later, the patient became somewhat anæmic, the dyspnoea and restlessness becoming more marked. Dr. Richardson, of Commercial Road, saw the patient with me and agreed that he was not fit to stand any operation; ergotin and morphine were administered. Marked tenesmus was present while in the hospital. Gradually the patient sank and died at 11.40 p.m., after one day's illness.

At the necropsy on August 11th, made with the co-operation of Dr. Richardson, nothing abnormal could be seen in heart, lungs (except adhesions of both pleurae), kidneys, spleen, or liver.

On opening the abdomen, the large intestine was found to be much distended (to a diameter of 2½ in. to 3 in.); the small intestine was not so distended. There was a little free sanguineous fluid in the peritoneal cavity, but no sign of peritonitis. The stomach and small intestine contained a yellowish fluid; the mucous membrane of the stomach being somewhat congested. The contents of the large intestine consisted of much gas and a dirty-brown fluid; the mucous membrane was somewhat congested.

In the upper part of the rectum a hard mass was found, which, on removal, was seen to consist of a concentric growth in the rectal wall, extending for 2 in. along the rectum and constricting its cavity. Behind this mass there were several enlarged glands, one of them being as large as a cherry. On opening the gut, the wall was seen to be thickened to ½ in., while the mucous membrane was eroded and ecchymotic; the stricture admitted the tip of the little finger; the growth and enlarged glands were all remarkably hard.

A report from the London Hospital Pathological Institute on the microscopical appearance of the growth states as follows:

"The section at one spot shows mucous membrane with thickened basement membrane and much round-celled infiltration; elsewhere it has been completely destroyed. For the most part the section consists of irregular fibrous septa enclosing spaces; these spaces are subdivided by delicate strands, which stain blue with Gram. Arranged irregularly in the spaces, and more densely along the fibrous septa, are polyhedral cells of an epithelial type whose nuclei stain well with haematoxylin. In one spot there are ducts lined with an epithelium, which is proliferating in irregular fashion, but in places has a columnar arrangement. In a few places colloid material is recognizable, but the method of preparation has destroyed it in most of the spaces. The peritoneal coat is thick and fibrous. The section is that of a colloid carcinoma of the large intestine."

A CASE OF HAEMATOMA OF THE LABIUM MAJUS FORMED DURING LABOUR.

By JOSEPH JONES, M.B., CH.B. VICT.,
Leigh, Lancashire.

In the EPITOME for September 16th, paragraph 168, an account is given of the spontaneous development of a haematoma during the seventh month of pregnancy. A record of the formation of one during labour may be of interest.

The patient was a primipara of 35 years, and rather unintelligent. At the beginning of the second stage the head was still at the brim of the pelvis and not properly engaged. As it remained there an undue length of time it was drawn into the cavity with the forceps which were then at once removed, and the case left to Nature. The child was born forty minutes later in the first cranial position, and weighed about 6 lb.

About two hours before birth a painful swelling the size of a hazel-nut was observed in the mother's right labium majus, at the junction of its lowest and next fourth. There was no change of colour. It continued to increase in size until, when

the head had begun to distend the soft parts, it had attained the dimensions of a hen's egg. As it appeared probable that its bulk, in a part already on the stretch, would cause a bad tear, an incision ½ in. long was made into it, parallel with the vulva; some liquid venous blood was liberated, and more continued to flow in a thin but continuous stream for a few minutes, amounting to about 2 oz. altogether. The tumour, however, did not disappear until the vertex had become plainly visible through the vulva, when the pressure caused the vaginal wall to rupture over the tumour, which was ejected from the vagina as a clot measuring 2½ in. by 2 in. by ¾ in. Delivery soon followed. There was no further bleeding of importance.

The tear thus started continued for an inch on the skin surface of the perineum, and to the extent of 2½ in. up the posterior wall of the vagina. It was duly sutured.

The subsequent history of the patient was satisfactory, except for a rise of temperature beginning about the end of the second week, and lasting four or five days, which was traced to an attempt to douche the vagina with an enema syringe, using only the bone rectum nozzle. The temperature fell from 102° F. to normal within twenty-four hours of this error being corrected.

The only etiological factors noted, apart from such increased blood pressure as always occurs in parturition, were: (1) The advanced age of the primipara, the effect of which in causing degeneration of the blood vessels was probably negligible; (2) the prolonged lodgement of the head at the brim, which apparently obstructed the venous return by pressure on the common iliac, since it was in this stage that the tumour began to form; and (3) the mental condition of the patient, which, perhaps, may have had some analogy with the state of mind of lunatics in whom haematoma of the ear is relatively very common.

The nature of the influence of insanity in producing blood tumours is unknown, and in the absence of more obvious cause it would appear reasonable to attribute to the patient's general stupidity a possible share in the bringing about of the haematoma in this instance.

RATS IN RELATION TO PLAGUE.

By BRUCE SKINNER,
Lieutenant-Colonel, R.A.M.C.

III.

Those who have become sufficiently interested in this question will desire to have available some means of distinguishing the tick which may prove to be the vehicle of plague to men and rats.

Before proceeding, attention must be called to an address¹ on Plague in India, given by Dr. Creighton before the Society of Arts on May 18th last. There are two points in this address which have a bearing on the subject under consideration. Dr. Creighton points out that there are two great seats of plague, "one the plains of the North-West, the other the alluvial valleys of the Deccan and Gujerat." These areas, marked on a map of India on p. 811 of the report of his address, show "fairly enough where the interest really lies." The other point is that Dr. Creighton draws attention to the comparative freedom from plague of stone-built villages, as shown in Ratnagiri; while the mud-built villages of Satara have suffered heavily. It can be readily understood that in the latter there would be every facility for the operations of ticks. The bearing of the point first mentioned above will become obvious on reference to the accompanying chart.

Out of some 250 species of tick noted by Professor Neumann in the *Mémoires de la Société Zoologique de France*, vols. ix, x, xii, the distribution of *Hyalomma aegyptium* forces itself upon the observation as coincident with that of the endemic plague areas, Yunnan excepted. Its occurrence in Bombay points to the West as the source of its arrival in India; from Bombay its spread may be found to coincide with the extension of plague. But there is another focus. Kumáon and Garhwal have long figured in the Indian reports² as plague centres in which an advance of the area of the disease was noted at each fresh outbreak. "This disease, which first commenced in the snowy range, is undoubtedly coming lower and lower every year."² We probably shall find an extension southwards from Lob Nor, across Thibet, of the *Hyalomma aegyptium*. The early history of the latter invasion will never be detected step by step, but it may be safely inferred that it entered Garhwal,