the pelvic colon directly; most valuable information can thus often be obtained. I believe that a careful examination of the pelvic colon with the sigmoidoscope ought always to be made in cases in which symptoms of colitis exist. Serious errors in diagnosis can thus often be avoided and definite information can be obtained on which a rational line of treatment may be based. Those cases in which the symptoms of colitis are due to a tumour (simple or malignant) situated high up in the bowel can by such an examination be separated from those in which the cause of the colitis is different, and a growth perhaps detected at a date when its removal is still possible. The two following cases are good examples of this:

CASE II.

A woman, age 42, under my care at St. Mark's Hospital, had suffered from diarrhoea for about three months before I saw her. The bowels acted several times a day and the stools consisted largely of slimy mucus. She passed considerable quantities of mucus at times. During the last month the bowels acted six or seven times in the twenty-four hours. At no time had she ever seen any blood in the stools. When I first saw her she had been under treatment at another hospital for colitis for six weeks. A careful digital examination of the rectum revealed nothing abnormal. I was inclined to believe that the case was one of true mucous colitis, especially as she had never passed any blood, and had a history of attacks of pain in the abdomen, followed by diarrhoea and the passage of much slimy mucus. I have recently made it a rule, however, always to examine such cases with the sigmoidoscope as a routine. And in this case the examination revealed the presence of a malignant stricture situated about 7 in. or 8 in. from the anus.

A careful examination seemed to indicate that the growth could be successfully excised from the abdomen. At the operation, however, I found on opening the abdomen that there were several secondary malignant glands up the side of the aorta, and I was therefore obliged to abandon the excision and perform colotomy.

CASE III.

The patient was a man, aged 63, who was under my care at St. Mark's Hospital in November, 1904. Eighteen months before I saw him he had an attack of diarrhoea, accompanied with considerable pain and tenesmus. This lasted for about six weeks, and then stopped. There was no blood in the stools. Two other similar attacks of pain and diarrhoea had occurred between this and the time when I first saw him. During the last attack he noticed a little blood with the stools on one or two occasions. On examining him with the sigmoidoscope, I found a papilloma about the size of a five-shilling piece situated on the posterior wall of the rectum between the middle and upper Houston's valves. I subsequently removed the growth, after excising the coccyx, and restored the rectal wall. The patient made a good recovery, and has remained well since.

A case is recorded by Dr. Pye-Smith in the *Transactions of the Pathological Society* (vol. xlvii) which is very similar to my first case:

The patient was a woman, aged 50, in Guy's Hospital. She had been suffering for over a year from constipation with distension of the abdomen and occasional attacks of diarrhoea. During the whole time she had been passing, at frequent intervals, large soft gelatinous shreds, and more or less complete mucous casts, per anum. She had frequent gripping pains, and there was visible peristalsis of the intestines. A microscopical examination of the casts showed that they consisted of mucus. The patient died, and at the *post-mortem* examination it was discovered that she had a carcinomatous stricture at the commencement of the sigmoid flexure. The colon both above and below the stricture was greatly thickened and distended. After removing the adherent mucus, the mucous membrane of the colon was found to be normal except at the site of stricture.

Dr. Pye-Smith mentioned two other similar cases, one of Dr. Goodhart's and one of Mr. Hutchinson's.

These cases show, I think, that typical symptoms of mucous colitis may exist with a malignant growth of the bowel and that without a careful local examination it is often impossible to distinguish between a case of simple mucous colitis and one of malignant disease of the bowel.

In other cases, again, an examination with the sigmoidoscope reveals the presence of ulceration and further often enables us to ascertain the position and limitations of the ulcerated area and thus to form a definite opinion with regard to the best method of treating it. In fact the examination with the sigmoidoscope replaces a more or less guess-work method of making a diagnosis in such cases by an exact observation.

CASE IV.

The patient was a young man, a patient of Dr. Bruce Porter, aged 27. Three years ago, while in South Africa, he had a bad attack of diarrhoea (said not to be dysentery), for which he was treated in hospital. His stools at that time consisted of blood and slime. He got slightly better, but more or less ever since he has suffered from diarrhoea and has passed blood and slime. On returning to England he got much worse, and when I saw him with Dr. Bruce Porter he was much emaciated, was suffering from more or less continuous diarrhoea, and was

passing much blood and mucus. An examination with the sigmoidoscope revealed the presence of numerous shallow ulcers with a granular base extending throughout the whole of the sigmoid flexure. The appearance of the ulcers was not tuberculous. A careful examination of the discharges showed the presence of pus, but there were no tubercle bacilli or bilharzia parasites. Considerable improvement followed a carefully-regulated diet and irrigation of the colon.

In cases of real mucous colitis an examination with the sigmoidoscope is sometimes the only method by which we can be certain that we are dealing with that condition and not with a new growth in the pelvic colon. This is well shown in Case v. The similarity between the symptoms in this case and in Case i, where the symptoms were due to cancer, show how impossible it is to make a correct diagnosis without a direct examination of the parts such as can be obtained with the sigmoidoscope.

CASE V.

The patient, a man aged 35, was employed as a chef. For about three months he had suffered from diarrhoea and constipation. The bowels often acted six or more times in the day, and the stools consisted largely of slimy mucus. He had never passed any blood. The attacks of diarrhoea were generally initiated by colicky pains in the abdomen. There were often periods of constipation between the attacks. A careful examination was made of the abdomen, and the pelvic colon and rectum were also examined with the sigmoidoscope. Beyond some slight congestion of the mucous membrane, however, nothing abnormal was found. I came to the conclusion that the case was one of mucous colitis, and he was placed upon von Noorden's diet. The symptoms disappeared almost at once, and he remained free from his trouble until some months later, when he got a recurrence, as the result of dietary indiscretions. A return to the diet, however, was again effectual in getting rid of his symptoms.

In the present position of our knowledge of the numerous conditions of the large bowel which are often grouped under the name of colitis, it does not seem possible, or even desirable, to attempt to classify these various conditions too exactly. But I think it is most important that the greatest care should be taken to exclude carefully the presence of any definite lesion of the bowel before attempting to treat a patient with symptoms of colitis. The cases I have quoted seem to me to prove how utterly fallacious a diagnosis of colitis based entirely upon symptoms may be. And I believe that a careful examination of the pelvic colon with the sigmoidoscope should be made a routine practice as a preliminary to treatment in all cases of colitis.

WANDERING SPLEEN: HAEMORRHAGE WITHIN THE CAPSULE: SPLENECTOMY: RECOVERY.

By CHARLES P. CHILDE, B.A., F.R.C.S.,

Assistant Surgeon, Royal Portsmouth Hospital; Senior Surgeon, Medical and Surgical Home for Women, Southsea.

Cases of wandering spleen are, generally speaking, of sufficient rarity and sufficient interest to place on record. The following instance presents some unusual features even in this uncommon condition, which invest it with additional clinical and surgical value.

REPORT OF CASE.

History.

P. L., aged 54, sent to me by Dr. Knott, of Southsea, was a spare woman of healthy rosy complexion; she was married, and had had nine children, five of whom were living; four had died of children's complaints. She had always had good health and normal confinements; had never suffered from any abdominal or other injury, and till the present illness had never felt any pain or inconvenience in the abdomen or pelvis.

She stated that about two months previous to her admission to the Royal Portsmouth Hospital on November 5th, 1904, she had influenza, though in the light of future developments it is more probable that her illness was associated with some pathological changes in the tumour. While recovering from this attack, which, whatever its nature, was accompanied by the usual symptoms of feverishness, she was seized with sudden pain in the lower abdomen, retching, and faintness. She stayed in bed for a week, during which time she had fever and profuse perspirations, but no return of the pain after the first day. Dr. Knott attended her during this illness, and discovered a tumour in the lower abdomen. The acute symptoms subsided quickly, and did not return up to the time of my seeing her two months afterwards.

When I first examined her, at the end of October, a largish tumour occupied the false pelvis and lower abdomen, extending

up higher and involving the left side more than the right, as shown in diagram (Fig. 1). The left part (a) was completely

Fig. 1.—Diagram of condition before abdomen was opened: (a) entirely dull and tender; (b) partially dull.

dull, the portion (b) which extended into the right iliac fossa being only partially so, owing to overlapping intestine. It could be moved, to a certain extent, from side to side, but could not be pushed up at all. The loins and upper abdomen were resonant, and the tumour did not extend into the kidney region on the left side. It was smooth, rounded, and felt of fluid consistence, but there was no distinct thrill. No tumour could be felt in the pelvis on vaginal examination, but there was a sense of resistance in front and to the left of the uterus, and bimanually an impulse could be communicated to the finger on this side, which could not be felt on the other. The uterus was retroverted and normal in length, and the tumour did not appear to be attached to it. The left part (a) was distinctly tender on pressure. There were no bladder symptoms and the urine was normal. Some constipation was present. The temperature was not elevated.

The diagnosis was made of ovarian cyst, the acute attack being attributed to torsion of the pedicle, with haemorrhage into its cavity, followed by peritonitis which had caused some adhesion and fixation of the tumour. It will be seen that from the history and physical signs this diagnosis was not unwarranted; indeed, the case seemed a fairly clear one.

Operation, November 10th, 1904.

After opening the peritoneal cavity in the middle line by the subumbilical incision, a rounded cystic tumour was exposed with omentum adherent everywhere over its front and sides. On partially separating the adhesions and plunging a trocar into the cavity dark blood escaped to the amount of about a pint, and the diagnosis was thought to be verified. The tumour could not be brought forward into the wound, being firmly fixed in the pelvis. It was also remarked that the cyst wall tore very easily. After extending the abdominal incision above the umbilicus, enlarging the opening in the cyst, and sponging out its contents, and further separating omentum adherent to its sides, it was still found that the tumour could not be delivered into the wound, being bound down especially in the left iliac fossa. On separating further adhesions a dark mass that looked at first, as it appeared from among them, like a coil of deeply congested and distended intestine, was brought into view from the right side of the pelvis. This was soon seen to be the upper pole of the spleen, and the nature of the case was now for the first time evident—a wandering spleen, with a large blood cyst attached to it. The lower pole was found to be firmly fixed to the left wall of the iliac fossa, the spleen lying obliquely across the pelvis (Fig. 2) with the pedicle twisted on itself, and the lower part of it very tense. The upper pole, which was lowermost, and turned towards the right, lay in front of the fundus of the uterus, contributing no doubt to the retroversion of that organ previously noted. The external and posterior surface of the spleen facing upwards formed the lower wall of the blood cyst. Its contents were included between this surface and the capsule, which had been torn off it by the haemorrhage. All adhesions having been freed, an attempt was made to peel off the

broad attachment to the left iliac wall, so as to get the tumour out of the wound, and deal with the pedicle last. In doing this the splenic tissue tore very readily, and the free haemorrhage that ensued enforced the abandonment of this procedure. A sponge was therefore plugged into the tear, and pressure kept up by an assistant, while the pedicle was dealt with *in situ*. It was attacked from above and tied in sections. It was very wide, and the lower part of it very tense, so that I was in considerable fear of the ligature slipping here. Each section, as it was tied, was divided between the ligature and a large Spencer Wells forceps placed on the distal side. Finally the lower pole was peeled off the left iliac wall, and spleen and blood cyst removed. The operation lasted 1 hour 10 minutes. A gauze drain was inserted in Douglas's pouch, which was removed in 24 hours.

Fig. 2.—Diagram of condition found at operation: (a) blood cyst; (b) spleen. The spleen was below and behind the cyst, and not visible till the latter was evacuated.

No shock followed the operation, and a normal recovery ensued, the temperature not rising above 99° F.

The length of the spleen after removal was 8 in., breadth 4 in., and weight (after emptying the blood cyst) 8 oz. A section of it was reported on by the Clinical Research Association as follows:

The tumour has the characteristics of a spleen. The capsule is somewhat thickened, and the vessels traversing its walls have hyaline thickened walls and sheaths. There is also some increase of the fibrous stroma of the splenic pulp.

REMARKS.

As regards the diagnosis, the likeness of this case in its history and physical signs to an ovarian cyst, with twisted pedicle and haemorrhage into its cavity, is manifest. The abdomen, however, has a nasty habit of presenting a "surprise packet" to the operating surgeon, and the one under notice serves to illustrate the fact that not every unilateral cystic tumour rising out of the pelvis of a woman is of ovarian or tubal origin. The more or less transverse position occupied by the spleen in this instance is the one which has been usually noted in cases of displacement. How this comes about is readily explained if we consider the normal position of the spleen and its attachments prior to descent. It lies very obliquely in the left hypochondrium, with its upper pole much nearer the middle line than the lower. It rests in this position on the pleuro-colic ligament, stretching from the diaphragm to the splenic flexure of the colon, and is attached to the stomach by the gastro-splenic fold of peritoneum, which reaches from the hilus to the left border of the great omentum. Its normal position is, therefore, oblique, and its attachment toward the middle line. As the spleen descends in cases of displacement, this oblique position and attachment to the stomach cause the upper pole to tip forward towards the middle line, and the organ to assume a more and more horizontal position. The upper pole turned toward the right therefore comes to lie eventually on a level with the lower pole, turned towards the left, and a still further descent will result in the spleen assuming an oblique position with the upper pole lowermost. Moreover, this change of position must cause a rotation of the pedicle from left to right, and from above downwards, so that its upper border will eventually lie lowermost. Where the

descent and rotation are gradual and accompanied by stretching no serious symptoms may ensue; but it is easy to imagine that the rotation may be sudden or suddenly aggravated, in which case, as in other abdominal tumours with pedicles, peritonitis or haemorrhage, or both, may occur. It is not remarkable, considering the close attachment of the spleen to the left end of the stomach, that these patients should commonly suffer from abdominal pains and indigestion; and one would expect this to be so especially in traumatic cases, where the displacement, or the initial stages of it, at all events, occur suddenly. My patient suffered from no abdominal symptoms whatever—no dragging pain, no feeling of sickness, or indigestion—and but for the accident of haemorrhage and peritonitis would have probably been going about now, quite unconscious of any internal abnormality. The most careful inquiry failed to elicit any history of injury, and her absence of previous symptoms is explicable on the assumption that the displacement was a gradual one, so that the gastro-splenic omentum had time to stretch and lengthen out, and the stomach was thereby

Fig. 3.—From a photograph of the tumour on removal (two-thirds natural size). The blood cyst when distended was the size of a large cocoa-nut.

enabled to accommodate itself to the abnormal drag upon it. Finally, some sudden increase in the rotation of the pedicle took place, resulting in the haemorrhage and peritonitis which first directed attention to her condition. It may be mentioned that the patient was rather a spare woman, and that she had had nine children, not by any means an unusual number among the class that can least afford to support them.

It must be admitted that splenectomy is not the ideal treatment for wandering spleen. At the same time, owing to practical difficulties in adopting the alternative and more rational measure of splenopexy—these difficulties being, first, that of replacing the organ, and, secondly, that of keeping it there when replaced—the former is frequently the only feasible plan of dealing with the condition. In the case here quoted

it was obviously the only method open. The position the spleen had come to occupy low down in the pelvis, its broad adhesion to the iliac wall making separation before the pedicle was tied impossible, the attachment to it of a large blood cyst, and the urgent and dangerous symptoms the patient had suffered from in connexion therewith, made splenectomy imperative.

Several methods of splenopexy have been adopted. Kenwar in 1895 was the first surgeon to perform the operation, making use of tamponnade for the purpose of keeping the organ in position. In one case he was obliged to remove the tampon for ileus. Rydygier in the same year made a pocket in the abdominal parietes in which he fixed the lower end of the spleen, suturing it also to the tenth rib and parietal peritoneum. Basil Hall, acting on similar lines, ingeniously made use of the notch (situated low down in his case) by bringing the parietal peritoneum and the aponeurosis of the abdominal muscles round it, so as to grasp and fix the lower end. Such an unusual conjunction of circumstances as a notch low down in a wandering spleen will be as rare as honesty in politics.

Balch's case, published by J. Collins Warren, resembled my own in some particulars. The spleen was adherent by one pole to the pelvic brim, and the uterus was tipped backwards by the other, its position being very similar to that found by myself. There was, however, no blood cyst in connexion with it, and Balch had to deal with the displacement only. He succeeded in freeing the adhesions to the pelvic brim, using ligatures and the cautery to arrest haemorrhage. The separation of these left plenty of raw surface, and the organ was then pushed up into position, gauze packing being used to keep it there and promote adhesions. I attempted with great gentleness to separate the broad attachment of the spleen to the iliac wall, not, however, with the idea of performing splenopexy, for I considered that from the first not feasible, but of dealing with the pedicle more easily. The bleeding was, however, so fierce from the torn splenic tissue, welling from it as from a squeezed sponge, that this had to be abandoned. No cautery or ligatures would have been of any use here. I was obliged to keep up sponge pressure in the tear, and to deal with the pedicle as expeditiously as possible, whereupon, of course, the bleeding ceased spontaneously. Thus it will be seen that splenopexy is the ideal treatment for this condition, to be adopted where the special circumstances of the case admit of it, but that splenectomy is frequently the only feasible course on account of the practical difficulties of placing the spleen in its natural position and of keeping it there, or on account of the trouble from haemorrhage, or, lastly, from some such complication in connexion with the displacement as occurred in the case under notice.

With regard to blood cysts in connexion with the spleen, Giulano gives a record of 15 cases of haemorrhagic cysts, including his own. They were apparently of a different nature from the one recorded here, occurring in the enlarged spleens of malaria, a portion of which softened and became diffuent, so forming a blood cyst. The origin of the cyst in my case was, as the specimen shows, entirely different. The spleen itself is seen to be entire, and the cyst is not within its substance, but between the capsule and the pulp. It was formed apparently by a sudden haemorrhage occurring from the rupture of a vessel at the time of the acute attack above described, the symptoms of which were those of an internal haemorrhage. This tore up the capsule from the splenic pulp, a blood cyst being thus formed between the two. It was probably caused or favoured by the torsion of the pedicle, and was followed by a localized peritonitis, and is comparable to haemorrhage within an ovarian cyst from the same cause, a condition from which in its symptoms and physical signs it was undistinguishable. Giulano recommends, in the cases described by himself, opening the cyst, stitching the wall to the parietes, and drainage. This treatment was not applicable here. As far as I have been able to ascertain by a reference to the literature on the subject, there is no other recorded instance of the condition found in this case, namely, a spleen fixed in the pelvis, and associated with it a large blood cyst of the nature described above. Referring to the immediate risks of the operation, during the last decade there have been 43 recorded cases of splenectomy for wandering spleen, with 3 deaths, a mortality of 7 per cent. Inasmuch as in a novel and rare operation there is a not unnatural disposition to record every successful case, while the contrary are prone to shrink publicity, we may assume that the mortality is somewhat higher than this. So that on the score of immediate risk

alone this operation is not to be recommended, unless it is clearly causing the patient great inconvenience and suffering, or possibly, as in my case, danger to life. A suitable truss would, of course, too, in any ordinary case be tried and found wanting prior to the consideration of any operative procedure.

As might have been expected, after the removal of an organ of the apparent importance of the spleen, though very little is certainly known of its functions, ultimate ill-effects have been recorded in some cases. After the patients have made a good recovery from the operation, they have begun to lose ground and to manifest symptoms of weakness, emaciation, thirst, and drowsiness. Thus, of the three cases brought before the Clinical Society in 1896 by Messrs. Ballance and Pitts, two suffered in this way. The unfavourable symptoms were successfully combated by the administration of sheep's spleen, raw bone marrow, cod-liver oil, and arsenic, and the patients were eventually restored to good health. All three of them, however, exhibited enlargement of the superficial lymphatic glands. In the case recorded here nothing except ordinary food, and no medicine whatever, has been administered since the operation. One year has now elapsed, and the patient has suffered no inconvenience at all from the removal of her spleen, nor does she appear to miss it in the least. There is also no enlargement of any lymphatic glands that can be palpated. A report on the blood kindly made for me by Dr. Leon on February 8th, 1905—that is, three months after operation—showed a decrease of white corpuscles (7,000 per cm.), and the lymphocytes relatively decreased. Possibly in cases where no after-effects have followed splenectomy, spleniculi may have existed which were not observed. None were observed in this case, and they were certainly not present in the situations where they have been usually noted—namely, the gastro-splenic and great omentum. At the same time, as no definite search was made for them, they may have been present and have escaped notice. The Malpighian bodies, situated on the minute branches of the splenic artery, are closely packed with lymphocytes, and their removal, along with the spleen, seems to throw increased work on the other generators of lymph corpuscles—the lymphatic glands, which consequently become hypertrophied. The other chief supposed rôle of the spleen is the purification of the blood by the destruction of effete red corpuscles, and the numerous extractions such as formic, butyric, lactic acids, etc., which are obtained from a chemical analysis of the spleen, are probably derived from the destruction of these worn-out constituents. It might be anticipated, therefore, that the removal of this great purifier, so to speak, of the blood would be followed by some such toxic symptoms as have been enumerated above, owing to the introduction into the system of poisonous matter, of which the spleen normally cleanses it. The ultimate disappearance of these symptoms may have been due to the remedies given. More probably they were owing to compensatory action becoming established in some other gland, or to the presence of spleniculi, which have stepped into the breach and gradually restored the normal balance. In the case recorded here no remedies have been given, which might have anticipated, so to speak, the want in the economy produced by the loss of the spleen. Yet its removal has been followed by no apparent deviation from normal health, nor is there evidence of any other organs of the body—for example, the lymphatic glands—being burdened with increased work. Whatever be the explanation, it is remarkable that the spleen, which, *a priori* from its size, its large blood supply, its structure—which alone of all organs in the body allows the blood to come into actual contact with the living cells of the tissues (Starling)—can be removed without the patient apparently suffering any inconvenience or deterioration of health. The probability is that though perfection of health requires the presence and normal working of every organ of the body, a degree of it which is by our imperfect methods of observation satisfactory, may be maintained on a much shorter allowance; and that by some compensatory action which may serve the purpose, but be less than perfect, the human organism may be able to do without some of its component parts apparently without serious inconvenience or disability. I say apparently, because the question as to whether these mutilations eventually shorten life cannot be shelved, and the after-histories of such operations as gastro-enterostomy, removal, or, what amounts to the same thing, exclusion of the large intestine, etc., has yet to be written. It is well to remember, however, that there is an after-history, of which complete thyroidectomy and cophorectomy have afforded

instructive and somewhat disconcerting object-lessons in the past. The mere ability to show a patient apparently in perfect health after the entire removal of some organ which *a priori* might be assumed to be essential to its well-being, though it may flatter our pride, should not in our surgical efforts lessen our respect for every part of the body, or diminish our consideration for any part, because we happen to be ignorant of its functions, or because our patient seems able to do without it. The assumption, for instance, that the large intestine is a cesspool is, to say the least of it, not a bashful one, and is, moreover, contrary to our experience of its uses. On theoretical grounds, I think, every surgeon would treat such an organ as the spleen at all events with respect, and would not remove it unless absolutely obliged to do so.

BIBLIOGRAPHY.

Annals of Surgery, 1903 (J. Collins Warren). *Arch. Prov. de Chir.* Year Book of Medicine and Surgery (Gould). *Elements of Human Physiology* (Starling).

SOME OBSERVATIONS ON PUERPERAL ECLAMPSIA.

By REGINALD DUNLOP, M.D. GLASG.,
Late Senior House-Surgeon, Glasgow Maternity Hospital.

WHILE house-surgeon in Glasgow Maternity Hospital, I had an opportunity of studying 8 cases of eclampsia, and of analysing 42 others which had been treated during the three preceding years. As much interest has been manifested of late in this condition, I should like to draw attention to a few points in its course and treatment which particularly impressed me, and especially to the use of saline infusion and *accouchement forcé*.

Premonitory Symptoms.—Oedema was present in 30 of the 50 cases, but it is no criterion of the imminence of convulsions. It may be present for weeks or months—a symptom of chronic renal disease—but an intense toxæmia may occur so rapidly that convulsions forestall oedema. The swelling is more significant when affecting the cellular tissues of the face. It is a fairly frequent history that oedema of the legs and feet has been present for weeks; then, without increase of this swelling, the face becomes oedematous, and eclampsia supervenes within forty-eight hours. This sequence is well illustrated in 2 of the 8 cases. In the first, oedema of the lower extremities was present for eight days; the eyelids then became swollen, and convulsions ensued in twenty-four hours. In the second, oedema of the feet commenced three weeks prior to the attack, became gradually worse, and was succeeded by swelling of the face. Only eight hours later eclamptic seizures followed.

One of the most important, because one of the most suggestive and unequivocal, of the prodromals is pain in the epigastrium. The pain is at first slight, or there may be only a sense of discomfort; but when this symptom is well marked, it becomes later so intense that the woman is rendered prostrate and is forced to cry out. It may be associated with pain in other regions of the abdomen or in the loins, and may be accompanied by vomiting or difficulty and oppression in breathing. These concomitants are most frequent when the pain has become intense and continuous, and betoken the imminence of convulsions.

Epigastric pain was present in 6 of the 8 cases, but either this is a larger proportion than usual, or less attention has been paid to this symptom than its importance deserves. Munro Kerr¹ in an analysis of 80 cases found it mentioned thrice. Dührssen² in 200 cases found it mentioned once. Parvin³ called it the "least frequent" of the prodromata. It is similarly mentioned in other obstetric works as one of the least important and most inconstant of premonitory symptoms, or is not mentioned at all. Of these 50 cases, 33 are carefully reported as to the presence or absence of prodromals. In 17 of these 33 epigastric pain is mentioned as present, and in 16 is either not mentioned or said to have been absent. In my own 8 cases, when present, it became continuous and very severe shortly before the onset of convulsions. To my mind its significance is very similar to that of headache: both may be present for some time intermittently and not very acutely, but when either becomes intense and constant it is unmistakable evidence that the patient is far advanced in toxæmia.

Duration of Paroxysm.—Herman⁴ says that the duration of a fit is seldom longer than a minute, and that the interference with respiration is such that a fit which lasts longer than two or three minutes must be fatal from asphyxia. He explains