CASE I.

The patient came under my care during the absence from home of her own medical adviser. There was considerable oedema and albuminuria, and also valvular disease of the heart. Her doctor had given a very poor prognosis of the result of her confinement. She sent for me one evening suffering from severe headache; the next morning convulsions set in. I kept her well under morphine hypodermically administered. The convulsions were controlled, but she remained in a comatose condition all day. Next morning spontaneous labour came on, with complete relief of the symptoms, followed by an uninterrupted recovery.

CASE II.

On my first seeing this patient there was albuminuria and convulsions. Spontaneous delivery was followed by cessation of the fits; the patient, however, remained comatose, and died in a few days.

CASE III.

A primipara with severe convulsions. I kept her well under chloroform, and when I found a slight dilatation of the os—no help being at hand—I ruptured the membranes, turned, and delivered. The fits then practically ceased, and the patient made an uninterrupted recovery.

CASE IV.

This case I saw with a friend. The patient had severe convulsions. I advised inducing labour, and while my friend administered chloroform I commenced manual dilatation, and after some perseverance managed to deliver. The convulsions then ceased, and I was told that the child lived and the mother made an uninterrupted recovery.

CASE V.

I was asked by a friend to see a case of puerperal convulsions with him. The attacks were frequent, and I advised inducing labour. My friend, however, seemed to prefer temporizing. We therefore decided to try the effect of the hypodermic injection of the hydrobromide of hyoscyne. The patient went from bad to worse, and later in the day my friend agreed to the advisability of inducing labour. The result was everything we could wish as regards the convulsions, but I understand that the patient afterwards died from other causes, remaining conscious to the end.

CASE VI.

A primipara. When first seen the patient was in a comatose condition, with occasional convulsions. She also had albuminuric retinitis. She remained a day or so in this condition, when spontaneous labour set in. Her condition immediately improved, and she made a good recovery, as far as the eclampsia was concerned, although the eye trouble persisted for some time.

It will be noticed that in each of these cases evacuation of the uterus was followed by relief. To my mind, the course of treatment in every case of puerperal eclampsia is clear and definite. First, try to control the convulsions by the administration of morphine or chloroform (I have never found difficulty in getting the patient under the influence of chloroform; in the various convulsive movements of the mouth and chest enough is drawn into the lungs to produce the desired effect). If the convulsions pass off, well and good; if not—you cannot keep the patient indefinitely under the influence of these drugs—and labour does not soon ensue, then induce delivery. For the latter purpose I prefer Nature's instrument—the human finger—as much the safest, for it is endowed with sensation, and knows what it is doing, and is, moreover, effectual earlier than any other form of dilator. If it is impossible to commence the dilatation with the finger, the sound may be used to start the process, there being no time, of course, to wait for the action of tents.

I offer these suggestions and experiences for what they are worth. I am convinced that if other general practitioners would give us their experiences, it would be useful to all who have these appalling cases to deal with, and would be immensely to the advantage of those most concerned—namely, the poor mothers who are thus so grievously afflicted.

REMOVAL OF IRON FROM INTERIOR OF THE EYE BY ELECTRO-MAGNET.*

By A. S. PERCIVAL, M.B., B.C. CAMB.,
Surgeon, Northumberland and Durham Eye Infirmary.

It was in 1881 that Mr. Snell of Sheffield devised the small electro-magnet that I have here. If the foreign body lie in front of the lens—that is, in the anterior chamber—it is an efficient little instrument. If it be behind the lens, in the vitreous, or embedded in the retina, I have found Snell's instrument to be very unsatisfactory. Suppose that there is a perforating wound in the sclera, Mr. Snell tells one to boil the probe-pointed terminal to sterilize it, then to attach it to

the magnet and insert it into the wound carefully, and to listen for a "click"; this will indicate that the iron fragment has touched the probe end of the magnet. If no sound be noticed, the probe is to be gently moved about until the "click" is heard. As soon as this is heard, the magnet is to be withdrawn with the steel chip attached to it.

In practice I have found that these directions are usually impossible to carry out. If one does succeed in hearing the "click," on withdrawing the instrument the foreign body is wiped off by the edges of the wound; if one enlarges the wound sufficiently to avoid this, there is such a prolapse of vitreous that the eye is subsequently lost; and even the mere act of moving the probe about in the vitreous leaves little hope of preserving a useful eye. The vitreous is a tissue consisting of cells and a fluid matrix, and bacteriologists tell us it is one of the best nutrient media for germs, so that it requires to be treated with even more reverence than most of the other parts of the body. Finally, there is a great physical objection to this little magnet: (1) A screw junction makes a most inefficient magnetic connexion; (2) the pull of an electro-magnet decreases very rapidly as the distance from the coil is increased.

The dimensions of the bobbin are $\frac{3}{4}$ in. by $\frac{1}{2}$ in. The diameter of the core is $\frac{1}{8}$ in.

Haab and Hirschberg have used giant magnets weighing 250 lb. to 600 lb.; these are certainly powerful enough, but they are unwieldy, and require the greatest care in their application, or irreparable damage may be done. Some six years ago I devised this magnet for eye work. It can be held in the hand, and when required the plunger can be removed and a pair of forceps inserted. I will endeavour to show, on this "phantom eye" how the magnet is to be worked.

[The "phantom eye" was represented by a wineglass filled with water, its mouth being covered with a piece of india-rubber sheeting. Part of a needle was passed through the rubber into the glass, from the bottom of which it was then removed through the minute hole of entrance by the electro-magnet.]

The greater number of iron particles enter the eye either through the cornea or the scleral border, and as we must at all costs prevent the foreign body from getting entangled in the ciliary body, in the majority of cases the best way out is through the pupil into the anterior chamber. The eye must be carefully watched the whole time, and if the foreign body lie behind the iris, it must be coaxed round the anterior surface of the lens, until it can enter the anterior chamber through the pupil. In some cases, where the wound is in the sclera, and is quite recent, one is tempted to extract the foreign body by its wound of entrance. There is then, however, the great risk of injuring the ciliary body. The specimen I send round was extracted in this way. I saw the case three

* A paper read before the all-day scientific meeting of the Newcastle-upon-Tyne Division of the British Medical Association, July 16th, 1905.

hours after the injury. The wound was 1 mm. beyond the scleral border, the vitreous was full of blood, the lens was apparently not injured, but no indication of a foreign body was observed. The patient was pretty certain that there was nothing in the eye; he thought he had been struck by a large piece of metal. I may mention that a patient's account of the injury should always be accepted with the greatest caution. I have frequently found foreign bodies present when the patient has been quite certain there were none. With the electromagnet I drew out this fragment, which I have sent round for inspection.* In a few days the vitreous haemorrhage cleared up almost entirely, leaving him with vision about $\frac{1}{4}$. Subsequently, however, a detachment of the retina occurred, so that I am afraid his sight will eventually be but slight.

I have seen it stated that if no pain is felt on the approach of the magnet, there is certainly no particle of iron in the eye. This is a "study table conclusion"; it sounds reasonable, but it is absolutely untrue in practice. I have been so often deceived by it that I never now pay the least regard to it.

It is of very great advantage to be able to cut off the current with ease, as a foreign body imbedded in the retina is more easily loosened by attracting it in jerks.

In conclusion, I should like to say a word about the indications for removal of an injured eye. Many books may be read on the subject, and leave one undecided what one should do in the case that presents itself for immediate decision. I think the following considerations will aid one in forming a correct conclusion, provided that the other eye is sound. It may seem unnecessary to mention this proviso, but I well remember once being consulted by a patient about an injured eye, which his doctor had made all preparations to remove the next day for fear of sympathetic ophthalmia; I found the other eye to be absolutely blind.

In almost every case of sympathetic ophthalmia the exciting eye has been previously perforated either by injury or by ulceration, hence every perforating wound which is not known to be aseptic is always a source of great anxiety. The risk is extreme:

1. If the wound be in the ciliary region—this dangerous area may be defined as the zone $\frac{1}{4}$ in. in width that surrounds the margin of the cornea.

2. If the wound be septic—this can often only be recognized by its consequences. It should, however, be noted that wounds giving rise to suppurative panophthalmitis are not liable to induce sympathetic ophthalmia.

3. If a foreign body be retained in the eye.

Immediate excision of the eye is indicated by the concurrence of two of these conditions, even though some sight be retained; by the occurrence of one of them, if sight be irreparably lost, provided that the vision of the other eye is good.

Under other circumstances excision of the eye may be delayed until symptoms arise without much risk, but operative interference is necessary when there is persistent, severe, or recurrent iritis, when the eye is tender or becomes soft, or when there is sympathetic irritation in the other eye. If irido-cyclitis have already occurred in the sympathizing eye when first seen, excision of the exciter should only be performed if vision is lost beyond hope of recovery, as not infrequently the injured eye will retain the better sight of the two.

If the other eye be blind or absent, intractable pain is practically the only condition of the injured eye that demands its excision, and neurectomy may sometimes be advantageously substituted for this.

* The fragment was $\frac{3}{8}$ in. long, by $\frac{1}{16}$ in.

REQUESTS TO MEDICAL CHARITIES.—The late Miss Mary Elizabeth Gorst, of Chester, whose will has now been proved, left £500 to Chester Infirmary.

INSANITY AMONG NEGROES.—Dr. W. F. Drewery, Superintendent of the Central State Hospital, at Petersburg, Virginia, gives some figures showing the increase of insanity among the negro population. The number of patients sent to the hospital during the year was 170 more than the previous year, showing an increase of about 50 per cent. The number in the hospital was 1,449, or 115 greater than the year before. The smallest number under care at any one time was 1,101 and the largest 1,305. The number present at the end of the year was 1,181, or 80 over the previous year. The growing prevalence of mental disease among negroes is ascribed by Dr. Drewery to physical conditions and degeneracy induced by indolent, unhygienic, and immoral modes of life, as well as to the influence of heredity, rather than to mental stress or overwork.

SEVENTY-THIRD ANNUAL MEETING

OF THE

British Medical Association.

Held at Leicester, July 24th, 25th, 26th, 27th, and 28th, 1905.

PROCEEDINGS OF SECTIONS.

SECTION OF LARYNGOLOGY, OTOTOLOGY, AND RHINOLOGY.

FREDERICK W. BENNETT, M.D., President.

PRESIDENT'S INTRODUCTORY REMARKS.

[AFTER some introductory remarks the PRESIDENT proceeded:]

It is interesting to look back upon the great change which has taken place during the last twenty years in the importance attached to the treatment of diseases of the throat, nose, and ear. Then only the larger centres of population were even moderately provided for. At the present time there are few important hospitals where there are not special posts, and, what is still more important, there are few practitioners who do not take some degree of interest in these studies. But the general public still remains far too indifferent to the importance of the early treatment of these diseases. What we shall look back upon with the most profound pleasure when our medical life is finished will be probably the effect our work has had in the prevention of disease and in advancing knowledge with a view to averting from the community dangers to which its members would otherwise have been exposed. To me at least this aspect of our work has always appealed.

For seventeen years I have taken an active interest in the educational work of this borough, and I have always been impressed by the great wastage due to the unrecognized beginnings of diseased processes. The results of Dr. Cheate's examination of the hearing power of a large number of school children was in entire accord with the impression I had gained from very frequent visits to schools. Amongst the backward children and the very large number who fail to keep up with the average of school work we find a large percentage suffering from defects of the special senses, which prove a severe handicap not only in early years but throughout life. I am glad to say that at last this is recognized by our Educational Committee, so that we are now about to appoint a medical officer for the inspection of the children, a prominent part of whose duties will consist in the detection of such errors of sight and hearing as prove detrimental to the child's interests. With the increasing competition among the nations, it will be more and more necessary to ensure that each citizen shall start life with the fullest possible development of his faculties.

A DISCUSSION ON THE LINES OF TREATMENT IN PREVENTING ACUTE MIDDLE-EAR SUPPURATION FROM BECOMING CHRONIC.

I.—WM. MILLIGAN, M.D.,

Honorary Surgeon, Manchester Ear Hospital; Aural Surgeon, Manchester Royal Infirmary; Lecturer on Diseases of the Ear, the University of Manchester.

If justification is required for the selection of this particular subject for discussion, a glance at any of the principal otological clinics of the country will, I think, afford an ample answer. Why is it that we meet with so many chronic cases of septic middle-ear disease in the course of practice? Is it that the virulence of the initial infective process is such as to baffle all attempts to arrest the disease, or is it that the methods of treatment usually adopted in its early stages are inadequate? Undoubtedly cases occur—more especially during the course of one or other of the exanthemata—where the microbic invasion of the tissues is so severe and the general want of resistance on the part of the patient so feeble, that all our efforts to restrain the progress of the disease