selects not only the function of tonicity but also that of conductivity of certain fibres.

Digitalis and the Function of Tonicity.

In my observations on the effect of digitalis on the human heart, I pointed out that when a function such as conductivity or contractility was depressed, digitalis had a tendency to increase this depression. Notwithstanding this I found that while the digitalis depressed these functions a patient's condition may be distinctly improved, and it seems at first sight very anomalous that benefit should be derived from a drug which manifestly injures certain functions of the muscle fibres. But the drug digitalis affects more functions than one, and from such experiments as those of Gaskell already mentioned, we learn that digitalis primarily affects the function of tonicity. In cases of heart failure where there is dilatation of the chambers, the contraction of the heart takes place at a disadvantage, for presumably efficiency of contraction will depend on a certain normal distension of the chambers, and an over-distension or dilatation will interfere with the efficiency of contraction. In such cases the digitalis by restoring the tonicity will reduce the cavity to a size more fitted for the effective accomplishment of its contraction. Herein, probably, lies part of the explanation of why it is that the heart's condition is improved with a depression of certain functions manifestly due to the digitalis.

INEBRIETY AS A PHYSICAL DISEASE.*

By H. W. MANN, M.B. EDIN.,

Resident Medical Officer, Larkfield, Nairn, N.B.

I VENTURE to ask your attention to this question of alcoholism, as it concerns us of the medical profession, and more particularly with regard to its treatment by the hypodermic method by injections of atropine. Perhaps next to tubercle there is no question more pressing or more discussed at the present time than this question of inebriety and the inebriate; and certainly there is no one on which the charlatan and the advertiser, lay and political, have thriven more abundantly. To this may in some measure be attributed the fact that medical men as a rule look sceptically and askance on the "cures" and specifics so lavishly offered in the daily press and monthly magazine.

Happily of late years medical men of note have given this matter scientific study and attention, and by regarding inebriety as a physical disease, in many cases capable of cure, they have rescued it from the advertising quack, who too often exploited the inebriate to his own purposes. The late Dr. Norman Kerr was the pioneer in this, and his work has been carried on by such men as Sir Victor Horsley, Professor Sims Woodhead, Mr. McAdam Eccles, Sir James Crichton-Browne, Dr. Clouston, to mention but a few of the more prominent.

In psychological medicine the great forward step in our times has been recognition of the insanities "as a physical disease with mental manifestations," as a condition cerebral in its essence rather than psychic, and on this recognition the modern treatment of this disease is based. Inebriety is one of the phases or varieties of the insane diathesis, and is akin to the *folie circulaire* or alternating insanity. It has its term of excitement, its depression, and its apparent equilibrium.

Inebriety has its etiology, its pathology, and its treatment. It is essentially an irresistible, uncontrollable, and periodic desire for alcohol. It is not thirst, and while the "crave" is present the appetite knows no satiety. In his book on *Mental Diseases*, Clouston places it under "States of Defective Inhibition," and while conscious of its gravity as a social evil, clearly regards it as a neurosis of defective inhibition, as a lack of that control which is the highest function of the cortical centres.

Briefly considered, the following have a place in the causation of this condition: traumatism, concussion, shock, injuries to the head. Epilepsy is frequently associated with inebriety, and, like it, is a periodic nerve storm. Perhaps no association is more commonly observed than tubercle or the tuberculous habit, and it is remarkable in how large a number of cases a family history of tubercle is found to occur.

Among what may be termed "moral" causes there are the numberless social influences, adversity, prosperity, education imperfect and education to excess, ambition, worry, and all

those undefinable states comprehended in the word "Society." Though in such it is sometimes difficult to discriminate between cause and effect. Lastly, heredity has a prominent place in the etiology of alcoholism, and in this case undoubtedly the children do suffer for the sins of their parents.

The pathology of inebriety has received much attention of late years, and if any one fact in pathology is established, it is this—that alcohol, even in physiological quantities, is a direct nerve poison, affecting the cortical layers in the cerebrum, the cerebellum, and the peripheral nerve endings. In a recent lecture Sir Victor Horsley has pointed out how even by the ingestion of small quantities of alcohol the brain functions are affected, how the reaction time is lengthened, both simple and complex, as estimated by Kraepelin's method. And he has demonstrated that in continued absorption the nerve cell degenerates, the granules become obscured, the nucleus and nucleolus gradually disappear, and the active protoplasm is replaced by pigmentation, until finally the cell is inert and functionless. In the peripheral nerve endings the same toxic process goes on, and so in the confirmed alcoholic there are the pains of neuritis, affecting his peripheral nerves, while the tremulous hands and ataxic shuffling gait indicate cerebellar mischief, and the confused misty intellect corresponds to and is co-extensive with the destruction of the cortical layers in the grey matter of the cerebrum.

Ford Robertson, in his work on *Cerebral Pathology*, describes this destructive process graphically when he compares each successive bout of the dipsomaniac as resembling a hurricane of wind sweeping through a forest, which when it has passed shows to the casual onlooker no change, though the trained eye can see the broken stems, the torn branches, and the disarranged undergrowth. But, unlike the forest, to the alcoholic there is no new growth; no proliferated cells replace those lost.

With such a pathology of destruction it may well seem futile to talk of treatment, but with the 600 million of nerve cells which has been calculated to be the average endowment of the average brain there may be always the possibility of calling up from this vasty deep of subconscious cerebration some reserve and reinforcement of those altered and destroyed.

And so the treatment of inebriety, always granted the abolition of alcohol, amounts to a re-education. It must be borne in mind clearly that no treatment will help an inebriate unless he is willing to help himself. He cannot be cured against his will. If one can but control or abolish the insane impulse for alcohol, then the case resolves itself into a dependence on time and the ordinary processes of assimilation and metabolism. What is destroyed cannot be replaced, but what remains may be utilized.

Various drugs have been used for the purpose mentioned—notably the bichloride of gold. An antiethylic serum was put forward some years ago, but with what success I know not. The treatment by hypodermic injection of atropine combined with strychnine has in my hands proved of great value in relieving the intensity of the appetite, and when thoroughly pushed it does confer an indifference to alcohol. I may be mistaken, but I think it is to Dr. McBride of Norwood that the profession owes the first clear record of this method of treatment.¹ The following notes of cases will illustrate this:

A. R., admitted July 24th, a case of chronic alcoholism with degenerative nerve and tissue changes, ataxic gait, tremulous twitchy fingers, patellar reflex absent on both sides, plantar greatly increased, has had neuritis very badly, great mental confusion of time, place, and person; appetite for alcohol intense and continuous. He was given a calomel purge, and put upon atropine gr. $\frac{1}{15}$, with strychnine gr. $\frac{1}{15}$, given at 9 a.m., 3 p.m., and 9 p.m., with a cinchona mixture sandwiched midway between each dose of atropine. July 28th. Slept well, mental condition improved, quiet, calm, and clear; alcohol crave diminished. July 31st. Much better, complexion clearer, cheerful and pleasant, weight 9 st. $2\frac{1}{2}$ lb. August 7th. Atropine increased to gr. $\frac{1}{10}$. August 23rd. Atropine gr. $\frac{1}{10}$; atropine stopped. October 2nd. Continues well, no relapse, no desire for alcohol, weight 9 st. 13 lb., is on parole.

Another case: August 29th. A.M., drinking for weeks, mainly solitary drinking; for a time he would square up, and then have a deeper and deeper debauch. Confesses his weakness, laments it, and promises reform. Put upon atropine and strychnine with cinchona, as in the other case. September 11th. Continues to improve, appetite for alcohol largely gone, is allowed to cycle out and go to and fro practically uncontrolled; has had no relapse. October 3rd. Improvement maintained and

* Read at the meeting of the Northern Counties Branch of the British Medical Association at Dingwall.

continuous, atropine and strychnine given regularly and to the full physiological limit; alcohol appetite disappeared entirely.

It is a remarkable fact that the dryness and consequent thirst created by atropine is a thirst for ordinary liquids, and is allayed by them, and is not a thirst for intoxicants. The use of the cinchona given as tinct. cinchon. co., with spirits of chloroform and ol. cinnam., is of great value. It cleans the tongue, braces up the appetite, and given in the early morning it assists in allaying the desire which in the morning is at its keenest. Beyond this drug treatment there is the dieting, abundance of vegetable food, and less of flesh meats. Milk in quantity and fish, exercise in the open air, and indulgence in some hobby, especially when that can take the form of any outdoor occupation. I have found sanatogen to be of great value in the early weeks, and when there is, as there often is, gastric trouble with vomiting or sickness, it is in my experience the best of artificial foods, easily retained and easily assimilated. I could give at the risk of wearying you similar examples of the influence of atropine in this combination on the "crave" which is the single real symptom of this disease. Its use is not wholly empirical, and the progress of such cases is well worth inquiry and investigation. It is no secret treatment, and it has the advantage that it is quite in the power of any medical man to pursue it in his own practice. There are, unfortunately, too many cases ready to the hands of the ordinary practitioner for trial, and I would urge upon the members of this Branch to give this method their interested study, and to look upon this condition not so much as a serious social evil, but as a physical disease depending on physical causes, and capable of responding to physical treatment.

Behind this lies the great question of permanence. It is not possible to repair damaged tissue, or to alter a man's character and disposition. No drug can do that, but that many cases of inebriety are curable I am thoroughly satisfied. I personally have too short an experience to permit of saying that the cure is permanent, but I know from a study of the case-books in a well-known Retreat, that cases under this method have been traced, and known to keep well without relapse for more than a dozen years, and the permanency of the result is largely determined by the completeness of the treatment. I had one case, which came to me in April last, in whom the agony of the crave was painfully intense. He had squandered his fortune, ruined his home, and in a fit of depression nearly killed himself. To-day, a sadder and a wiser man, he is steadily and soberly trying to regain something of his lost life, and is happy in his home. Such cases are common knowledge to us all, and it is precisely in such cases that the ordinary general practitioner can by this method, if regularly carried out, benefit his patient and those depending upon him.

It may be necessary for the State or the community to segregate the inebriate. It may even be desirable to adopt Dr. Rentoul's heroic method and sterilize him in the interests of the race, but neither segregation nor sterilization will cure him. But if the impulse, the crave, can be removed, if the inhibition can be made more complete, if the control can be regained, then he may return to the world and take part in its work. It may seem absurd and incredible to lift against this Goliath the tiny hypodermic syringe, the fresh air, and the simple life; but when one reflects how the lancet has controlled the horrors of small-pox and the serum treatment has modified diphtheria, then one may hope that medicine has not yet said the last word regarding inebriety as a physical disease controllable by physical treatment.

REFERENCE.

¹ BRITISH MEDICAL JOURNAL, April 30th, 1904, p. 1006.

SOME TOXIC EFFECTS OF ASPIRIN.

By J. S. DOCKRAY, M.D. VICT., B.Sc. LOND.,
Bishop's Stortford.

As aspirin is a comparatively new drug, very useful in many forms of rheumatism and ill-defined rheumatic pains, and consequently very freely given, the following case, illustrating some of the toxic effects, may not be without interest:

History.—On February 8th, 1905, I was called to see a lady of about 50 years of age suffering from sharp pains in the regions of the loins. The temperature in the mouth was 100° F., the patient was extremely restless and excitable, far more so than the pain accounted for, but the pulse was steady, of good tension, and the pulse-rate was 80. This was

about eleven o'clock in the morning, and she informed me that she had taken a dose of 10 gr. of aspirin a short time before. As she had taken it for lumbago previously, and it had always relieved her rapidly, I ordered the drug to be continued in the same doses, 10 gr. every three hours. Next morning (the 9th) the pain had disappeared, the temperature was 97.6° F., but the patient complained of singing in the ears and drowsiness. The excitability of the day before had disappeared. The pulse was unaltered. The drug was still continued. On the 10th I found the following symptoms: Numbness of the gums, tongue, and pharynx, causing a great difficulty in swallowing even fluids, and a sensation of impending suffocation and choking. There was no swelling of the mucous membrane, but it looked dry and glazed, and in that covering the gums and tongue sensation was practically lost, this anaesthesia being much more marked on the right side than the left; the teeth were not tender, but the patient said she felt as if she had lost them; very severe pain in the right ear but no tenderness or swelling over the mastoid; nausea but no vomiting; temperature subnormal, and the drowsy condition of the day before had now given place to one of extreme excitability and restlessness; the patient was constantly moving up and down in the bed and never still a moment; the pulse was unaltered. On inquiry, I found that she had in a previous attack of muscular rheumatism been ordered 20 gr. of aspirin every three hours for three doses and then three times a day for two or three days. After taking 120 gr. there was such a marked feeling of numbness all over the body, accompanied by a feeling of intoxication, that the drug was stopped.

Progress.—As evidently the uncomfortable symptoms were all due to the aspirin, I stopped it at once. More for the sake of doing something, I ordered warm olive oil to be dropped in the ear and warm linseed-meal poultices over the ear. No drugs were ordered, but the patient was advised to drink plenty of fluid, preferably hot water and hot tea. Had not the pulse been so good I should have ordered strychnine. At 8 p.m. the same evening the patient felt better. She was sleepy and quiet, but the pain in the ear was very troublesome. There was now also pain in the supraorbital region on the right side, with distinct yellow vision again confined to the right eye, vision in the left eye being quite normal. The nurse in charge stated that there had been distinct drooping of the upper right eyelid during the day, but I did not notice it myself. There was no squint, and the pupils were equal and moderately contracted. There was no flushing or pallor of right side of face. Temperature 100°, pulse good and steady.

February 11th. The patient much better. The right ear was much less painful, but completely deaf, and there was a slight discharge of blood-stained fluid; no yellow vision; no ptosis; severe pain was still felt in the supraorbital region, and now also along the facial and mental nerves. Total loss of smell and taste was now noticed for the first time, but there was no pain in the tongue. For the first time, also, the patient complained of shooting pains down the right arm and leg. The urine was very pale and greatly increased in quantity; specific gravity 1000, acid, no albumen or sugar, no coloration with FeCl₃. Temperature 100° F.

February 12th. Again much better. Temperature normal. The pain in the ear and the neuralgic pains in head and face were much less. The smell and taste had returned slightly, but the mucous membrane of the mouth was dry and scaly. The difficulty in swallowing was almost gone, and the patient could now take nourishment comfortably.

Result.—During the next two days recovery was rapid, the mouth became moist, desquamation occurred over the tongue and lips, and the smell and taste became normal. On February 16th she was practically well, with the exception of the total deafness of the right ear and some supraorbital neuralgia. There was no more discharge from the ear, and no treatment was carried out, as the hearing gradually returned, and by the end of the month was quite well. The neuralgia was rather persistent, but disappeared when the hearing was restored.

REMARKS.

The interesting points to notice in this case are:

1. The marked idiosyncrasy to the drug. The patient had only taken 100 gr. in all, in 10 gr. doses.
2. The marked effect the drug had on all the sensory cranial nerves, causing, when the patient was most fully under its influence, numbness and anaesthesia, followed, as the patient was recovering, by pain. Except for the slight ptosis, which only lasted a few hours, and only occurred at the very height of the attack, there was no loss of muscular power. The facial muscles were not affected, there was no squint, and the difficulty in swallowing was probably due to the loss of sensation of the mucous membrane. Most cases (1 and 2) of reported intoxication by this drug differ from this one in that the chief symptom was a transitory oedematous swelling of the face and scalp. No mention is made of any effect on the sensory nerves.
3. The heart was not affected at all. Had not the pulse remained good and steady compared with the extreme restlessness of the patient, the case would have assumed an alarming aspect. Franke² says, in reporting his own case, that his pulse-rate "rose to 60"—not a very alarming or very high pulse-rate. Barnett³ records quite recently two cases where a single dose of 15 gr. caused "violent palpitation, difficult respiration, a feeling of extreme weakness and gradually approaching unconsciousness"—these symptom