Records

OF

THE LOCK HOSPITAL.

BY

HENRY LEE, Esq., F.R.C.S.,

SURGEON TO THE HOSPITAL.

V.—THE CALOMEL VAPOUR BATH.

FOR many years past, I have discontinued the use of mercury internally in the treatment of syphilis at the Lock Hospital. The class of patients which there present themselves have their constitutions often impaired by excesses of various kinds; and I became very early deeply impressed with the belief that such patients would ill bear long and protracted courses of mercury administered internally.

Mr. Pearson's experience led him to the conclusion that, when mercury is given under the direction of an unskilful man, it will often fail in effecting the cure of syphilis; and that then the complaint will be exasperated by it, and rendered more intractable than if no mercury had been given.* Mr. Pearson says, "a long course of observation has convinced me that, if this alarming truth were as closely understood, and as deeply impressed upon men's minds, as it deserves to be, the great number of mutilated victims, whose sad condition excites horror, while it calls for commiseration, would be reduced within much narrower limits than I can venture to express." It is true that in the present day there is little danger of mercury being used in the way it was in Mr. Pearson's time; and, consequently, the very severe results of the medicine to which he alludes are seldom witnessed. But there are other effects consequent upon the internal administration of mercury, which require the most thoughtful consideration on the part of the surgeon.

In Sir Ranald Martin's admirable work, *On the Influence of Tropical Climates*, we read that "more than sixty years ago, Dr. Dick, an intelligent physician of Calcutta, noted that 'liver-attacks very often succeeded long courses of mercury.' Dr. Girdlestone made the same observation in Madras in 1781-83. He ascribes hepatic disease to the abuse of calomel, even when used 'as a prophylactic'; but not so to mercurial frictions, which, however long continued, did not, he thought, produce the same results." Sir Ranald Martin relates the following case: "Dr. — had never been in a hot climate; and he stated that his case had originally been one of syphilitic ulceration treated without mercury. Secondary and tertiary forms of syphilis followed; and for these, three protracted courses of mercury were had recourse to, with no benefit. When I saw him, his constitution was entirely destroyed, and his liver was of enormous size, having become so after the last mercurial course."

Again, with regard to the internal administration of mercury in the treatment of syphilis, Sir Benjamin Brodie says: "You may patch up the disease by giving

the remedy internally, but it will return over and over again."

It will, indeed, seldom happen that in private practice the surgeon can induce his patient to continue the internal use of mercury sufficiently long to cure real syphilis. The patient either "catches cold," or has an attack of diarrhoea, or finds that he is losing flesh, or feels languid and uncomfortable, and incapable of much bodily or mental exertion; and then he discontinues his medicine, and, feeling subsequently better, is convinced that he acted properly. But, as observed by Sir Benjamin Brodie, if, in a case of well marked syphilis, a short course of mercury only be given, "the disease is sure to return; you have then to repeat it; and again the disease reappears. Thus, you have repeated courses; and not only is the system weakened by the mercury, but the disease, when it does return, assumes a more formidable character than before." (*Lectures on Pathology and Surgery*, pp. 248-9.)

To remedy the evils attending the prolonged internal use of mercury, the plan of introducing the medicine into the system through the skin by means of mercurial frictions was introduced; and although this plan, like other modes of administering mercury, was carried to a great excess, and much abused, yet, as practised by the late Mr. Pearson, when, by mature experience, he had learned the proper use of the remedy, it proved by far more successful than any other plan previously employed.

By introducing mercury into a patient's constitution in this way, by inunction, its deleterious action upon internal organs is avoided. The amount absorbed into the blood produces its influence equally throughout the system, and is not conveyed direct to the liver, as when the medicine is administered internally.

But, great as the advantages of the inunction of mercury are as compared with its internal administration, it, nevertheless, is attended with certain inconveniences which prevent its very general use. The inunction of mercurial ointment, so as to ensure the proper effect of the remedy, requires considerable labour and perseverance on the part of the patients; and it is with difficulty that they can be induced to continue its use for any length of time. It sometimes produces a pustular eruption on the skin, which proves troublesome. It is a dirty remedy, and patients in the upper class will avail themselves of any excuse to avoid its use.

Fumigation of the surface of the body by means of certain mercurial preparations, possesses the advantages of inunction without some of the objections to which this is liable. But, like other modes of using mercury, it was tried in a variety of ways before a safe and efficient mode of administration was adopted.

In the years 1786-7, Mr. Pearson had a fumigating machine constructed according to the directions given by M. Lalouette. This apparatus, although it was thought to be new at the time, differed in no material respect from that described by Nicholas de Blegny in the year 1683.

Mr. Pearson made a considerable number of experiments with this fumigating machine, and found that the gums became turgid and tender very quickly, and that the local appearances were sooner removed than by the other modes of introducing mercury into the system. But, to counterbalance

* A similar remark may be made with regard to the prolonged internal administration of mercury in any form to those whose constitutions have become impaired.

these advantages, it was found that the mode of treatment adopted induced debility, and that ptalism was often excited rapidly, and at an early period. Mr. Pearson found that he was, consequently, obliged often to discontinue his course of treatment.

Sir Benjamin Brodie's experience coincided with that of Mr. Pearson. He found that it was difficult to regulate the mercurial action; and he observes that by using mercurial fumigation "you may affect the system too much or too little; and you may be taken unawares by the patient's gums becoming all at once excessively sore." (*Lectures on Pathology and Surgery*, p. 246.)

From observations and comparative experiments which I made at the Lock Hospital during the years 1855-6, I feel satisfied that the irregular results noticed by Mr. Pearson and by Sir Benjamin Brodie, depended upon the difference in the chemical composition of the powder used for the purposes of fumigation, both before and after it was raised into a state of vapour. The gray oxide of mercury (the preparation generally used) varies much in colour as obtained at different shops. Some specimens will not volatilise at all at the temperature produced by an ordinary spirit-lamp under a metallic plate. Other specimens of a lighter colour volatilise quickly enough. When the darker specimens are sublimed, they are decomposed in a greater or less degree. A deutoxide of mercury is formed by the addition of an equivalent of oxygen from the air; and if the temperature be much increased, then the oxygen is driven off altogether, and metallic mercury is sublimed. Under these circumstances, with a mercurial preparation of uncertain composition, and undergoing different changes according to the degree of heat applied, there is no wonder that very different effects should have been produced in different cases. With some samples of the gray oxide it is necessary to use a considerable quantity of the powder in order to ensure any effect; with other samples, the same quantity produces much more action than is desirable. The gray colour of the powder depends upon the admixture of a certain proportion of calomel with the protoxide, and the temperature at which any particular specimen will volatilise will depend upon the relative proportions of the two. The bisulphuret of mercury, again, which has been extensively used for the purposes of fumigation, gives off, when exposed to heat, a vapour, probably the sulphurous acid gas, which has sometimes caused very considerable irritation of the lungs; and all forms of mercurial fumigation have, in consequence, been condemned. Now, all the inconveniences above-mentioned may be avoided with certainty, by using a mercurial preparation which is always of the same chemical composition, which does not irritate the lungs, and which is not liable to be altered by an increase of temperature. Such a preparation is calomel. We have here a definite chemical compound. It is altered in composition, neither by heat, nor by moisture, and may be relied upon when used for the purposes of general fumigation, for producing its action, as certainly as any medicine administered internally. A very small quantity (as compared with the other mercurial preparations) will ensure the required effect, and this may be regulated and controlled with great precision.

[To be continued.]

Lectures

ON THE

DIAGNOSIS AND TREATMENT OF DISEASES OF WOMEN.

DELIVERED AT ST. MARY'S HOSPITAL MEDICAL SCHOOL.

BY

GRAILY HEWITT, M.D.LOND., M.R.C.P.,

PHYSICIAN TO THE BRITISH LYING-IN HOSPITAL; LECTURER ON MIDWIFERY AND DISEASES OF WOMEN AND CHILDREN AT ST. MARY'S HOSPITAL MEDICAL SCHOOL.

DISCHARGES FROM THE GENERATIVE ORGANS OF NON-SANGUINEOUS CHARACTER. (*Continued.*)

GENTLEMEN,—The diagnostic significance of certain other qualities of these non-sanguineous discharges remains to be considered.

Sanious Discharges from the generative passages are not unfrequently the subject of observation; that is to say, there is a discharge of a reddish tinge, not hæmorrhage, but evidently containing a certain admixture of blood-elements. In women the subjects of profuse menstruation, as the discharge of blood is becoming less, there is generally to be observed a period when there is sanious discharge. The most ordinary causes of that kind of sanious discharge to which our attention is most commonly directed are, the presence of morbid growths within, or organic disease of, the uterus, malignant ulceration of the os uteri, etc.; and we find combined, leucorrhœa, and very slight but continuous hæmorrhage. In polypus of the uterus, such sanious discharge, alternating with hæmorrhages or with colourless leucorrhœal discharge, is observed. Whatever, in fact, is capable of giving rise to hæmorrhage may occasion discharge of a sanious character. It is important to be aware of the fact that the presence of a more or less continuous sanious discharge is a condition of things requiring careful examination.

Offensive Discharge. This quality of the discharge is important in reference to the determination of the disease present in certain cases; but it is one on which too much reliance must not be placed, or serious mistakes may be made. Discharges of an *offensive character* have been usually considered as absolutely indicative of the existence of cancer. Now, it is true that, in almost all cases of cancer of the uterus, there is to be remarked a particularly offensive odour of the discharge proceeding from the vagina; but it is also true that it may be absent. In cases of profuse purulent leucorrhœa accompanied by hectic and a generally low state of the constitutional powers, the discharge, when retained in the vagina for any time, is apt to be offensive, particularly if the retention of the discharge be favoured by a contracted state of the ostium vaginæ. (Tyler Smith, *op. cit.*, p. 93.) The offensive nature of the discharge is so often looked upon as a material point in the diagnosis of cancer uteri, that a good deal of practical importance attaches to this observation. The smell of cancerous discharge is peculiar; there is not simply fœtor, but fœtor of a peculiar quality. In fact, the smell is so peculiar that it can hardly