of our human nature; and the extent of this ignorance and amount of this degraded weakness must, alas! be measured by the amazing spread of quackery amongst the masses of mankind. The homœopath talks of infinitesimal doses, and flourishes a few unmeaning cabalistic phrases; and finds abundant listeners, because he flatters, he promises, and, above all, gabbles to the people of that which is utterly incomprehensible to them.

If the bright face of medicine be ever obscured by the shadow of quackeries and delusions, it can only be in proportion as our professional morality has grown weak. But woe is it to our art when men are found amongst us who are willing to shake hands with a wicked imposture, because, clothed in the garb which it has surreptitiously stolen from medicine, this imposture has found favour in the eyes of the wealthy! Woe is it to our art when men barter their professional honour for the sake of great gains, or juggle their consciences with jesuitical rules of ethics!

But, gentlemen, thank Heaven! our body is sound at heart. If a loose professional morality be here and there seen floating about amongst us, we know and feel, as members of a medical corporation, that our path and course are clear and fixed amidst this scene of ignorance and vicious reasonings. It is no fault of ours if deception in the cure of diseases have a strong hold on the human mind. Let us only keep our hands clean of the contagion of pandering to it. We have no reason to fear the conclusion. Truth is stronger than deception.

Illustrations

or

HOSPITAL PRACTICE:

METROPOLITAN AND PROVINCIAL.

ST. MARY'S HOSPITAL.

CASES OF MALARIOID DISEASE: WITH COMMENTS.

By C. HANDFIELD JONES, M.B., F.R.S., Physician to the Hospital.

CASE I. W. C., aged 22, an excavator, was admitted Sept. 16th, 1861, having been ill four months. His illness came on gradually. His aspect is that of a man labouring under serious disease. He is anæmic; his speech is slow and hesitating, and his whole manner of the same kind. He seems to get worse every week. He complains of great debility and headaches at night. He is very ill in the morning, can hardly draw his breath, and has cramps in his legs. Tongue glazy; skin cool; pulse feeble. The heart and lungs are sound. There is not much cough nor expectoration. He can take food fairly well. He never has had ague, nor been in the way of it, that he is aware of. He was ordered to take five grains of quinine three times a day.

Sept. 19th. He feels a deal better. His breath is better; the urine is very clear, not albuminous. The medicine was continued.

Sept. 23rd. He is improved. He has pain over the left orbit, in the situation of the frontal nerve. He was ordered to have, three times a day, ten grains of citrate of quinine and iron, and ten minims of tincture of nuxvomica. He ceased attendance.

The general depressing, enfeebling character of obscure malarious disease, and its special implication of

the cerebro-spinal nervous system, was here well marked. The episode of brow-ague aids in completing the picture.

CASE II. A. S., aged 37, coachman, was admitted July 29th, 1861, having been ill three or four months. He feels very weak. The tongue is very coated. He has pain at the mid-sternum, worse after food; no thirst. His appetite is bad in the morning, ravenous by midday. The epigastrium is not tender. He has cough and expectoration in the morning. Pulse natural. He cannot sleep at night, and sweats very much. He was ordered to have four grains of disulphate of quinine three times a day.

He improved steadily, and ceased attendance Aug. 29th, having had iron and quinine the last ten days. The perspirations at night had almost ceased, and he had gained strength.

This case differs from the preceding, in that the disorder was less restricted to the cerebro-spinal nervous system, and involved the sympathetic; there was, in fact, bronchial and gastric catarrh. The malariousness was evidenced by the nocturnal sweating without existing organic disease—a symptom on which I have before dwelt, as highly significant of such affections. It is highly interesting to note in these two cases the existence of general debility—of special depression of nervous power; and then, in the latter, the same as manifested in the paralysis of certain vaso-motor nerves, and its results in bronchial, gastric, and cutaneous exudation.

CASE III. G. H., aged 36, was admitted Aug. 16th, 1861. He is a blacksmith, residing at Sheerness. He has never had ague or fever. He is very anæmic, and looks spare and weakly. He has been ill since last November, when he was seized with pain in the back during sleep, having been perspiring copiously before. He is extremely weak, and has drenching night-sweats. His legs are so weak that he cannot walk without crutches. He was treated till Aug. 29th with tincture of sesquichloride of iron. The drenching sweats continued, but he could walk better. He was then ordered five grains of quinine three times a day. He had ordinary diet and porter from the first.

Sept. 5th. He is improved, sweats less at night, and is stronger; his appetite is better. The spleen, liver, and lungs are normal. A blister was applied to the loins on Aug. 22nd, with benefit.

Sept. 9th. He is much improved; has no night-sweats now. He was ordered to take one-sixteenth of a grain of strychnia and four grains of sulphate of iron three times a day.

Sept. 16th. He is not so well without the quinine, but has pain in the back. The legs are not improving. The quinine was repeated, and a blister was applied to the loins.

Sept. 23rd. He was improved.

Oct. 12th. He looks bigger and better, is strong in himself. He can walk well; but his ankles feel weak, and feet tender. He went out soon afterwards, apparently quite well.

The locality where the patient resided, the debility without sufficient cause, the drenching night-sweats, and the good effect of quinine, indicate very strongly the malarious nature of his disorder. It may be surmised that the poison had caused congestion, or low inflammatory action, of the membranes of the cord; and that this was concerned in part in causing the incomplete paraplegia. The two blisters acted beneficially in removing this congestion. As, however, the good effect of the quinine was evident, I think it must be granted that this remedy exerted a direct action upon the spinal nervous centre, increasing its tissue-power, and making it more capable of fulfilling its normal function. Compare a more acute case recorded by me in the BRITISH MEDICAL JOURNAL, Nov. 28th, 1857. Such instances as the above

illustrate the point, already notorious, that the action of malarious poison may be general or local. It may involve the whole nervous system, and produce an ague, or limit itself to a frontal or some other nerve. It may cause an hemiplegia, or, as in this case, a paraplegia. It may affect the heart prominently, as in syncopal tertian; or concentrate itself on the large intestine, as in dysentery.

CASE IV. W. D., a boy, aged 8, was admitted Sept. 6th, 1861, having been ill, on and off, since he had an attack of the same kind fifteen months ago. He has blue eyes, light hair, an intelligent face, a rather prominent forehead. He is very fond of reading. He has lost much flesh. At times he has a "nasty dry cough". He complains of uneasiness at the lower part of the sternum. There is good breathing in the backs; some crepitation is audible under the right clavicle. The abdomen is normal and painless. The skin is hotish, with perspiration on the face and forehead. He does not sleep well at night; rambles and talks. He varies much; one week, his mother says, he is all life and spirits; the next, much depressed. He was ordered to take three minims of dilute hydrocyanic acid in half an ounce of citrate of potash mixture three times a day.

Sept. 9th. He lies sleeping very much, is pretty quiet at night, and takes his food (broth diet) fairly. Râles are heard all over both fronts and backs. He was ordered to have a grain of disulphate of quinine three times a day.

Sept. 12th. He is evidently better; his skin is cool; he sleeps less by day, and is quiet at night. But little râle is heard in the chest. He was allowed half ordinary diet.

Sept. 16th. He seems quite well, running about. A drachm of cod-liver oil was ordered to be taken three times a day. The quinine was exchanged for citrate of iron and quinine, with chloric ether; and he went out on Sept. 23rd, apparently quite well.

Familiar as I have been for some years with the various forms of malaroid disorder, there was much in the above case to make one very suspicious that one had to deal with a much less hopeful condition. The physical and mental constitution of the child, the emaciation, the cough, the subclavicular râle, and the febrile movement, pointed very much in the direction of tuberculosis, and made me forbear from tonics at first. When, however, these were brought into play, the response of the system was such that the true nature of the disorder became at once apparent. Truly may it be affirmed in this case, that "naturam morbi remedium ostendit".

CASE V. W. Y., a boy, aged 8, was admitted July 15th, 1861, having been ill one week. He is losing flesh fast; has no appetite, and has a bad cough. The skin is cool; tongue clean. He faints away at times. He is "like a fire at night", so feverish. Two grains of quinine were given at 12 M., and at 6 and 8 P.M. He was much better in a week, and in fourteen days was reported to sleep well. Five grains of citrate of iron and quinine three times a day were now given, which he continued till Aug. 26th, when he was fairly convalescent.

This was a not very severe case of nocturnal fever, with general depression and emaciation, which, on account of its being so markedly under the control of quinine, I have termed malaroid remittent. For other and more severe examples of the same affection, I refer to my pamphlet on *Neurolytic Affections*, etc.

CASE VI. H. W., a boy, aged 2, was admitted July 8th. He has been ill, on and off, from his birth. He has been complaining during the last week of his head; he lies down, and puts his hand to the right side of it. He has no rest at night. The head is not hot; the bowels are relaxed; the urine is thick and dark coloured. The following draught was ordered to be taken three times a day.

R. Tincturæ opii m̄ij; potassii iodidi gr. j; tincturæ cinchon. m̄x; sodæ carbonat. gr. ij; aquæ cassiæ 3ij.

July 11th. He sleeps much better, and does not complain of his head. He is "nothing near so cross". His appetite is poor. After this he took a grain of quinine three times a day; and, at a later date, citrate of iron and quinine and cod-liver oil; and improved very much. He remained under observation till Sept. 26th, and had no relapse.

CASE VII. J. T., a boy, aged 2, was admitted Aug. 26th, having been ill fourteen days. He has had diarrhœa lately. His aspect is distressed. He has all his teeth. The appetite is bad. His bowels are properly open. He is very thirsty. He screams by night and day, and seems to have pain in the face and head and feet. He is constantly whining and crying. He is cool now, but very feverish at night, and is worse then. He was ordered to have half a grain of iodide of potassium, and two grains of citrate of quinine and iron in two drachms of water, three times a day.

Aug. 29th. He is a great deal better, and quiet.

Sept. 9th. Pain recurred on the 7th. He has had no medicine the last four days. The medicine was repeated.

Sept. 16th. He is better. The medicine was repeated. He did not attend again.

The last two cases are both instances of a rheumatoid head-affection, probably implicating the dura mater. The symptoms of this disorder may often suggest grave suspicions of the existence or the approach of tubercular meningitis. However, the absence of vomiting and convulsions, and of drowsiness, the irritability, and, above all, the non-progressive course of the affection, indicate usually its true nature. Dr. Coldstream has already pointed out the advantages of using iodide of potassium in various inflammatory affections of the head, especially in cases of less sthenic character. (*Edin. Med. Journal*, Dec. 1859.) He does not, however, seem to regard the remedy as specially applicable to rheumatic cases; and in this his experience justifies him. Rheumatism of the head, though familiar to us in adults, is not much noticed, if at all, in works on children's diseases. Dr. West does not mention it in his valuable *Lectures*. I am, however, quite sure that the practitioner will find it well worth while to bear in mind the possibility of the occurrence; and that in many instances, where leeching and calomel would avail little, the use of that great remedy, iodide of potassium, will be attended with gratifying success.

I am anxious, in conclusion, to say a word or two with regard to the remittent fever of children. This is generally regarded as identical with the enteric of adults; and no doubt there are many cases in which this opinion is correct. There is, however, also, I am convinced, another form, which, from its general features, its prolonged course, and its controllability by quinine, approximates closely to malarious fevers. It is not a very common disorder, not so frequent lately as it was two or three years ago; but yet I have met with it much oftener than with the graver (perhaps typhoid) form. To be aware of its possible occurrence, and to be able to distinguish it, is certainly worth while, as it is positively curable.

THE WAGES OF HUMANITY. All of the prisoners speak in terms of the deepest gratitude of the kind attention of Dr. Higginbotham, a well known physician of Richmond. He was constant in his attentions to them from first to last—so much so, indeed, that his loyalty was suspected, and he was placed under arrest by the military authorities. He was subsequently released, however, by the civil authorities, and resumed his kind ministrations.