such serious danger by its locality, but rapidly recovered from by reason of the slight morbid alteration which has taken place, I show you here a larynx taken this week from the body of a man who died of pulmonary consumption. See how extensively it is disorganised. The mucous membrane covering all the upper part is white, thick, and ragged; just below the vocal cords on the right side, there is a ragged deep ulcer, which has bared the arytenoid cartilages, and caused them to become necrosed by killing the perichondrium; a piece of the necrosed cartilages projects into the bottom of the cavity. There is another superficial ulcer in a corresponding place on the opposite side. Yet here the symptoms referable to the larynx were of the most minor importance. The man's voice was indeed hoarse and weak, but not more so than you find in most sufferers dying with a very large vomica in one lung and the other filled with crude tubercles, which we found in this autopsy. There was a considerable secretion of pus also, and pain in the larynx when pressed; but no difficulty of breathing, in whatever position the patient lay. Consequently, no local applications were made to the organ. There was no demand for local medication, and certainly it would have been useless to the prolongation of life. The deficiency calling for restorative treatment was the important one of chronically deficient nutriment to the whole system. The larynx was ulcerated and degenerated, for the same reason that the kidneys were shrunken and granular, and that the whole person was emaciated to a skeleton: sufficient aliment was not absorbed to compensate waste. A despairing effort was made to supply this by endeavouring to restore digestive and absorbent powers to the alimentary canal by means of quinine and iron, and to supply muscle and fat with meat and cod-oil. But in vain. Sentence of death had long before gone forth; and it was difficult to see that it was in any degree delayed by treatment, however appropriate. My object in quoting this case now is not by it to commend the remedial agents employed, but to draw your attention to the rule, that the immediate danger of diseases is mainly traceable to their locality; the final danger, to the extent and degree of the pathological changes.

I would therefore lay it down as a rule that, in cases where the danger is immediate, the attention of the medical attendant should be directed to locally active remedies; where it is more remote, it should be directed to those of general agency. Thus in acute laryngitis you are to think of the larynx alone, to the temporary neglect of other parts; in chronic laryngitis, of the general nutrition, in preference to the local injury.

GRINDING. The tutor 'crams' his pupil, not with knowledge, but with promising bits of information adroitly selected, and teaches him how to reproduce them by the most rapid process and in the most specious shape. He carefully studies the papers that are set, divines, if he can, the theories or tastes of the Examiners, makes 'shots' at the subjects most likely to 'pay,' and discovers or constructs short cuts to the knowledge required. The proficiency obtained under such a system is not likely to be genuine, but whether it be real or simulated signifies little, so long as the object is gained. If the pupil appears in the list of the successful candidates, he is satisfied, and his tutor too.

Lectures

ON THE

DIAGNOSIS AND TREATMENT OF DISEASES OF WOMEN.

DELIVERED AT ST. MARY'S HOSPITAL MEDICAL SCHOOL.

BY

GRAILY HEWITT, M.D.LOND., M.R.C.P.,

PHYSICIAN TO THE BRITISH LYING-IN HOSPITAL; LECTURER ON MIDWIFERY AND DISEASES OF WOMEN AND CHILDREN AT ST. MARY'S HOSPITAL MEDICAL SCHOOL.

DISORDERS OF MICTURITION. (*Continued.*)

GENTLEMEN,—We have now considered the cases in which micturition is difficult, and those in which it is painful. The next class of cases are those in which

c. Micturition is frequent. There is, perhaps, no one diseased condition of the vagina, uterus, bladder, or adjacent organs, which may not, at one time or other, give rise to frequency of micturition, to say nothing of the varying conditions of the urine which may occasion the same phenomenon. Frequency of micturition can hardly, then, be considered as characteristic of the presence of any one diseased or altered condition.

Frequent micturition is often an early sign of pregnancy. During the first two months of gestation in primiparae it is very generally present. Towards the latter end of pregnancy also, it is pretty frequently observed. In hysteria, frequent micturition is often present during the attacks.

Displacements of the uterus may occasion frequent micturition; but more often difficulty and pain during micturition are produced thereby. In retroflexion of the uterus the micturition was most frequent during the night in two cases observed by Ashwell. In prolapsus uteri, there is frequency of micturition, especially during its early stages; it is diminished by the horizontal posture. *Ovarian* or other pelvic tumours occasion frequent micturition, owing to pressure on the bladder before remarked. Urinary difficulties are more frequently present during the early than the later stages of these tumours; when larger, they rise out of the pelvis, and the patient suffers less. One of the most important causes of frequent micturition is retroflexion of the gravid uterus, a condition in which urinary difficulties are rarely absent. There may be difficulty alone; but more generally difficulty and frequency of micturition are noticed; the latter may alone be observed. *Organic affections* of the uterus, as cancer, fibroid tumour, polypus uteri, or simple hypertrophy, or an inflammatory or hyperæsthetic condition of the organ, may, each of them, give rise to frequent micturition. Cancer would be recognised, after its very earliest stage, by the other symptoms present; fibroid tumour and hypertrophy might be unaccompanied by other symptoms; in polypus, menorrhagia is not often absent. Tenderness and pain are the characteristic symptoms of the inflamed or irritable uterus. Pressure on the bladder, and consequent frequent micturition, may be produced by abscess in the cellular tissue between the bladder and vagina, or by effusion of blood into the peritoneal cavity around the uterus in peri-uterine hæmatocele.

Dysmenorrhœa is often associated with frequent micturition; the tenesmus of the uterus extends to the bladder. But certain conditions of the bladder itself may give rise to frequent micturition. Calculus of the bladder, cystitis, cancerous disease of the organ, the condition known as the "irritable bladder," occasion this symptom, which is, moreover, observed in the early stage of the affections in question. The presence of the blood in the urine occasions frequent micturition, as do also various disordered conditions of the urine. Irritation propagated from the kidneys, when these organs are diseased, or from the rectum, as when hæmorrhoids are present, may occasion frequency of micturition. Cases in which hæmorrhoids have to do with disturbances of the function of the bladder not seldom remain for some time obscure.

Inflammation of the urethra, as in gonorrhœa, is a cause of frequent micturition: the urine is then passed in drops, with scalding pain. *Vascular tumour* of the meatus occasions frequency of micturition distinguished from inflammatory conditions by its chronicity.

d. Micturition Involuntary. The conditions under which this symptom may be observed are the following.

Fistulæ in the Vesico-Vaginal Septum. In such cases, the patient has hardly the slightest control over the evacuation of the bladder, the urine escaping from the bladder by the unnatural opening as fast as it is secreted. The condition dates from a definite period, at which time the vesico-vaginal septum was injured, and since when there has been involuntary micturition. The formation of these fistulæ is generally connected with the act of parturition; but *syphilitic* or *cancerous* ulceration may be the source of the evil. If the existence of fistula be suspected, the vagina and bladder must be carefully examined.

There are cases on record in which involuntary micturition was produced by the existence of a *vesico-uterine fistula*. Here the symptoms are very peculiar, and the nature of the case would be easily recognisable on careful study of the history of the case, combined with examination of the vagina. If the urine were seen issuing from the os uteri, this would conclusively determine the diagnosis.*

At the latter part of *pregnancy*, micturition is often involuntary, either entirely, or when the patient is in certain positions. The diagnosis of this would be obvious enough.

Retroflexion of the gravid uterus generally occasions great distension of the bladder; and not unfrequently a case of this kind comes before us in this form:—the patient complains of involuntary micturition; and, on examination, it is found that the condition really present is one of *retention of urine*, produced by retroflexion; small quantities from time to time escaping, owing to the extreme distension of the bladder. The period of pregnancy at which this distension of the bladder most commonly occurs is the fourth month. The distension of the bladder was supposed by William Hunter to be the cause of the dislocation of the uterus. Dr. Tyler Smith has shown that the retroversion (in many cases, at all events) is the primary evil; the fact being, that the

retroversion existed before the pregnancy occurred. (See vol. ii. of the *Transactions* of the Obstetrical Society.)

When the bladder is paralysed partially or entirely, as in the course of fevers, etc., great distension of the organ and *overflow* may occur, as in the case of retroflexion just noticed.

After parturition there is often involuntary micturition for a few days, which may extend to weeks or even longer. The muscular structure of the urethra has in such cases undergone undue pressure and injury during the act of parturition. In women who have had large families, the neck of the bladder occasionally becomes thus permanently weakened, and the control over the bladder is subsequently always imperfect.

Tumours of the ovaries now and then produce involuntary micturition; the tumour drags on the bladder, and mechanically interferes with the action of the sphincters. This effect is more commonly witnessed when the tumour is large. Other tumours in the pelvis or neighbourhood may have a like effect.

Great hypertrophy of the nymphæ was a cause of incontinence of urine in a case recorded by Breslau. Owing to the traction of the enlarged nymphæ, the action of the sphincter was interfered with.

Cancer of the uterus may extend to the neck of the bladder, and give rise to involuntary micturition (Kiwisch. See West, vol. ii, p. 195.)

Congenital defect of power over the sphincter of the bladder is very rare, but the possibility of its existence should not be forgotten. Congenital incontinence of urine may be due to *imperfect formation of the urethral canal associated with epispadias*, of which a very interesting case is recorded by Dr. Röser (*Wurt. Corr. Bl.*, 1861; Schmidt, *Jahrb.*, vol. cxii, p. 47). The case was that of a young woman, aged 18, who had had incontinence from birth. The clitoris consisted of two parts; the upper and anterior portions of the orifice of the urethra were wanting, and the orifice itself was very large. A cure was obtained by bringing the separated halves of the clitoris together by a plastic operation.

II.—MICTURITION NOT POSSIBLE.

In cases where the patient is absolutely unable to pass urine, it is evident that there is either an impediment to the escape of the urine from the bladder, or that there is no secretion from the kidneys. In other words, the case is one of *retention* or of *suppression of urine*. In the distinction of these two conditions, it is to be remarked that retention is, as a rule, accompanied by a desire to evacuate the bladder, which is for the most part absent in cases of suppression: the exception is in reference to cases of paralysis of the lower extremities, and some other instances where there is *sensational* as well as *motor* paralysis. Cases are rare in which there is a possibility of taking suppression for retention; but it might prove a dangerous mistake, and one more within the limits of possibility, to overlook retention, and set down the condition as one of suppression. Such cases occur in connexion with the presence of diseases producing great prostration, fevers being the chief of these. The patient may for a considerable time have no evacuation from the bladder; and, this circumstance escaping attention, the bladder is allowed to go on increasing in size. The obscurity of the case is often increased by the fact

* A most interesting case of this kind is related by Dr. Leishman, in the "Glasgow Medical Journal," October 1861. The patient in this instance could only retain urine within the bladder when lying on the side.

(previously alluded to) of a small quantity of urine escaping from time to time from the distended organ, the retention all the while persisting to a dangerous degree. The fact that the patient has expressed no desire to evacuate the bladder must be disregarded; and, after a certain time has elapsed, an examination should be made, in order to ascertain whether the condition present is one of retention or suppression. The definitive decision between retention and suppression depends, then, upon the results of examination.

Cases in which retention is combined with involuntary micturition have been already disposed of.

Retention produced by inability to evacuate the bladder, coupled with distress and strong desire for the same, may arise from mechanical pressure on the neck of the bladder, of whatever kind. Whatever, in fact, may produce mechanical difficulty may produce also retention. *Fibroid* tumours of the uterine wall, enlargement of the uterus by fluid, etc., act in this way, as do also, but more rarely, *ovarian tumours*. The explanation given by Dr. West of this fact is, that the uterus is central, and therefore more likely, when enlarged, to press on the neck of the bladder and produce retention, than an ovarian tumour, which is generally lateral. In such cases, difficult micturition has generally preceded the retention. *Retroflexion of the uterus*, or *retroversion* of this organ, when suddenly produced, may also cause retention, which either supervenes suddenly, or is not detected for a long time, in consequence of partial escape of the contents of the bladder occasionally taking place. In cases of *prolapsus of the uterus*, retention may occur during the catamenial periods, when the organ is larger and heavier, and in cases of prolapsus of the bladder itself, and chronic inversion of the uterus, etc.

Another form of retention, not by any means uncommonly observed, is that witnessed in *hysterical* patients, arising from spasmodic contraction of the sphincter, associated, perhaps, in some cases, with an erectile condition of the clitoris. Retention from this cause is accompanied with a good deal of acute pain in the hypogastrium. The attack is of a more acute character than in the cases before considered. There is generally a history of previous attacks of a similar character; and the previous history of the case affords, on investigation, pretty complete evidence of the presence of hysteria in some one of its multitudinous forms. In many cases, the nature and cause of the presumed retention cannot be made out without an examination.

Lastly, there are cases in which no urine is passed, because there is none in the bladder. This kind of *suppression* has been known to be produced by pressure of large ovarian or other tumours on the ureters. More ordinarily, however, suppression in the true sense of the word is due to other causes, the consideration of which does not come within the scope of the present inquiry.

TREATMENT.

There is comparatively little to be said respecting the treatment of disorders of micturition; the treatment usually consists in removing the cause of the disorder, whatever that may be. Respecting the majority of cases of this kind, therefore, the question of treatment will be more appropriately considered elsewhere. In many instances, the making of

the diagnosis is the essential thing, the mode of giving relief naturally suggesting itself when the cause of the pain or inconvenience has been once ascertained. There are a certain number of cases, however, in which palliative treatment may be indispensable; the removal of the disorder there and then present being necessary. Such are cases of *retention* of urine in which the patient is absolutely unable to evacuate the bladder, and requires temporary relief.

The catheter is the instrument by means of which the relief in question is obtained; and it may be well in this place to give a few directions as to the manner in which this instrument is best applied.

Mode of Introducing the Female Catheter. The first thing which it is important to be aware of is this—that ease in the use of the instrument is only to be attained by practice; and, consequently, that it is important that opportunities for practice in the manoeuvres necessary be not, as is often the case, neglected. The plan to be adopted is the following:—The patient is to be laid on her back; the operator is to stand on her right side; the right leg is to be flexed, the sole resting on the bed or couch. The operator then, by means of one finger of the left hand carried from the abdomen over the pubis, ascertains the position of the clitoris, and of the urethral orifice just beneath it; and, having done this, the right hand, holding the gum-elastic or silver catheter, is passed under the right leg, and the point of the instrument guided into the urethral canal. The introduction of the instrument is very easy, if these several points be attended to, and by one who is cognisant of the anatomical relations of the parts. The principal thing is to make certain, in the first instance, of the position of the clitoris and urethral orifice; the latter is known by the fact that the vaginal canal is immediately below it. If the finger be introduced into the vagina, the urethral canal must therefore be in the median line immediately above it.

It is convenient to have a slender India-rubber tube, five feet long, attached to the catheter. The urine then flows directly into the receptacle on the floor.

THE CHLORATE OF POTASH TREATMENT OF PHTHISIS. "We see, then," says Dr. Percy, of New York, "that Dr. Fountain's high expectations of the great value of chlorate of potash in phthisis, are not likely to be realized; for others who have preceded him in the same views, have experienced but little benefit from it, and those who have followed up his experiments have met with but little success. It is a strange fact, but we daily learn over again the same lessons, and many of us let them slip as quickly from our memories. Theories, and systems, and medical agents are brought forward, and have their day, and are for awhile the fashion; but when common sense reviews these claims, they soon assume their proper level. Time passes by, and another mind, reasoning from the same deductions, takes up the same remedy with the same earnestness, and again it becomes the fashion, and is again forgotten except by those who use it for its legitimate purposes. The thing occurs again and again, and the last man who vaunts it, through his imperfect investigation of medical literature, is equally honest with the first, and supposes he has made a most valuable discovery. Dr. Fountain has shown no more Quixotism in mounting his hobby, than Dr. Churchill has shown in riding his; but let us take a lesson from their experience." (*American Med. Times.*)