healthy, but the os was obliterated. In Dr. Coley's case, the patient died of peritonitis and purpura; and three pints of dark coloured, thin, and offensive fluid were found. The uterus was thinned to the consistence of a bullock's bladder. The os was closed by a large tuberculous mass about the cervix, and a plug of mucus. The entire lining membrane was softened, in many places ulcerated.

The treatment in these last three diseases is the same. The os must be opened to give free exit to the uterine contents. The opening may be made by a cannula; but some preliminary precautions should be taken. In one case I treated, where occlusion of the os was due to a severe instrumental labour, and in which, moreover, the uterus was retroverted, I pulled down the uterus by a vulsellum, so as to get the cervix and fundus in the same axis; and, using a very fine trocar, penetrated to about a line or two every day, keeping open the passage made towards the uterine cavity by a plug of wood covered with cotton wool, and secured in its place by a large plug of cotton behind; but in this case the adhesions extended about half an inch up the cervix. When once the cavity is entered, dilatation by a sponge-tent or the hysterotome is an easy matter. It is wise in these cases to keep the new opening made patent by a plug, changed daily, till the proper menstrual epoch occurs, when all danger of reclosure ceases.

[To be continued.]

Illustrations

OF

HOSPITAL PRACTICE:

METROPOLITAN AND PROVINCIAL.

ST. MARY'S HOSPITAL.

FATAL CASE OF GROUP IN WHICH TRACHEOTOMY WAS PERFORMED.

By EDWARD PARKER YOUNG, Esq., House Surgeon to the Hospital.

—, aged 4½, was sent to the hospital by Dr. Handfield Jones, on Feb. 16th, 1862, to have tracheotomy performed, as the symptoms were very urgent, and as tracheotomy appeared the only chance of saving the child from death.

It appeared that the child had been unwell for three or four days, and had been gradually getting worse up to the time of its admission; for when it was first admitted the face was livid, the head thrown back by every inspiration, the pulse quick but full, the alæ of the nostrils were expanded; and the child presented all the symptoms of impending suffocation. Tracheotomy was, therefore, performed; and it was with great difficulty that the tube was inserted, on account of the enormous venous hæmorrhage. When the trachea was opened, blood was drawn in, which was sucked out by the mouth of the operator, and nearly an hour and a half elapsed before good breathing was established through the tube. After this time, the breathing became very good, and the child took nourishment, and slept at intervals, till the next day, twenty hours after the operation; when alarming symptoms came on, the breathing became more laborious, the respirations quickened; the child appeared very restless, and vomited frequently; and expired twenty-eight hours after the operation.

A *post mortem* examination showed that the lungs were collapsed and pneumonic, and contained very little air. The trachea was lined by a false membrane, which was loose, and extended into the bronchi and their ramifications. As this state of parts explained the cause of death, the examination was not proceeded with further.

The remedies in this case were emetics, salines, and expectorants; but the *post mortem* examination clearly showed that it was impossible that the child could have survived with the respiratory apparatus in such a disorganised state.

WESTMINSTER HOSPITAL.

MEDIAN LITHOTOMY ON A BOY: RECOVERY.

Under the care of B. HOLT, Esq.

MEDIAN lithotomy has been practised occasionally during the last five years at many of the metropolitan hospitals; but the operation has been confined for the most part to adult patients. Mr. Holt has, however, performed it on several occasions upon children, with the best results. The case here recorded is interesting from the fact that, finding some difficulty in withdrawing the stone (which was of considerable size), Mr. Holt made a small incision in the side of the prostate, which allowed of the immediate withdrawal of the calculus, and was productive of no injury to the patient; and it serves to illustrate the fact that it is tearing rather than cutting which gives rise to inflammation and abscess about the prostate, as was so strongly insisted upon by Mr. H. Thompson in his Lettsomian lectures.

In this case, as in all Mr. Holt's other cases, the urine did not pass involuntarily through the wound, as it usually does after the lateral operation; but the patient was able to hold it for some hours, although in the act of micturating the urine came through the wound for the first few days.

John Stenson, aged 16, but small for his age, and of a pale, flabby appearance, although he has always lived in the country, was admitted into the Westminster Hospital, under Mr. Holt's care, with stone in the bladder, on Jan. 24th, 1862. He had suffered from symptoms of stone since birth; and four years ago a surgeon sounded him, and recognised the presence of a calculus; but no operation was performed, because it was supposed that the boy was labouring under kidney-disease.

Mr. Holt, having detected a stone, proceeded to perform median lithotomy on the 11th February, 1862. After opening the membranous portion of the urethra, Mr. Holt always passes a curved steel probe with a handle along the staff into the bladder, and then introduces the forefinger between the probe and the staff, upon both of which traction is made. Thus dilatation is accomplished with ease and certainty, both instruments being withdrawn when the finger reaches the bladder.

The stone was readily seized with the forceps, and withdrawn for a short distance; but it was found impossible to draw it through the neck of the bladder; nor could the forceps be disentangled, so as to remove it with the scoop. Mr. Holt, therefore, introduced a straight probe-pointed bistoury, and divided the tissues slightly in a lateral direction, when the stone was immediately and readily extracted. The calculus was an inch and three-fourths in length, an inch and one-eighth in width, and seven-eighths of an inch in breadth, and weighed six drachms, two scruples, and six grains. Its composition was phosphatic externally; but upon section there was found a mulberry (oxalate of lime) calculus, of the size of a hazel-nut, imbedded in the centre of a mass of phosphates.

The boy had a good night, and was able to hold his urine; when he passed it, however, it all came through