leaving in a few days sound skin beneath; or whether it shall produce the very different effects above mentioned. It is scarcely possible to doubt that similar events occur in other organs, and on a larger scale, and that senile gangrene may originate in this way. The trivial amount of neuralgia in the robust, and its fearful intensity in the weakly, afford a highly significant contrast. It seems impossible to consider the circumstances just mentioned, and not to recognise the vast importance of the state of the tissues, as compared with that of the poison or exciting cause. We know it is just the same with one of our most serious diseases; in the same family, one member may have scarlatina most mildly and favourably, while another may be smitten deathfully, and perish in a few hours; yet both imbibed the same poison and at the same time.

The treatment of herpes in healthy strong subjects is a very small matter. A saline purge, and a lead lotion or ointment to the affected part, are all that may be requisite. But in the weakly, from any cause, the case may be very different.

I shall not soon forget the case of an elderly lady, aged 70, of very calm, well governed mind, and robust frame, who was attended by me through a severe attack of herpes. She had, before the eruption appeared, internal pain through the back and side affected, which disappeared in three or four days; but afterwards, while the sloughing sores left by the eruption were slowly healing, and for many weeks or months afterwards, she suffered most distressing neuralgic pain of the back and side and arms. So severe was it that the gentlest handling of the side was scarce endurable; and her health suffered so grievously that I was almost apprehensive for her life. This pain continued at its full severity when the sores had healed; it was not constant, was of burning character, aggravated at night. Quinine to cinchonism abated it, as well as the local application of chloroform and belladonna liniment. Change from London to Brighton improved her general state; but she never, I believe, quite lost the pain; certainly not for more than a year.

The treatment of herpes zoster which I find usually indicated is to give quinine and iron, or bark and ammonia, to anticipate the depression and neuralgia, and to dry up the eruption speedily with lead lotion or lead and zinc ointment. Cazenave recommends in refractory cases of continuing pain a blister to the painful part. Von Bärensprung applies his blister in the neighbourhood of the spine, but with the endermic use of morphia. I think the subcutaneous injection of narcotics would be preferable.

With regard to herpes generally, I think the conclusions must be: 1. That there is nothing whatever of eliminatory effort in it; 2. That it has a distinct relation to nerve-disorder.

There is a form of eruption with which I am familiar, which corresponds precisely with none of the above. It consists of scattered sero-purulent vesicles, larger than those of herpes, more prominent and superficial than those of impetigo, developed on a slightly inflamed area. It chiefly occurs in cachectic children; and coexists with a tendency to whitlow. Arsenic, iron, and cod-liver oil, and the like, are its efficient remedies. Its chief interest consists in its furnishing one more instance of the great tendency in cachectic states to derangements of nutrition and quasi-inflammatory effusions which cease under a tonic regimen. I subjoin the following illustrative case:—

CASE XI. J. P., aged 4½, male, was admitted Dec. 23. He had been ill fourteen days with cachectic vesicular eruption on his head and all about him; even a scratch festered. He had been weak ever since a fever last summer.

The forms of eruption now considered evidently constitute, in some measure a natural group: 1. From their

superficial character; 2. From their pouring out more or less copious sero-purulent exudation; 3. From their being (except herpes) so markedly controlled by arsenic.

[To be continued.]

SOUTH DEVON AND EAST CORNWALL HOSPITAL.

LOOSE CARTILAGES IN THE KNEE-JOINT.

By W. J. SQUARE, Esq., Surgeon to the Hospital.*

SINCE I wrote the above case of loose cartilage in the knee-joint, my attention has been directed to an article in Mr. Chatto's Report on Surgery, in the *British and Foreign Medico-Chirurgical Review* for January 1862, entitled "On the Extraction of Articular Foreign Bodies from the Knee-joint, by M. H. Larrey (*Gazette des Hôpitaux*, No. 67)."

This article has such an especial and valuable relation to the case which I have reported, that I transcribe it, and shall venture to make two or three remarks on its bearings upon the mode of operation which I adopt.

"As the result of his bibliographical researches, M. Larrey has collected references to 167 published cases in which operations have been performed on the knee-joint for the extraction of articular foreign bodies. In 129 of these cases, the extraction was performed by the old or direct method, and in 38 by the indirect or subcutaneous method. The results are given for 121 of the former cases, being successful in 98, doubtful in 5, and fatal in 28; while of the 38 indirect operations, 19 were successful, 15 failed, and 5 were fatal. From these facts, and from others communicated to him orally by his colleagues, the author deduces the following conclusions.

"1. The operation of extracting these foreign bodies is a serious one when practised by direct incision, and a difficult one by the subcutaneous method. Many accidents have occurred even amongst the somewhat numerous cases in which recovery has taken place.

"2. Extraction is proportionally more dangerous than is the affection itself; that is, more redoubtable accidents may be induced by the operation than by the presence of the foreign body.

"3. While many cases of recovery may be verified, many instances of failure or of fatal results have doubtless never been published.

"4. The operation definitively would seem to be indicated only under the following circumstances: the certain and complete mobility of the foreign body within the joint; the persistence of the accidents caused by its presence—viz., pain, arthritis or hyarthrosis, and lameness; the failure of compression and other means for fixing the condition of the body; and the free consent of the patient, although made aware of the dangers and difficulties of the operation."

The statistics of M. H. Larrey distinctly maintain the truth of his conclusion that the operation of extracting foreign bodies from the knee-joint "is a serious one when practised by direct incision"; but I cannot agree with him in his conclusion that it is a "difficult one by the subcutaneous method". If the surgeon perform it exactly after the fashion which I have described, and use a long, narrow, strong tenotomy-knife, he will find that, in the great majority of cases, he will be able to execute it with much facility. M. Larrey states that "extraction is proportionally more dangerous than is the affection itself". In this conclusion I cannot concur; and for this obvious reason, that I have myself operated or assisted at operations by the subcutaneous method fifteen times without witnessing any evidence of

* These remarks were intended for addition to Mr. Square's paper in last week's JOURNAL, but arrived too late for insertion.

the least inflammatory or other morbid action. This fact also makes me object to the fourth conclusion, that "the operation definitively would seem to be indicated only under the following circumstances: the certain and complete mobility of the foreign body within the joint; the persistence of the accidents caused by its presence—viz., pain, arthritis or hyarthrosis, and lameness; the failure of compression and other means for fixing the condition of the body; and the free consent of the patient."

It is true that it is necessary to be certain of the complete mobility of the foreign body within the joint. Many years ago I saw an operation attempted on a case of the description to which M. Larrey alludes, and death was the result. Again, no discreet surgeon would operate unless the symptoms incident to the presence of a foreign body in the knee-joint were frequent or permanent. Still I must express my decided opinion that in every case, unless very exceptional, the foreign body should be removed by the form of subcutaneous incision which I advocate.

Original Communications.

MEDICAL PSYCHOLOGY.

By ROBERT DUNN, F.R.C.S.E.

III. On the Psychological Phenomena or Symptoms of Disease.

[Continued from page 522.]

Loss of Speech or the Power of Utterance. Among the abnormal phenomena of the intellectual consciousness, we may now turn from the noisy ravings of delirium, in febrile and other diseases, to the consideration of an opposite state, depending upon other and varying pathological conditions—that of dumb silence; not, however, of insensibility and coma, in which the phenomena of life are reduced to a mere series of automatic movements; but to that which results from the loss of articulate speech—a lack of the power to give utterance to the thoughts, feelings, and emotions of the mind. Now, the loss of speech or power of utterance may arise from a variety of causes, and may be quite independent of any cerebral lesion, or implication of the intellectual consciousness. For, aphonia is not unfrequently the consequence of disease of the larynx or vocal organs; at other times, the loss of voice is occasioned by pressure upon, or change of structure in, the lingual and glosso-pharyngeal nerves; either at their origin or in their course. From the close and intimate connection between the corpora olivaria and the nuclei of the hypoglossal nerves, Schröder van der Kolk has been led to infer that the former are auxiliary ganglia to the latter, in the production of certain combinations of movements in the acts of articulation and speech. "For," says he, "speech and the articulation of words require such a multitude of peculiar motions of the tongue, and such an infinite number of varying combinations of its muscular movements, that it cannot appear strange that two auxiliary ganglia should be required for the performance of these functions." After stating his conviction that all motor nerves take their origin from certain groups of multipolar ganglionic cells, he goes on to observe, "that he is acquainted with no other part of the extra spinal cord or brain where the multipolar ganglionic cells are so densely accumulated as in the nuclei of the hypoglossal nerves"; and then adduces pathological evidence in support of his opinion that the corpora olivaria are the organs for the articulation of the voice. But he by no means considers the corpora olivaria as the organs through which the will directly

acts; for these he admits to be, chiefly, the corpora striata, any injury to which is invariably followed by paralysis; and he adds that, "when the organs of the will are disturbed, the paralysis of the tongue and of the neighbouring parts renders the utterance of words wholly impossible."

During the past year, an interesting case came under my notice. The patient was an intelligent man, a law-stationer, of temperate, regular, but sedentary habits, aged 61 years when he died in September last. In April 1858, he had an apoplectic seizure, which left him, for some time, hemiplegic on the right side; but from this he gradually, and to all appearance completely, recovered; so that, in the course of five or six months, he was again actively engaged in his business. In May 1859, he had a second attack; but this left permanent effects behind it. He never properly recovered the use of his arm or leg. He could sign his name, and get about the house with assistance. He experienced some difficulty in deglutition, and there was an impediment in his speech. A third seizure in June last year left him utterly helpless as to his arm and leg, unable to articulate a word distinctly, or so as to be understood, and only able to swallow liquids, and these with difficulty, threatening, at times, suffocation. Still to the last he was sensible, and easily moved to tears, and suffered from emotional excitement. No *post mortem* examination was permitted, which I much regretted. As there had been three distinct seizures, it would have been interesting to have noted the records which they had left behind them; and to have seen how far the corpora olivaria were implicated.

In another case, a young lady, of a delicate constitution, who had suffered from repeated rheumatic attacks, and who had, in consequence, disease of the mitral valves, under a severe attack of bronchitis, became suddenly hemiplegic and speechless. She soon recovered consciousness and sensibility, so that she perfectly understood whatever was said to her, and noticed all that was going on about her; but she was hemiplegic and without power of utterance. A few days afterwards, and at the time quite unexpectedly and suddenly, she expired.

Here, too, no examination was permitted. But, from the suddenness of the death, and of the loss of speech, how interesting and instructive it would have been to have seen the actual condition of the cerebral arteries, and the *locale* of the rupture, in reference to the corpora olivaria and striata; and to the observations and researches of Dr. Kirkes. How often, in private practice, I cannot help remarking, does it happen that we are denied the satisfaction of a *post mortem* examination in cases where we have had the opportunity of carefully watching the patients during life, and of noting every appreciable change in their symptoms, so that they have, in consequence, become invested with an interest in our minds, which nothing short of a knowledge of the pathological conditions can fully satisfy. In private practice, we have neither the appliances nor opportunities for pathological investigation, which the physicians and surgeons of our hospitals can command and do possess; but still it is gratifying, and it ought to be encouraging, to reflect what may be done, and what an Abercrombie actually did accomplish single-handed in the walks of private practice.

Now, in all cases of loss of speech or the power of utterance, which are of cerebral origin, there is involved functional derangement or structural change in the nervous apparatus of the intellectual consciousness; for "perfect speech—that is, the power of expressing our thoughts in suitable language—depends upon the due relation between the centres of volition and of intellectual action." (Dr. Todd.) I had a striking illustration of this, in the young woman's case to which I have had occasion so frequently to refer; for she continued