careful subsequent treatment, less certain than the more tardy dilatation process. Care should always be exercised in dilating the cervix uteri; for although, as a rule, the organ is capable of bearing a tolerable amount of rough treatment with impunity, it is not by any means always so; fatal peritonitis has followed want of attention to this rule. If sponge-tents be employed, very great attention to cleanliness is essential. It is proper always to allow a day or two to intervene between each insertion of the tent, and it is better not to leave the tent in the cervical cavity more than four or six hours. After removal of the tent, tepid water should be very gently thrown up the vagina.

Illustrations OF HOSPITAL PRACTICE: METROPOLITAN AND PROVINCIAL.

ST. MARY'S HOSPITAL.

PRACTICAL REMARKS ON THE PATHOLOGY AND TREATMENT
OF SKIN-DISEASES.

By C. HANDFIELD JONES, M.B., F.R.S., Physician to the
Hospital.

[Continued from page 647.]

Erythematous Eruptions. The next group of affections I propose to examine are those which are characterised essentially by hyperæmic afflux to the skin—in fact, inflammation just stopping short of effusion, or attended with very slight. Roseola and erythema are the typical members of this group; but it may include also the slighter forms of erysipelas.

Commencing with the last, we must notice the justice of Mr. E. Wilson's remark that erysipelas serves to establish a link of transition between eruptive fevers and cutaneous exanthemata without constitutional affections of a specific kind. The graver forms resemble the eruptive fevers in their transmissibility by infection and contagion, and their being attended with fever often of a very low type; while the slighter approach very much to the common skin-eruptions in the great obscurity of their causes, their great liability to recur and to become chronic, the absence of fever, and the partial affection of the skin. Now, whoever reads over the lists given by authors of the causes of erysipelas (*Vide Copland's Dictionary*, art. Erysipelas, p. 826) must, I think, admit that the disease cannot be supposed to depend, like small-pox, on the introduction of a definite poison into the circulation. Evidently a certain state of system, a peculiar predisposition, goes for much more. With this, "the most trifling irritation" will cause the disease, which without it has no such effect. Foul atmospheres and depressing causes are potent in generating this predisposition. Erysipelas may be of the most sthenic or æsthenic type, and require the most opposite treatment. In the one, free bloodletting and purgation may be necessary, and little else. In the other, quinine, iron, and stimulants must be poured in lavishly. In London, at present, a sthenic case would be a *rara avis*; but I have heard of very marked ones in the country, on the best authority.

The general bearing of the above facts certainly tends to show that the state of the vital powers is the main thing to look to in endeavouring to appreciate the phenomena of erysipelas. The conception of a poison seems subordinate to that of a predisposition. It is the

latter that we have to deal with practically. As therapeutists, we renounce all idea of eliminating the poison, if we suppose there is any, seeing that in the worst cases all our exertions are directed to supporting the strength, and arresting the spread or obviating the effects of the inflammation. The phenomena in the æsthenic form have much resemblance with those of vaso motor nerve paralysis, and are modified very beneficially by a stimulant and tonic treatment calculated to raise and sustain nervous power. In a case of frequently recurring erysipelatoid inflammation of the face in a feeble old lady under my care, the only measures that do good are fresh air, good living, and citrate of quinine and iron. Certainly, at any rate, in the milder forms of erysipelas, we have little reason to believe in the existence of any definite poison, and still less to think that we have any power to eliminate it by treatment.

Roseola and Erythema are two minor skin affections, which it requires some artistic power to separate, and which, for all useful purposes, need scarcely be separated at all. They are mere hyperæmiæ of the skin, often of no great extent, and for the most part of transitory nature. Erythema intertrigo and erythema paratrimma are cutaneous congestions occurring usually in young children under slight irritation of the surface, aided by heat and moisture. Roseola infantilis is more dependent on general causes, which probably induce some amount of vaso-motor paralysis. I have seen this affection very marked in a child affected with malarious disorder. Roseola cholericæ is no doubt produced in the same way. Roseola æstiva and roseola autumnalis are in all probability the mere results of heat, or atonic relaxation of the vessels. They are just such transitory congestions of the outer integument as often befall in the same season the inner, and give rise to diarrhœa. Roseola æstiva is a very little step from eczema solare. Roseola variolosa, roseola vaccina, roseola rheumatica and arthritica, are minor results of the direct irritation of the poisons whose name they bear. Erythema nodosum is interesting as an example of a local inflammation invariably, according to my experience, of æsthenic character, and yielding readily to tonics. It is a like case to the cure of eczema by arsenic. The varieties of erythema are mostly associated with some catarrhal or febrile disorder, bronchitis, enteritis, etc.; and may be regarded as cutaneous irritation produced by the same cause that gives rise to the internal. Erythema fugax, by virtue of its fugaciousness, must be ranked as a neurosis; it is without example that any true tissue-inflammation should appear and disappear in an apparently causeless manner. This is the sole appanage of nerve-inflammations. Erythema papulatum and erythema tuberculatum are evidently related to urticaria, to lichen urticatus, and more remotely to lichen. The marginate and annulate forms of erythema and roseola are simply examples of that curious tendency in many skin-eruptions to spread excentrically, while they heal more or less completely in the middle. It is not at all clear on what this depends, nor why it is so often absent. The mode of distribution of the nerves does not seem in any way to explain it. So far is certain, that it is not a peculiarity belonging to any single form of eruption, and therefore probably depends on something in the skin itself. Erythema læve seems to me to have no claim to be considered as an eruption; it is for the most part a mere co-result of the condition that gives rise to anasarca.

Roseola and erythema are well described by the synonym given of the latter by Mr. E. Wilson; they are truly inflammatory blushes, and, like all blushes, are very often dependent on a nerve-paresis giving rise to arterial dilatation. In some cases, as in malarious and choleric roseola, this paresis directly affects the vaso-motor nerves of the part itself. The same doubtless

occurs in some of the erythemata, which are associated with rheumatism, bronchial or gastric catarrh. In others, the paresis may be the result of inhibitory action set up in some disordered part. This is very apparent in some persons of habitually weak digestion, who have considerable flushing of the face and neck immediately after a meal. The indication here is clearly to lessen the irritability of the mucous membrane, while in the former case it is to impart tone. In roseola variolosa and some other forms, the tissue may be directly affected. The general resemblance of these affections to erysipelas in its milder forms is very considerable; and the pathology of all is, I believe, very much the same. They depend upon no specific and constant cause, but may be set up by many. In some instances, a material poison in the blood may excite the phenomena; in others, some subtle atmospheric influence acting upon the nerves; in others, again, a remote irritation acting through nerves on an excitable skin. Sometimes, as in erythema intertrigo, the causes are purely local, and the treatment is clear. I am quite satisfied by experience of the value of tonics in some erythemata, and of astringent sedatives in others. The following cases are of some interest.

CASE I. Mrs. D., aged about 40, was seen on January 2nd. She had recovered some weeks ago from ague. She had been now for some days feeling very depressed, in spite of stimulants. She felt bilious; her loins ached; and she had an eruption on the face near the eyes, consisting of slightly injected patches, with some swelling. On the left of the neck, there was a circular spot the size of a shilling. She was ordered to take a draught of carbonate of ammonia, tincture of cinchona, arsenical solution, and infusion of cascarrilla, three times a day.

Feb. 4th. Fever came on on the 2nd, and continued till this morning. Pulse excited, 100; tongue clean; skin moist; urine very pale and clear. She had no appetite. The eruption on the face formed two symmetrical red patches, with subcutaneous oedema extending some way under the orbit. The circular spot on the neck had increased in size, and some small ones had appeared on the hands and fingers. The mixture was continued; and she was ordered to take a pill of mercury with aloes at night, and to use lead lotion (two parts to ten).

Feb. 6th. The eruption on the face was much less; on neck, was gone; there were traces still on the fingers. The medicine was continued.

Feb. 9th. She was much better. Her appetite was good; she was cheerful. Slight staining of the skin and desquamation were all that remained of the face-eruption; the spots on the fingers were now reduced to rings. Two grains and a half of quinine were ordered to be taken three times a day. A few days later, she was quite convalescent.

CASE II. L. B., aged 16, was admitted December 15th. She suffered with some eczematous eruption, and with great pain in the forehead, which came on regularly every morning, and lasted about an hour; it was attended with giddiness. The catamenia were regular, but very dark. The eczema yielded speedily to bark and arsenic; but on Dec. 20th it was noted that she looked wonderfully well—in fact, unnaturally so; the cheeks and forehead being over red and hot. She said this was from a flush which came on every afternoon. With strychnia and cinchonine, she improved very much. I have recorded a similar case in the *Lancet* for July 28th, 1855.

CASE III. E. J., aged 18, was admitted February 10th, having been ill three years. She had pain after eating at the chest; meat lay very heavy. There was no epigastric tenderness. The functions were all natural. After taking food, her face was apt to get hot and red, and to break out into a pimply rash; and her chest the same. She complained of lumbar pain. With nitrate of silver and lactic acid, the digestion improved, and the

eruption diminished a good deal. She only attended fourteen days.

CASE IV. M. B., aged 21, was admitted May 20th. She had been ill one year, on and off. She had pain and tenderness at the epigastrium, and pain in the sacral region. Her legs swelled towards night. The face was swollen and red, as from superficial erysipelas; a week later, it was desquamating. Her head ached. She felt very weak. There was much pain after food. Under treatment directed to soothe and tone the stomach, with a lead lotion to the face, her digestion improved, and her face became much paler.

CASE V. M. A. A., aged 14 months, was admitted April 21. She had been ill one month with an eruption consisting of slightly raised patches of diffused inflammation, on the arms, back, abdomen, and legs. Two or three small bullæ had formed on the wrist of the left arm. The eruption had been on the neck. She had just cut two canines, and was still suckled. She was a fat, puffy child, fretful. The skin was hot and dry. She was ordered to take four times a day a teaspoonful of a mixture containing two ounces of citrate of potash mixture and twelve minims of tincture of opium; and to have lead lotion applied.

April 24th. There was an aphthous patch on the tongue; both the hands were intensely oedematous; the skin was shining and whitish, evidently from subcutaneous effusion; the bowels were open. She was ordered to take two drachms of the following mixture three times a day.

℞ Misturæ potassæ citratis ℥ij; ferri et quinae citratis ℥j; liquoris potass. arsenitis mxxv; tincturæ hyoscyami m℥.

The lotion was repeated.

April 28th. She was much better. On the 26th, between 5 and 6 p.m., a large red patch appeared on and about the left knee, and was now of large size. The hands were well. She coughed at night. The mixture was continued, water being substituted for the citrate of potash solution; and the lotion was repeated.

May 12th. The eruption had affected the legs down to the feet; it had now quite disappeared everywhere, leaving only some desquamation. Bowels costive. Her health was much improved. The mother stated that the eruption, even when it swelled the hands so tensely, commenced as redness, which faded as the swelling took place. The mixture was ordered to be given twice a day, and castor oil to be taken.

REMARKS. Cases I, II, and V illustrate the advantage of the tonic treatment of some erythemata; Cases III and IV, that of allaying a remote irritation. Case V was one of great interest. There was so much feverishness and irritation that I forbore at first from the employment of tonics; and, when I began to use them, combined them with a saline. Yet they were signally useful, improving the health and abolishing the eruption. The existence of vesicles and of subcutaneous effusion shows how slight may be the essential difference between phenomena apparently very dissimilar. In ninety-nine children out of a hundred, under similar circumstances, the eruption would have taken the form of eczema. Some slight peculiarity in the vital condition of the skin determined it to be an erythematous oedema, yet with sufficient surface-effusion to show how readily it might have assumed the other form. I have recorded a similar, but more chronic case, in an adult. (See *BRITISH MEDICAL JOURNAL*, Dec. 19th, 1857.)

Purpura, though not purely a cutaneous affection, yet, in its less grave forms, is so much restricted to this tissue, and is of so much interest in various respects, that I must take some notice of it here. It is certainly not so much an exanthem as an hæmorrhage; but still the more sthenic cases are attended with more or less swelling and congestion, apart from the sanguineous effusion. By the former it shows its affinity with the

erythematæ. Rayer gives one case, indeed, of these in which sanguineous extravasation occurred, and which was consequently almost identical with purpura. It seems quite established that it does not depend on any uniform cause, but rather on a peculiar state of system, of which "debility of the nervous powers" (Eras. Wilson) and fragility of the capillaries must be considered as the chief characteristics. In one form, there is decided fever, and what is called high arterial action, which of course means cardiac excitement with relaxation of the arteries—a condition which paresis of the vagi and vaso-motor nerves would certainly induce. In the other, there is an absence of pyrexia; and the spots appear without tumefaction or congestion, or with but little. Treatment in many instances by tonics or astringents is very effectual (the severest case I ever saw was cured by matico); but in some cholagogue purgatives are essential. In these, the symptoms are either the result of morbid matter in the blood, or of accumulations in the liver affecting its nerves depressingly. In the other, the nervous and vascular disorder is primary. This is no unfrequent occurrence in pathology, that the same or very similar results are produced by very different causes. I have seen more than once subconjunctival effusion of blood as the result of malarious disorder; and mental depression and nostalgia alone have produced the allied disorder, scurvy.

CASE I. J. M., aged 12, was admitted April 5th, having been ill one week. He had numerous and largish livid spots of purpura on the legs, mostly on the right. He did not feel ill. Skin cool; pulse weak. He had not been without potatoes. The tongue was a little coated. He was ordered to take seven grains of citrate of quinine and iron, and seven grains of citric acid in an ounce of water three times a day. In fourteen days the eruption had quite disappeared, but he required tonics some time longer.

CASE II. J. W., aged 42, was admitted August 3rd. He had been ill four days, but had gonorrhœa a fortnight previously, for which he had taken copaiba. There was now a copious eruption of urticaria on all parts of the body. The right forearm was much swollen; the left was covered with red spots, which faded on pressure; but the thighs were thickly set with red livid patches like purpura. Pulse 87, of tolerable force; tongue rather white in the middle; bowels open. He attributed his disorder to getting wet when very hot, and drinking cold beer. He was ordered to have a scruple of ipecacuanha immediately, four grains of calomel at bedtime, and an ounce of white mixture three times a day.

Aug. 5th. There was much itching still; the patches on the legs were fading; the arm was less swollen; no fresh spots had appeared. He was relieved by the emetic. The bowels were not much purged. He was ordered to take two grains of calomel every alternate night, and a dose of white mixture on the following mornings; and to have six ounces of lime-juice daily. He was discharged well on the 24th.

[To be continued.]

ST. GEORGE'S HOSPITAL.

SERO-CYSTIC TUMOUR IN THE NEIGHBOURHOOD OF THE ANKLE-JOINT: REMOVAL.

Under the care of PRESCOTT G. HEWETT, Esq.

JOSEPH W., aged 50, a labourer, was admitted into Winchester Ward, under Mr. Hewett, on May 21st, 1862. He stated that about eight months ago, he noticed a small lump on the left ankle. He could assign no cause for it; he had never worn tight boots, or been subjected to pressure of any kind in this part. The tumour had gone on steadily increasing since its first appearance.

On admission, there was found to be a small tumour of about the size of a walnut over the inner malleolus;

it was circumscribed, firm and hard, but not osseous; it was quite moveable under the skin, and in no way connected with the bone. It was not painful; but caused great inconvenience in walking.

May 29th. Mr. Hewett made a longitudinal incision over the tumour, and dissected it out. It was adherent to the periosteum in one part; but it was carefully removed without injury to this membrane.

On examining the tumour, it was found to be composed of a number of small cysts collected together, and freely communicating with each other. The cysts contained a thick cheesy matter, and had little bodies growing within them, being evidently of the tubero-cystic variety. The man made a quick recovery; and left the hospital on June 14th.

Original Communications.

REMARKS ON LARYNGOSCOPY.

By EDWARD H. SIEVEKING, M.D., Physician to St. Mary's Hospital.

At a time when laryngoscopy is in its infancy among ourselves, it may be useful to multiply evidence of its practical utility; as the *cui bono* objection is already raised, and may diminish the confidence of the student of medicine in a method of examination, which, to those who have familiarised themselves with it, offers the most palpable evidence of its being a manifest advance in the diagnosis and treatment of laryngeal disease.

Having recently enjoyed the advantage of employing the laryngoscope under Professor Czermak's kind superintendence, I invited two patients suffering under hoarseness of doubtful pathology to submit to an examination by the Professor. The one offering symptoms of subclavian aneurism, appeared, both to Professor Czermak and myself, to be so peculiarly illustrative of the value of this new method of diagnosis, that I venture to submit it to the readers of this JOURNAL.

The following are briefly the leading features of the case:—

No. 39, June 21st, 1862. S. J., aged 40, of a spare habit, the father of a numerous family, has, for sixteen or seventeen years, been subject to frequent attacks of sore throat, but unaccompanied by hoarseness. In other respects, he has enjoyed good health till the commencement of his present illness, about a year and nine months ago. The first symptom was an attack of rigors, attributed to a damp bed used on a journey, followed, three days after, by hoarseness, which had continued ever since, with some dyspnoea; the latter feature has diminished during the last six weeks. He has had no pain in the throat; but has all along had a sensation of there being something in his throat. Ever since the commencement of the hoarseness, he has had a husky cough. For two months past, he has had a pain at the upper right thorax. The cough has not been worse since this pain has come on. He has never spat blood; but of a morning there is copious yellow expectoration. There is no dyspnoea at present; and position makes no difference in regard to respiration. There is a circumscribed dulness at the middle of the right clavicle, with a well marked systolic murmur, not heard over the heart, and extending feebly into the axillary and carotid arteries. The pulse of the right radial is feebler than that of the left.

On subjecting the patient to laryngoscopic examination, which he bore remarkably well, it became at once apparent that the hoarseness was due to the projection into the laryngeal cavity, from the anterior junction of