Illustrations

or

HOSPITAL PRACTICE:

METROPOLITAN AND PROVINCIAL.

ST. MARY'S HOSPITAL.

PRACTICAL REMARKS ON THE PATHOLOGY AND TREATMENT
OF SKIN-DISEASES.By C. HANDFIELD JONES, M.B., F.R.S., Physician to the
Hospital.

[Continued from page 625.]

Lepra and Psoriasis (continued). I will now refer to and relate a few illustrative cases.

Mr. Hunt's first case of lepra demonstrates satisfactorily the value of bloodletting and leeching, aperients, and the avoidance of all stimulants. His third is interesting, as showing how small a dose may sometimes be requisite. His second case of psoriasis shows the importance of a modified treatment, vegetable diet, and arsenic, trimming thus between sedation and stimulation. Mr. Green's first two cases of lepra and his two of psoriasis attest the value of bloodletting and aperients as preparatory to or assisting the operation of arsenic and the sulphur fume.

The following case shows forcibly the mischief which may be done by arsenic, if persisted with, in cases that are intolerant of it:—

CASE I. Miss —, aged 18; has suffered twelve years with psoriasis. No cause was known for its appearance. She had been lately under treatment by arsenic with purgation, which had rendered her stouter, but aggravated the eruption extremely. Her whole face, neck, most of her chest and arms, indeed most of her body, were covered; the skin was of a deep red, stiff and irritable, and cracked at the flexure of the left elbow, so that she could not extend the arm at all. She could not close her eyelids from the state of their skin. A copious desquamation was continually going on. Her general health was good. Wet packing was employed at first, with salines and wine of colchicum. Her sister informed me that the skin was quite pale for a short time after the daily morning pack. The sulphur-fume bath was used subsequently every night or alternate nights. She also took for some time an infusion of galium aparine.

In about three months, she had recovered from her former pitiable condition, with the exception of some scattered spots, which she had had before the arsenical course. The wet packing was applied by means of a mask to the face, and worn at night; it was of great service as long as the skin continued very irritable. The temporary pallor of the inflamed surface produced by the packing attests the sedative influence of moist warmth on the skin.

CASE II. W. D., aged 18, was admitted June 19th, with extensive psoriasis of eighteen months date. His health was good; the character of the eruption was non-irritative. Liquor potassæ (℥xx), chlorate of potash (gr. x), and bichloride of mercury (one-tenth of a grain), were each tried (being taken three times a day); but without good effect. Sulphur fume baths were also fully tried, but as fruitlessly.

On Sept. 9th, he began to take ten minims of liquor potassæ arseniatis, with three minims of tincture of opium in infusion of cascarrilla; and, as improvement was slow, the dose was gradually increased, till on Oct. 14th it reached twenty-five minims of arsenical solution and four minims of tincture of opium three times a day.

This was taken regularly, with the exception of one week's interval, till Dec. 23rd.

A week later, I saw him, and there were only remaining slight stains and congestion on the site of the patches. The medicine was then omitted; it had caused no apparent disorder the whole time, except, perhaps, one day. He took twenty minims of dilute sulphuric acid twice daily a month longer, and then ceased attendance.

How very different must the excitability of the skin (its vital property) have been in these two cases. In the one, a dose, I am nearly certain, not exceeding five minims three times a day, caused infinite deterioration; in the other, twenty-five minims three times a day caused as marked improvement. In neither were there injurious effects produced on the general health. The sulphur stimulus, decidedly beneficial in the first, was ineffectual in the last. Now, who can over-estimate the great and essential difference in these cases, on which the whole treatment turns? Is it not clear that, in a disease of this kind, there is question of much more than the state of the blood, which was presumably the same in both (according to the poison theory)? Is not the vital condition of the tissue practically the important matter?

CASE III. E. W., aged 13, was admitted June 7th, 1855, with very extensive psoriasis, guttata et diffusa, on her arms, legs, and thighs; some, but less, on the trunk and forehead. There was much itching. The eruption had lasted four or five months. Her health was fairly good, except some headache. She did not like animal food. She took for fourteen days one-twelfth of a grain of bichloride of mercury in decoction of elm-bark without benefit; and then was ordered a grain of hydrargyrum cum cretâ every night, and half a drachm of sulphate of magnesia in an ounce and a half of citrate of potash mixture thrice daily. She improved steadily, and was discharged well August 6th.

The next year she had a relapse, was admitted June 23rd, was treated by arsenic, and recovered pretty readily. Since then I have not seen her. It is clear that her pathological state was very different on these two occasions.

In the case of a girl, aged 12, who had suffered ten years with psoriasis, and who recovered ultimately under the use of arsenic with an issue in one arm, there was the unusual complication of chorea. According to the mother's account, the chorea had appeared on several occasions when the eruption had got better. The chorea was rather of the paralytic kind while she was under my observation; both it and the psoriasis got well together.

Does the fact that arsenic is a real remedy for chorea alone point at all to the concernment of the nervous system in psoriasis? Does it not in some measure countenance the opinion I have expressed as to the action of arsenic on the cutaneous nerves? In one of my cases, a weakly female, aged 24, rather menorrhagic, weakly, suffering with palpitation and considerable dyspepsia, arsenic was well borne by the stomach, and cured the eruption. In another female, also treated beneficially by arsenic, fresh spots of eruption always appeared at the catamenial periods. As this is a time when the general power is lowered, and all sorts of disorders are intensified, it is only what I should expect that psoriasis would be so too. The fact, however, is not very accordant with eliminatory views.

In some cases of asthenic character, citric acid has appeared useful. I have employed it mostly when, from some cause, arsenic was inexpedient. I have often remarked that, after the patches have all but returned to their healthy state, remaining only a little stained and congested, it is extremely difficult to get rid of this last trace of the disease. In fact, the capillaries seem to have become dilated, and lost their healthy tone; and

pro tanto there is actual organic alteration of the tissue. In one of my cases, well marked psoriasis was associated with well marked eczema. Both were ultimately cured by arsenic; but its administration had to be interrupted for about a month, in consequence of some asthenic ulcers which refused to heal while it was being taken, and soon got well afterwards. The ulcers would not form granulations as long as arsenic was given; was this owing to the contraction of the cutaneous arteries and diminished supply of blood? Sulphur ointment, ointment of iodide of sulphur, bismuth ointment, glycerine, bichloride of mercury lotion, have all appeared to me useful occasionally. Cazenave speaks favourably of the mercurial iodide.

Pityriasis. Pityriasis may be shortly described for our purpose as essentially a chronic somewhat modified psoriasis. The patches are more extensive, less congested, and covered with thinner and smaller furfureous scales. Mr. E. Wilson thinks it owes its origin to some unknown modification of innervation of the cutaneous textures; and is developed, mostly, in persons remarkable for the delicacy and susceptibility of their skin. Local irritants, he admits, may give rise to it on the face. As far as I can see, there is no important difference between this and the other squamous affections; their pathology is essentially the same; and, of course, the same principles must guide the treatment.

I quite coincide with Mr. Wilson in his appreciation of Rayer's remark, that "it is with general pityriasis, as with almost the whole of the chronic diseases of the skin that are independent of appreciable causes; a solid and enduring cure is only to be obtained by a general change of the constitution, brought about by dietetic means long and regularly pursued, effected naturally by the progress of years, and the modifications undergone by the organisation; or accidentally induced by some intervening disease, such as measles, scarlatina, etc. Here it is plainly laid down that no mere elimination is either possible or sufficient; but that an entire renovation, so to speak, of the nutritive action of the skin must be effected. I say of the skin, because there is often no disorder elsewhere. This is, of course, no easy matter; and treatment by drugs alone must often fail. In-veterate cases of squamous disease seem to me more analogous to superficial lupoid affections than to any other. They are, to a great extent tissue-affections, and much more enrooted in the vital actions than the exudative. These have more the character of a temporary disturbance from general causes; the other of a more permanent and local change. The following illustrative case is worth perusal:—

Miss —, aged 14, had had skin disease, more or less, since she was six weeks old. She had been often treated, but nothing had done good. She was very short, stout, and puffy looking. Her limbs and face had a plumpness which seemed to proceed rather from œdema than from fat, at least in part; her eyelids swelled at times. All her scalp, the back of her neck, and the upper part of her forehead, were more or less congested, rough, and covered with squamous formation. The hair was thin, but did not fall off. She was markedly pallid, and rather languid. The skin-disease was hereditary for three generations; the great-grandfather had it, but not the grandfather. The urine, examined five days later, deposited well marked oxalate of lime octohedra with vesical epithelium. It was not albuminous, moderately acid, of normal colour. She took at first ten minims of tincture of sesquichloride of iron three times a day; which was changed, after a month, for potassio-tartrate of iron in infusion of calumba. She used the sulphur fume bath steadily, and, besides, a spirit of rosemary wash and dilute citrine ointment. In less than four months, the skin had quite recovered, and her health was improved.

Ichthyosis. Of this disease, it is only necessary for

me to say that I quite coincide in the view taken by Dr. Waiburton Begbie, that white ichthyosis is a distinct affection from *ichthyose brune* and *ichthyose porc-épic*, the terms employed by M. Devergie. The former belongs certainly to the scaly diseases; the two latter to affections of the sebaceous glands. White ichthyosis is mostly congenital, affects the whole surface, is not attended with heat, itching, or inflammation; but with a dry, unperspiring state of skin. The general health is unaffected. Treatment is futile. In all these particulars, it seems to me that we have evidence of primary disorder of the nutritive action of the cell-bearing surface of the skin; but nothing to show that this is occasioned by any abnormal state of the blood. It is more like a malformation than a disease.

[To be continued.]

Original Communications.

INCARCERATED INTUSSUSCEPTION IN A CHILD SUCCESSFULLY TREATED BY INFLATION.

By EDWARD COUSINS, Esq., F.R.C.S.

[Read before the North London Medical Society, March 14th, 1862.]

FEBRUARY 23rd, 1862, A.M. A male child, aged 13 months, at 10 A.M., apparently in good health, after having had the breast, suddenly vomited all the contents of the stomach, including the undigested first meal of boiled bread, and immediately fell into deep syncope. Half an hour later, it vomited a thin serous fluid, and relapsed into a semicomatose state; surface generally pale, but rosy over scalp. The gums were lanced over two molar teeth which were approaching the surface. Castor oil was given, and at once vomited. After this, the child vomited immediately everything it took, and repeatedly vomited serous liquid when not recently fed; at last it refused food.

From one o'clock until the evening, it had several fits of tormina every ten or fifteen minutes. During these, its face was pallid, anxious, and distressed; it writhed, and pulled violently at its nose, ears, lips, and penis. The fits of tormina were not accompanied with tenesmus; they lasted two or three minutes, and were succeeded by semicoma, during which the eyelids met imperfectly, and the child was vexed if disturbed. Sinapisms were applied. Enemata were retained for a time, and then escaped without fecal matter.

5 o'clock. The abdomen was flaccid, and not oversensitive to pressure. The tormina were not accompanied with tenesmus or dejections. Examination *per anum* discovered nothing; but the finger, on being removed from the anus, was followed by some slightly bloody mucus. There were no febrile symptoms or thirst. A half-drachm of castor oil had previously been given, with a minute portion of Dover's powder, and at once rejected. The absence of more marked symptoms of intestinal obstruction led to a momentary error in diagnosis; and hæmatoxylin with opium was prescribed. Aromatic spirits of ammonia, carbonate of soda, and brandy, were given occasionally.

A later examination, at 7 P.M., exhibited increased prostration; absolute inappetency; shrunken face; cold extremities; finger-nails somewhat livid; pulse running, small, and frequent; fontanelle depressed; tormina frequent, without tenesmus; no dejection. An elongated horizontal firm tumour was felt to form and harden under palpation in the region of the right half of the transverse colon, then vanish gradually, and again become sensible, with contemporaneous indications of