and then a marked result followed. For nine days, the patient took two drachms of the acid daily; and then, the urine being still slightly turbid and milky, the quantity was increased to two drachms and fifteen grains daily, and was continued for seven days. On the fourth day of this dose being taken, the urine became perfectly natural, and the chylous condition did not return.

Now, had I been content with the ordinary dose of the remedy in this case, I believe that, although I might possibly have cured my patient, the time occupied in the cure would have been much longer than it was. And a very remarkable circumstance occurred in the case, showing the tolerance of the system of this large quantity of acid under the influence of the disease—a tolerance which ceased when the disease was cured. As soon as the urine assumed a healthy character, lithic acid appeared in it in large quantities, a circumstance which had never happened whilst the urine was chylous. The diminution, and subsequent stoppage, of the gallic acid caused a disappearance of the lithic acid deposit, showing the relations which the administration of the one bore to the formation of the other.

[To be continued.]

Illustrations

OF

HOSPITAL PRACTICE:

METROPOLITAN AND PROVINCIAL.

ST. GEORGE'S HOSPITAL.

HISTORY OF A RECENT EPIDEMIC OF PHAGEDÆNA OR HOSPITAL GANGRENE.

By THOMAS P. PICK, Esq., late Surgical Registrar.

MANY writers have observed that, individual epidemics of phagedæna differ essentially in their characteristics; it becomes, therefore, of interest to the surgeon to note each attack of this disease, so that he may be prepared for any phase in which it may show itself.

St. George's Hospital has, within the last few months, been visited by an epidemic of phagedæna which presents several points of interest, and which differs in many particulars from recorded accounts, particularly those of recent writers, and more especially from those cases recorded by Dr. Macleod as having occurred after the storming of the Redan, in June 1855, and by Dr. Hennen during the Peninsular War.

In order to discuss the subject systematically, it will be advisable to consider it under three heads; 1, the Cause; 2, the Symptoms; and 3, the Treatment of the disease.

1. *Cause.* In 1785, the first accurate description of this disease was given by Mr. Gillespie; and about two years later, Dr. Trotter described the same disease, and distinctly denied its contagious nature; more modern writers, however, have not agreed with this gentlemen. Mr. Blackadder, who wrote largely on this disease, believed in the complaint being *only* communicable by the direct application of the infectious matters to some part of the body from which the cuticle had been removed. The observations of Professor Thomson convincingly prove that hospital gangrene may be communicated by direct contact. He

adduces the following arguments. 1. It may be communicated by sponges, bandages, and clothing to persons at a distance from those infected with it. 2. It has been observed to attack the slightest wounds of surgeons, nurses, and others who were employed in handling and dressing infected persons, and that even in circumstances where the medical man so employed lived some distance from the patient. 3. We are often able to trace its progress distinctly from a single individual through a succession of patients. 4. We are able to prevent the progress of the disease in particular situations, by removing the infected person before the contagion has had time to operate. 5. It continues long in one ward or room without infecting others, if pains be taken to prevent intercourse between the infected and the non-infected.

Though these observations, if correct, remove all doubt as to the contagious character of this disease, there can be no doubt, at the same time, that it is also often due to the agency of some unknown poison, which finds for itself a soil fit for propagation; and I shall endeavour to prove that the late outbreak we have had in this hospital has been entirely due to infectious causes.

And to me there seems nothing incongruous in saying that a disease may be both contagious and infectious; since in all cases the disease is conveyed from one person to another by particles of matter, whether those particles be in a solid or in a gaseous form, whether they be imparted by direct contact of the two human bodies, or wafted through the air.

But the epidemic at present under notice appeared to partake most of an infectious character; and in order to prove this, I shall take the same arguments that Professor Thomson adduced to prove its contagious character in the cases he observed. 1. In no instance was it proved that it was communicated by sponges, bandages, or clothing to persons at a distance from the infected. 2. In no case did it attack any surgeon or nurse, though no special care was taken to guard any slight wounds in those dressing the patients. 3. The disease could not be traced from a single individual through a succession of cases. 4. We were not able to prevent the progress of the disease by removing patients from an infected spot. 5. It did not continue in one ward particularly without infecting others, but invaded the whole house.

The following is a brief sketch of the history of this epidemic, as it occurred among the patients who fell more especially under my notice.

In the early part of March 1863, a woman was admitted into the Hospital for necrosis of the tibia; she was placed on the operating table, and the diseased bone removed; a few days afterwards, the wound took on a phagedænic character; this, however, soon yielded to treatment. Nothing more was seen of this disease for four months. On July 14th, 1863, a patient was admitted into the accident ward, with a slight wound on the leg; on the 21st, seven days after admission, phagedæna attacked the wound. This was the first case which had occurred since March. The following day an out-patient was admitted with phagedæna. Two days afterwards, July 24th, two more cases were admitted, and were placed in separate wards. On the 27th, another case was admitted, and on this day a case broke out in the house; but, strange to say, it did not occur in any of the wards where there were phagedænic patients, but in a private room, where a patient had been totally isolated from the other patients, after undergoing amputation, for fear of the supposed contagious nature of the disease. The next case which was admitted, was on August 3rd (a large gangrenous sore on the back of the arm.) On August 5th, two cases were admitted; on

August 7th, a case broke out in the house; on August 9th, another case was admitted; on August 11th, another; on August 12th, another; on August 14th, a case broke out in the house; and so on, taking an average, three cases being admitted for every one which broke out in the house. Out of the total number of cases; viz., 92, 73 occurred out of the hospital; and of these 33 were treated from first to last as out-patients; so that out of 92 cases, only 19 broke out in the house. These rough facts are, to my mind, sufficient evidence of the epidemic nature of the disease.

One word ought here, I think, to be said upon an objection which might possibly be raised; the following theory might be proposed. Might not all the cases which were admitted have come from the same neighbourhood, and so explain how the disease might have been contagious, the virus being communicated from one person to another, possibly through the intervention of a third? The idea suggested itself, and careful inquiries were made as to the residence of the patients, and it was found that they came from all parts of the surrounding neighbourhood; from Westminster on the one side, and from Hammersmith and Barnes on the other, and between the two, from Pimlico, Chelsea, Fulham, etc.

Another objection might be raised; and that is, that the morbid matter might have been communicated from one out-patient to another, during the process of dressing, on sponges, etc.; but this fallacy has also been guarded against. For instance, in August of last year, a man was admitted with a crushed foot, for which amputation was performed. He was discharged with the wound nearly healed; he went down to Henley-on-Thames and remained there perfectly well, with the stump healing, till October 23rd, when phagedæna appeared in his stump, and he returned to London, and was readmitted. Again, a man was admitted suffering from phagedæna, who resided at King's Cross, and who had not been near the Hospital. Many other cases might be cited. It appears to me, then, that sufficient evidence has been adduced to prove that this epidemic did not arise from "overcrowding the hospital, want of cleanliness, ventilation, or from that accumulation of animal exhalations, arising from the sick and wounded, which is the source of various low forms of fever and allied diseases;" but that it was entirely due to some morbid matters in the atmosphere, some poison or ferment in the air, aided, no doubt, by want of cleanliness and ventilation among the poorer classes.

2. *Symptoms.* Regarding the symptoms, in different works, we find great diversity of opinion; and more especially as regards the constitutional symptoms, which, according to Blackadder, were secondary to the local disease, whilst Macleod and Hennen assert that the disease was always ushered in by premonitory febrile symptoms, such as pain in the head and eyes, loss of sleep, want of appetite, and quickness of the pulse, etc. In no instance, in the cases under review, was any constitutional fever observed, which might lead one to suspect the advent of phagedæna, before the local symptoms manifested themselves. The only premonitory sign which I observed was, in some cases, a total cessation of pain twenty-four hours prior to the attack; but this was by no means constant, and could not in any way be relied upon.

The constitutional symptoms, when they did appear, which was for the most part about the second or third day, were inflammatory in the first place, but quickly subsided into irritative fever, especially characterised by an extremely rapid and weak pulse.

The local symptoms presented themselves in two perfectly distinct forms, which for the sake of clear-

ness and distinction, may be designated the gangrenous and the ulcerative.

When a wound, which has been previously healthy, is about to take on the gangrenous form, it becomes dry, and, losing its florid colour, becomes of an ashy grey; the surrounding parts are swollen, tumid and tense, of a bright red hue, and present a glossy and shining appearance; the colour soon becomes darker, and eventually black; the ulcer now spreads rapidly, and has a tendency to assume a circular form; the edges become hard, everted and rugged, and the surface covered with a thick dark grey or blackish spongy slough, which can only be compared, in appearance, to a yeast poultice. There is great tendency to cedema of surrounding parts, and in some instances, to inflammation of the absorbents, and even the veins. The pain is often extremely severe, of a sharp, stabbing character, but sometimes it is comparatively slight. The disease may extend, principally, however, confining itself to the skin and cellular tissue, until large surfaces of bone and muscle are exposed; it rarely, however, attacks these structures; vessels may be ulcerated into, and alarming hæmorrhage take place. After a time, however, its progress is arrested; upon visiting our patient, we notice a little row of tiny bead-like projections around the margin of the sore, or it may be in one part only; wherever these little granulations, for such I take them to be, are found, we may rest assured that the disease is stopped. Soon, now, the sloughs are thrown off in the form of reddish-brown, or greyish viscid masses, leaving a very sensitive surface, covered with pale flabby granulations, and it is astonishing with what rapidity these ulcers heal, when there is no bone implicated; they have, however, a tendency to a recurrence of the disease.

CASE. William W., aged 24, labourer, was admitted as an out-patient, on October 1st, under Mr. Hewett. Three days before admission, he crushed his finger with a heavy stone, lacerating it very much; he applied water-dressing, but did not attend at the Hospital.

On admission, there was found to be a compound fracture of the middle phalanx, with considerable laceration and bruising of the soft parts, the injuries being such as to preclude the possibility of saving the finger. Amputation was therefore performed, and water-dressing applied.

The wound went on well, discharging healthy pus, till the sixth day, when he came complaining of intense pain in the part. The wound was found to be dry; it had lost its florid colour, and assumed a glossy coat. He was directed to apply a lotion, consisting of a drachm of solution of diacetate of lead and half an ounce of tincture of opium, with water to six ounces; and to take ten grains of chlorate of potash in an ounce and a half of distilled water every four hours, and five grains of compound soap-pill every night.

On the 9th, the whole of the flaps were black, and separated from the surrounding parts by deep excavated edges. The skin around was purple; and he complained of intense pain. He was ordered to take, every six hours, a cinchona draught with twenty minims of tincture of opium and ten grains of chlorate of potash; and to use a lotion of a drachm of carbolic acid, an ounce of glycerine, and a pint of water.

Oct. 12th. The whole of the finger was gangrenous, and the sloughing extended rapidly. The sore, which completely surrounded the finger, had acquired hard, prominent edges, the bottom of the cavity being lined with a blackish slough. His constitution had become affected; he suffered from cold sweats, had an anxious expression of countenance, and a quick, weak pulse. He was admitted into the Hospital. The

treatment was continued. He was ordered to have ordinary diet, two pints of porter, and one pint of beef-tea daily.

Oct. 15th. The sloughing did not appear to have spread since his admission, but there was no tendency to a separation of the sloughs; and there was still considerable tumefaction and purple discoloration of the surrounding parts. He was ordered to have four ounces of brandy daily.

Oct. 17th. There were two or three black spots formed in the finger beyond the line of demarcation, which had been set up. The medicine was ordered to be taken every four hours.

Oct. 19th. The sloughing was again extending. He was in great pain; tongue furred; pulse very quick and weak. There was considerable redness and swelling of the arm, and the absorbents were inflamed as far as the axillary glands, which were enlarged. Fifteen grains of sulphite of soda were added to each dose of his mixture.

October 22nd. The gangrene still continued; the whole of these tissues were destroyed down to the bone, which protruded through the end of the wound. There was still great swelling of the arm. The dose of tincture of opium was increased to twenty-five minims.

Oct. 23rd. There was still no appearance of arrest of phagedæna. The dose of laudanum was increased to thirty minims.

Oct. 25th. There was very much less swelling of the arm; and the sore had not extended during the last twenty-four hours. He did not appear to suffer at all from the large doses of opium; in fact, he slept very little.

Oct. 27th. All sloughing had ceased, and a line of demarcation had been set up. Green ointment and a linseed-meal poultice were ordered.

No further extension of the disease took place. The process of separation, granulation, and cicatrisation, ran their ordinary course; the exposed bone was removed; and he was discharged on Nov. 14th.

This form of disease which I have attempted to describe, and which I have termed the gangrenous, because it is always accompanied by the formation of a slough, is the one commonly described in our works on surgery; but there is also another form of this disease, in which there is no slough formed, in which there is rapid molecular death, or in which there is simple melting away or liquefying of the parts. When a healthy sore takes on this form of disease, it assumes a greyish appearance; the surface is covered with a semi-gelatinous, greyish-blue fluid, without any appearance of slough; the edges, instead of being hard, rugged, and everted, are levelled and are gradually lost in the ulcer, melting away as it were, so that one is almost at a loss to say where the skin terminates; the discharge is very profuse, and covers in a thick layer the surface of the sore; no doubt giving it its peculiar appearance, which once seen could not be mistaken, but which it is difficult to describe. There is no redness or œdema of the surrounding parts; in fact, they are usually possessed of an unnatural pallor. The constitutional symptoms are of a very low, asthenic type, and the disease spreads with fearful rapidity; the parts liquefying, as it were—melting away much in the same way as a lump of ice in a warm room. The same appearance of a row of minute granulations around the sore, as in the gangrenous form, indicates the arrest of the disease; when these are once formed, the sore begins to lose its greyish-blue tinge and assume a reddish colour; granulations spring up; and the ulcer cicatrises in the ordinary manner.

CASE. Joseph Short, aged 59, a pensioner, was admitted June 13th, 1863, under Mr. Lee, for necrosis

of the great toe, for which amputation was performed. On July 9th, sloughing and secondary hæmorrhage supervened, necessitating the ligation of the dorsal artery of the foot, and afterwards, upon a recurrence from another source, of the posterior tibial artery. As a consequence of this, gangrene occurred, for which amputation was performed in the lower third of the leg on July 25th. Secondary hæmorrhage again came on, owing to the atheromatous state of the vessels, in the evening of the same day, and required the application of several ligatures. On the 27th, two days after the operation, the edges of the wound were found to be extremely pale, of a greyish ashy appearance, covered with a semi-gelatinous layer, which swayed backwards and forwards on being touched, but was adherent to the structures beneath. There was no defined edge; but the skin, where it adjoined the sore, became gradually shaded off, till it eventually assumed the colour of the sore. There was no swelling or tension of the limb, which was unnaturally pale. He was in no pain; the tongue was very dry; pulse 120, very weak, and occasionally intermittent. He was ordered to apply a lotion of permanganate of potash; to take a cinchona draught with ten grains of carbonate of ammonia every six hours, and ten minims of tincture of opium in camphor mixture every two hours; and to have daily six ounces of brandy, eight ounces of port wine, ordinary diet, and one pint of porter.

July 28th. The ulceration was extending with great rapidity. The whole of the anterior flap had gone, and the skin and cellular tissue of the lower, leaving the muscles exposed, and the bone projecting out of the end of the wound. He felt better and stronger in himself. The tongue was moister; the pulse stronger. He took all his nourishment. The medicine was continued; the wine was increased to twelve ounces daily; and he was ordered to have six ounces of brandy mixture (*Pharmacopœia*) in place of the brandy.

July 29th. The extension of the disease was now arrested. Around the margin of the bone, some granulations were springing up.

From this time the disease stopped; the wound gradually cicatrised; the bone exfoliated; and the stump was all but healed, when, unfortunately, he was attacked with erysipelas, from which he died.

3. *Treatment.* In considering this, we shall pass briefly in review the various modes of treatment, both local and constitutional, which were adopted here during the late outbreak; and then sum up with a few facts gathered from the cases under observation.

Local Remedies. Of these remedies, the one which usually ranks first, and is most highly extolled, is *nitric acid*. We are told that "the extension of the sore must be stopped by the free application of fuming nitric acid." If this application did stop the disease, we should be justified in using it; but, unfortunately, it appeared to be perfectly inert to effect this end. In spite of a thorough destruction of the sides and edges of the sore with the strongest acid, the disease still progressed. In one case in which it was used, the edges were destroyed twice, and still the disease continued; and, as it causes excruciating pain, it does not seem to me advisable to use it.

Carbolic Acid, as a remedial agent, has been brought before the profession during the last two years; and two eminent French physiologists, MM. Gratiot and Lemaire, have made the important observation that, whilst it does not interfere with chemical fermentation, it completely arrests all vegetable and animal fermentations which arise from cryptogamic life. This remedy has been much extolled in checking sloughing and altering the character of ill-conditioned ulcers; and, though it does not appear to

have any power of arresting phagedæna, it is an extremely useful application in these cases in destroying the offensive smell of the secretion, and in accelerating the separation of the slough. In several cases, this remedy was applied locally, all general treatment being avoided; and in no one instance was there sufficient evidence to justify one in assuming that it arrested the disease.

The transfusion of a stream of *chlorine gas* through the ward was tried in one or two instances, though not sufficiently often to establish any definite result. On August 6th, there were three cases of phagedæna in the Fitzwilliam Ward of this hospital. They were of different duration, and were all spreading, though in one there was some slight tendency to arrest. A gentle stream of chlorine gas was passed through the ward for eighteen hours, at the end of which time the phagedæna was arrested in all three cases. A man, having undergone amputation, was placed in a separate apartment. Two days after, he was attacked with phagedæna, which rapidly extended. A stream of chlorine was passed through the ward, and in forty-eight hours the disease was arrested. Though these facts are satisfactory, they are not conclusive, and are far too meagre to arrive at any just conclusions. Whether this remedy has any control over the disease, or not, there is no doubt that it must act beneficially, if in no other way, at least in destroying the disgusting effluvia, which it does in a preeminent degree, and is thus especially useful, particularly in hospitals, where so many sick are collected together in one room.

General Remedies. There is no doubt that *opium* is our sheet-anchor. After watching the wonderful effects of this drug, not in one or two cases, but in dozens, and after every other conceivable remedy had been tried and failed, I am confident that, if there is such a thing as a specific, opium is a specific for phagedæna, if properly administered, and in sufficient quantities. I have never seen it fail. The most obstinate case was that of William W., whose history is given above. He took laudanum in gradually increasing quantities for fifteen days, till at last he was taking nearly half an ounce in the twenty-four hours. This, however, subdued the disease. But this must not be regarded as a typical case. Instances have been seen in which a rapidly spreading sore has presented a perfectly clean and healthy surface under twenty-four hours' treatment by opium; and two or three days may be generally considered as ample time to stop the most rapidly spreading sore.

Chlorate of Potass. This remedy, from its known properties in checking unhealthy and gangrenous ulcerations, was largely tried in phagedæna, and with some amount of success; for, though it did not appear to arrest the disease, at all events not with the same certainty as opium, still it appeared to act beneficially in the cleaning of sores and the separation of sloughs; and, from a knowledge of its properties, it was supposed that it might act as a prophylactic, and prevent a recurrence of the disease, and was accordingly prescribed largely.

Ammonia possesses no power over the disease, but was often a necessary addition, on account of the very rapid prostration and the great deficiency of nervous power which is often observable in these cases.

Dr. Polli of Milan has lately introduced a class of medicines before the profession, as having the power of arresting putrefactive fermentation; viz., *sulphurous acid*, in combination with potass, soda, and lime. These remedies were tried in phagedæna; and the drug did not appear to exert any influence. It was given in several cases, and in no one instance was the slightest benefit obtained.

CASE. A man was admitted with a fracture of the

leg on August 26th. On September 4th, phagedæna set in. He was ordered to take fifteen grains of sulphate of soda in water every three hours; and to use, for a lotion, a drachm of sulphite of soda to four ounces of water. He went on steadily with this treatment till September 12th, the disease still progressing; on that day, he was ordered opium, and the disease was entirely arrested in forty-eight hours.

The treatment, then, that has been adopted during the late outbreak, has been in the main opium, sometimes combined with ammonia or chlorate of potass, at other times alone; and a liberal diet. And, whatever may be the result of treatment in other epidemics, there is no doubt that, in the one under consideration, this plan was followed by the best results; in fact, it was the only one which really subdued the disease. Of forty-seven cases of which accurate records are kept, the average time which the opium took to subdue the disease was forty-eight hours, the shortest being twenty-four hours.

British Medical Journal.

SATURDAY, MARCH 5TH, 1864.

GERMANIC MEDICAL EDUCATION.

As the subject of medical education is at present exciting so much interest in the profession, our readers will be doubtless interested to know how they manage these things in foreign countries. We, therefore, introduce a summary of a report by Dr. Jaccoud, who was last year sent by the French Minister of Public Instruction into Germany to study the organisation of the Faculties of Medicine in that country. His report—the result of his labours in that direction—has just been published. In an introduction, he shows that the term of dreaminess no longer applies to Germany, in so far at least as the study of the sciences, and of medicine in particular, is concerned. Germany has thrown off the transcendentalism of Schelling and of Hegel, and is now steadily following in the footsteps of Locke and Condillac.

A German University is formed by the union of the four faculties—theology, law, medicine, and philosophy. Of such universities, Germany has now twenty-five; viz., six in Austria, six in Prussia, one in Hanover, two in Saxony, three in Bavaria, two in Baden, one in Württemberg, one in Mecklenburg-Schwerin, one in Holstein, and two in Hesse.

In each Faculty of Medicine, the instructors consist of three kinds of masters: ordinary professors, extraordinary professors, and private masters.

The number of ordinary professors is limited. The highest number varies between twelve and fourteen, and is only found in Berlin, Vienna, Prague—that is, in first-class faculties.

The mode of nomination of an ordinary professor is the same in all universities. The presentation is made by the ordinary professors themselves, and the nomination made by the crown. The faculties are