Published in final edited form as:

Prog Neuropsychopharmacol Biol Psychiatry. 2008 April 1; 32(3): 603–612. doi:10.1016/j.pnpbp.2007.07.029.

Panic, Suffocation False Alarms, Separation Anxiety and Endogenous Opioids

Maurice Preter, M.D.^{1,*} and Donald F. Klein, M.D.²

¹New York State Psychiatric Institute, Columbia University College of Physicians&Surgeons

²New York State Psychiatric Institute, Columbia University College of Physicians&Surgeons, 1051 Riverside Drive, New York, NY 10032, Phone 1-212-543-6249, e-mail DonaldK737@aol.com

Abstract

This review paper presents an amplification of the suffocation false alarm theory (SFA) of spontaneous panic (Klein, 1993). SFA postulates the existence of an evolved physiologic suffocation alarm system that monitors information about potential suffocation. Panic attacks maladaptively occur when the alarm is erroneously triggered. That panic is distinct from Cannon's emergency fear response and Selye's General Alarm Syndrome is shown by the prominence of intense air hunger during these attacks. Further, panic sufferers have chronic sighing abnormalities outside of the acute attack. Another basic physiologic distinction between fear and panic is the counter-intuitive lack of hypothalamic-pituitary-adrenal (HPA) activation in panic. Understanding panic as provoked by indicators of potential suffocation, such as fluctuations in pCO₂ and brain lactate, as well as environmental circumstances fits the observed respiratory abnormalities. However, that sudden loss, bereavement and childhood separation anxiety are also antecedents of "spontaneous" panic requires an integrative explanation. Because of the opioid system's central regulatory role in both disordered breathing and separation distress, we detail the role of opioidergic dysfunction in decreasing the suffocation alarm threshold. We present results from our laboratory where the naloxone-lactate challenge in normals produces supportive evidence for the endorphinergic defect hypothesis in the form of a distress episode of specific tidal volume hyperventilation paralleling challenge-produced and clinical panic.

Keywords

Affective neuroscience; Endogenous opioids; Panic disorder; Respiratory physiology; Separation Anxiety

Introduction

We extend the suffocation false alarm theory (SFA) of Panic Disorder (PD) (Klein, 1993) by hypothesizing that an episodic dysfunction in endogenous opioidergic regulation - a phylogenetically old system that co-regulates breathing as well as social-affiliative behavior - explains this adaptive failure. This makes it possible to integrate separation anxiety disorder,

* Corresponding author. Mailing Address: 1160 Fifth Avenue, Suite 112, New York, NY 10029. Phone 1-212-713-5336, Fax 1-212-713-5336, e-mail MP2285@columbia.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

CO₂ and lactate hypersensitivity, and a range of respiratory phenomena and pathology with Panic Disorder.

Experimental Challenge Studies: Lactate Infusion and CO₂ Inhalation in Panic

Panic Disorder is unique among psychiatric disorders in that its salient component, the panic attack, can be reliably incited in laboratory settings by specific chemical challenges as well as having challenges specifically blocked by anti-panic agents, e.g. imipramine. We can experimentally turn panic on and off, producing trenchant causally related data rather than inferences from naturalistic data. These challenge studies, using intravenous lactate infusion and carbon dioxide inhalation led to a number of unexpected laboratory findings that have advanced our understanding of clinical panic pathophysiology (see Klein, 1993 for details).

Patients who panic to CO₂ are a subset of lactate panickers (Klein, 1993). It was the recognition that increasing brain CO₂ and lactate are both harbingers of potential asphyxiation that prompted the suffocation false alarm theory of panic disorder. This theory is consonant with many recent observations detailed and expanded below.

Air Hunger (Dyspnea) and Panic Disorder

That panic is distinct from Cannon's emergency fear response (Cannon, 1920) and Selye's General Alarm Syndrome (Selye, 1956) is shown by the prominence of intense air hunger during these attacks. Acute air hunger rarely occurs in acute, external-danger initiated fear (Klein, 1993; Preter and Klein, 1998). Further, PD patients have chronic sighing abnormalities outside of the acute attack. Smoking and pulmonary complaints are independent, multiplicative risk factors for PD, but not for other anxiety disorders (Pohl et al., 1992; Amering et al., 1999). Panic is highly prevalent in lung disease (asthma, chronic obstructive pulmonary disease) and in torture victims who specifically suffered suffocation torture rather than other assaults (Bouwer and Stein, 1999).

Although increasing hypercapnia is the salient indicator of potential suffocation, hypoxia also serves this function. Beck et al. (1999; 2000) showed that panic patients respond with increased panic symptoms not only to CO₂ inhalation, but also to normocapnic hypoxia, as predicted by SFA. Patients with prominent respiratory symptoms during attacks, showed greater fluctuations in tidal volume during and after the challenge, as well as overall lower levels of end-tidal CO₂ than those whose clinical attack did not include respiratory symptoms. Equivalent increases in anxiety and panic symptoms were noted, although the sample size (seven patients in each group) limits conclusions from this particular null result. These findings support the centrality of the suffocation alarm system as a detector of the range of suffocation predictive data.

Pulmonary Conditions, Suffocation, and Panic

PD is a frequently comorbid - if not the most prevalent - psychiatric disorder among patients with pulmonary disease (Goodwin and Eaton 2003; Goodwin et al., 2004; Katon et al., 2004; Nascimento et al., 2002; Klein 2001; Roy-Byrne et al., 2006; Valenca et al., 2006; Wingate and Hansen-Flaschen, 1997; Yellowlees and Kalucy 1990; Yellowlees et al., 1987; 1988). Lung disease, including asthma and COPD may predispose to PD (Craske et al., 2001; Goodwin and Eaton 2003; Hasler et al., 2005; Karajgi et al., 1990; Perna et al., 1997; Verburg et al., 1995), or present solely with panic symptoms (Edlund et al., 1991; Sietsema et al., 1987).

Asthma and PD are both characterized by acute episodes, salient respiratory symptoms and anxiety with avoidance of situations related to acute attacks (Klein, 1993; Yellowlees and

Kalucy, 1990). There is a significantly higher (6.5 to 24%) prevalence of PD in asthmatics (Goodwin et al., 2005; Shavitt et al., 1992; Yellowlees et al., 1987; 1988) than the 1-3% reported in the general population (Kessler et al., 2006; Weissman, 1988). Perna et al. (1997) found a significantly higher prevalence of PD, sporadic panic attacks, and social phobia in asthmatics than the general population. In 90% of asthmatics with PD, asthma appeared first. Panic symptomatology during the asthmatic attack predicted longer hospitalizations in asthmatic patients (Baron et al., 1986; Brooks et al., 1989; Jurenec, 1988).

Serotonergic and tricyclic antipanic compounds may symptomatically benefit asthma (Smoller et al., 1998; Sugihara et al., 1965; Yellowlees and Kalucy, 1990). The antipanic drug, sertraline, was reported useful for comorbid anxiety and depression in COPD (Papp et al., 1995).

Cigarette Smoking is Risk Factor for Panic

Smoking and PD have been positively associated in several epidemiological studies (Amering et al., 1999; Isensee et al., 2003; Pohl et al., 1992). Breslau and Klein (1999) and Breslau et al. (2004) found that current daily smoking increased the onset risk for panic attack and PD. Quitting smoking sharply reduces risk of panic onset. Pulmonary complaints in both smokers and non-smokers increased panic risk; however, no significant risk for onset of daily smoking in persons with prior panic attacks or disorder was found. That the risk is unidirectional - from prior smoking to panic attack onset - was confirmed, after controlling for alcohol and drug use, anxiety, depressive disorders during adolescence, and parental smoking (Johnson et al., 2000).

In PD patients grouped by symptom profiles into respiratory and nonrespiratory subtypes (Biber and Alkin 1999), the respiratory group was significantly more sensitive to 35% CO₂ and smoked more cigarettes.

A large twin study (Reichborn-Kjennerud et al., 2004) found little common genetic liability for panic disorder and smoking, whereas “shared or familial environmental factors accounted for 75 % of the association between the phenotypes”. Smoking-induced lung pathology, whether manifest or sub-clinical, impairs gas exchange which may trigger panics in those with a low alarm threshold.

Separation Anxiety and Panic

Klein and Fink (1962) posited a developmental pathophysiological link between separation anxiety and PD and subsequent agoraphobia, since 50% of hospitalized agoraphobics reported severe early separation anxiety that often prevented school attendance. Further, panic, in this group, was frequently precipitated by bereavement, or separation. Therefore the anti-panic drug, imipramine, might be effective in childhood separation anxiety disorder (SAD). This was confirmed (Bernstein et al., 2000; Gittelman-Klein and Klein, 1973). These observations, coupled with attachment theory and ethological views of anxiety (Bowlby, 1973) fostered contemporary anxiety disorder classification.

Separation anxiety correlates with increased familial loading and early onset of PD (Battaglia et al., 1995). Patients highly comorbid for multiple anxiety disorders are particularly likely to recall childhood SAD (Lipsitz et al., 1994). Claims that separation anxiety equivalently antecedes other anxious states (Van der Molen et al., 1989) may be due to diagnostically ambiguous limited symptom attacks and the unreliability of the questionnaire method. However, in the only controlled, long-term, direct, blind, clinical interview follow-up of separation-anxious, school-phobic children, the only significant finding was an increased PD rate (Klein, 1995).

Silove et al. (1996) concluded that “the weight of studies support an association between early SA and adult PD; and SA appears to be linked to PD rather than to agoraphobic symptoms, but the specificity of the link remains unresolved, particularly in relation to the other anxiety and depressive disorders”. Manicavasagar et al. (2000) showed the persistence of separation anxiety symptoms from childhood to adulthood, raising the question of persistent SAD (Manicavasagar et al., 2001; 2003).

Panic disorder is more frequent in women than men (Gater et al., 1998) and commonly presents with symptoms of air hunger (Sheikh et al., 2001). The onset of panic is often triggered by separation, loss or bereavement (Faravelli and Pallanti, 1989; Kaunonen et al., 2000; Klein, 1993; Milrod et al. 2004).

Pine et al. (2000; 2005) documented a relationship between respiratory dysregulation and specific childhood anxiety disorders. Respiratory hypersensitivity to 5% CO₂ was significantly present in children with separation anxiety disorder, to a lesser degree in generalized anxiety disorder, but not in social phobia.

Panic, Cortisol, and Challenges

Another basic physiologic distinction between fear and panic is the counter-intuitive lack (possibly suppression) of hypothalamic-pituitary-adrenal (HPA) activation in panic (Sinha et al., 1999).

That the panic of PD does not trigger the hypothalamic-pituitary axis (HPA) may be explained by a deduction from the existence of a suffocation detector. Under suffocation circumstances, since acute HPA activation would counterproductively increase catabolic activity and oxygen demand, the fear response should be modified to allow energy conservative activation for possible swift escape. Relying on vagal withdrawal for rapid enhancement of cardiovascular performance (Cevese and Verlato, 1985) while suppressing HPA release seems appropriate (Porges, 1995).

A test of this idea would be measuring the response to transient hypoxia, under circumstances where fearful apprehension and motor activation is avoided. A trenchant example is provided by withdrawal of nasal continuous positive airway pressure (CPAP) in patients with obstructive sleep apnea. Strikingly, this did not trigger cortisol release despite recurrence of sleep apnea and oxygen desaturation (Grunstein et al., 1996). Gautier (1996) emphasized that hypoxia is associated with a reduced metabolic rate. This hypometabolism may be accompanied by a lowering of the thermoregulatory set point during hypoxia, both mediated by the hypothalamus. These data stress that metabolic shutdown, rather than activation, may promote survival under specific conditions such as potential suffocation.

Similarly, probands flown to high altitudes, but not stressed by exercise, did not develop hypercortisolemia unless suffering from altitude sickness (Larsen et al., 1997). When exercised, probands had an HPA response, showing that high altitude does not prevent HPA activation. Similarly, chronically hypoxic and hypercapnic patients with chronic obstructive pulmonary disease (COPD) had ordinary levels of cortisol (Hjalmarsen et al., 1996), unless they decompensated into acute respiratory failure.

Two other circumstances where hyperoxidation would be counterproductive are hyperthermia and starvation, where HPA axis activation occurs only if the subject is distressed or cachectic (which requires the lipolytic action of cortisol). In isolation, neither pleasant hyperthermia (sauna) (Kukkonen-Harjula and Kauppinen, 1988; Jokinen et al., 1991) nor brief fasting (Adamson et al., 1989) causes HPA activation. Conversely, sudden cold exposure (Hiramatsu et al., 1984) adaptively triggers HPA activation and heat production. Therefore, different

stresses produce appropriately adaptive HPA responses rather than a generic activation. This modular view of brain adaptive systems for specific dangers is supported by Corfield et al. (1995) who showed activation of cingulate gyrus and cerebellar structures in a PET study of CO₂-stimulated breathing in normal subjects. Similarly, PET scan and fMRI neuroimaging data implicate the cerebellum in the hypercapnic production of air hunger (Brannan et al. 2001; Parsons et al., 2001; Evans et al., 2002), a “compelling primal emotion like severe thirst” (Liotti et al. 2001). Parsons et al. (2001) have argued that the cerebellum regulates the overriding emotional activation that occurs under conditions of air, food and water deprivation. Notably, thirst and hunger are generally not considered as alarm systems since mortality is considerably delayed. This is not the case for air deprivation.

Opioids as Physiologic Regulators

Since separation anxiety and CO₂ sensitivity are both under opioidergic control (see below), we hypothesized that PD may be due to an episodic functional endogenous opioid deficit (amplified SFA theory).

The endogenous opioid system was unknown until the early 1970s. Naloxone prevents exogenous opiate effects, but has little effect on normal animals (Akil et al., 1998). This hindered a search for endogenous opioids. However, electrical stimulation of the periaqueductal gray (Mayer et al., 1972) produced naloxone-reversible analgesia, strongly suggesting the existence of an endogenous opioid system.

Opioid molecules are among the oldest evolved signaling substances. Remarkably conserved structurally, they are involved in diverse functions, e.g., pain perception, respiration, homeothermy, nutrient intake and immune response (Stefano et al., 1996). Their reward-signaling function may have evolved from anti-nociceptive properties. Currently three peptide groups, comprising over a dozen molecules, are identified. All arise from prohormones: Proenkephalin contains Met- and Leu-enkephalin; prodynorphin contains dynorphin A, dynorphin B, and neo-endorphin. Enkephalins and dynorphins may be the predominant central transmitters. β -Endorphin is cleaved from the prohormone, pro-opio-melano-cortin (POMC) and co-released with ACTH from the anterior pituitary. It is considered the major circulating endogenous opioid agonist.

The opioids interact with three major classes of receptors, the δ , κ and μ receptors (Reisine 1995), each with several subtypes (Connor and Christie, 1999). The enkephalins and β -endorphin have a high affinity for the μ and δ receptors, whereas dynorphin A may stimulate the κ receptor. The receptors have different affinities for the prototypical opioid antagonist, naloxone, with the μ receptor exhibiting the highest affinity. Novel opioid receptors and corresponding agonists are still regularly discovered. Morphine and codeine are synthesized by vertebrate species, including humans (Glattard et al., 2006; Stefano and Scharrer 1994; Stefano et al. 2000; Zhu et al. 2001). Our knowledge of this system is still quite incomplete.

μ Receptor activation has been seen responsible for the analgesic, respiratory and addictive effects of opioids and opiates, but more recently, δ blockade leading to reversal of μ agonist-induced respiratory depression without loss of analgesia has been described (Su et al., 1998; Verborgh and Meert, 1999). Therefore, the effects of μ active agents may partly depend on δ receptor activation. The dose of naloxone (2mg/kg) that induced panic-like reactions to lactate in normals (Preter et al., 2007, in preparation; Sinha et al., 2007) is well beyond the point of μ receptor saturation and is at the level required for δ blockade (Sluka et al., 1999).

The cranial nerves and muscles for expressing affect all evolved from the primitive gill arches that extract oxygen from water (Porges, 1997). The extent to which endogenous opioids participate in respiratory control in non-stressed human adults, i.e. under normoxic,

normocapnic conditions, remains controversial. However, their role in fetal and neonatal respiration, situations in which even small gas exchange abnormalities may be devastating, is clear (Santiago and Edelman, 1985). Endogenous opioids are activated in hypoxic or hypercapnic respiratory distress (Santiago and Edelman, 1985; Olson et al., 1996) and are inhibitory to CRH release (De Souza and Nemeroff 1989; Dunn and Berridge 1990). Opioids decrease respiratory sensitivity (Eldridge and Millhorn, 1981; Iasnetsov et al., 1984; Akiyama et al., 1990) and increase survival under hypoxic and hypercapnic conditions. Opioid modulation of CO₂ sensitivity may be of particular importance during sleep, when plasma CO₂ concentration becomes the primary breathing stimulus.

Dyspnea is modulated by central and peripheral opioid levels in both rodents and humans (Santiago and Edelman, 1985). Mice exposed to severe, intermittent hypoxia prolonged their survival during subsequent lethal suffocation (Mayfield and D'Alecy, 1992). Naloxone blocked this effect, suggesting that endogenous opioids increase adaptability to low-oxygen environments. Opioids lowered body temperature in mice, thus slowing counter-productive metabolic activity during hypoxia (Mayfield and D'Alecy, 1992). Stark et al. (1983), in a placebo-controlled trial in normal human subjects, showed that codeine allows high levels of carbon dioxide to be tolerated during breath holding. Opioid receptors, including 'non-conventional' ones, are located throughout the respiratory tract. Nebulized morphine is being investigated as a chronic dyspnea treatment (Baydur, 2004; Bruera et al., 2005; Zebraski et al. 2000).

Polyvagal Theory (Porges, 1995; 1997; 2003; 2007) may add an important structural element to SFA. Porges argues that Cannon unduly emphasized that emergency adaptations were due to sympatho-adrenal excitation. In mammals, the vagus evolved into two separate branches, both involved in the mammalian procreative process (feeding, nursing, copulation etc.). The phylogenetically older, unmyelinated dorsal vagal complex (DVC) regulates digestion and responds to novelty or threat, specifically to hypoxia, by reducing metabolic output. Oxytocinergic hypothalamic projections activate DVC output, whose sensory component monitors circulating neuropeptide levels. Porges hypothesizes that this vagal component has evolved to support, in conjunction with neuropeptide systems, mammalian bonding and attachment.

The ventral vagal complex (VVC), unique to mammals, carries myelinated vagal axons and portions of other branchiomeric cranial nerves (V, VII, IX, XI). Together, these pathways control facial expression, sucking, swallowing, breathing, crying, and vocalization. Further, the myelinated vagus (VVC) controls resting heart rate by tonic inhibition of the sinoatrial node.

Thus, VVC inhibition provides a rapid response system without the need to immediately activate the metabolically costly sympatho-adrenal system (Porges et al., 1996). The mechanism of acute tachycardia during lactate-induced panic has been attributed to vagal withdrawal (Yeragani et al., 1994) rather than sympathetic discharge. For unknown reasons, the vagal withdrawal response seems excessive in panic disorder. For instance, patients with PD show vagal withdrawal on standing in contrast to normal and depressed subjects (Yeragani et al., 1990; 1991).

The Separation Cry and Opioids

Following birth, mammalian infants cannot survive independently. Survival requires reliable distress signaling mechanisms to elicit parental care and retrieval. Distress vocalizations (DVs) are a primitive form of audio-vocal communication (Panksepp, 1998). A common neuroanatomy subserving DVs may be shared by all mammals, although substantial functional variations depend on the ontogenetic niche. The latter (West and King, 1987) signifies the

ecological and social legacies (“the inherited environment”) in which a given set of genes develops. For instance, isolated altricial (developmentally immature) infants do not emit DVs as much as many other species, since it is not likely they will stray from the nest (Panksepp et al., 1992).

Human infants are born immature and practically never get lost for their first six months. Despite frequent maternal absence, separation anxiety in humans develops only after their motor system matures. Young rats are not specifically attached to their mother, i.e. any mother will do as heater or feeder. Only once mobile do they socially bond, but their responses do not compare with the vigor seen in other species. Rats also differ from other species, including primates, dogs and chicks in their greater DV suppression by benzodiazepines (Kalin et al., 1987; Panksepp et al., 1992; Scott, 1974). Since benzodiazepines differentially alleviate anticipatory fear, social isolation in young rodents, as compared to many other mammals, may activate fear more than separation distress. Thus, when using cross-species analogies, it is important to keep in mind that the type and degree of social separation distress depends on ecological and developmental parameters (Panksepp et al., 1992).

Separation anxiety serves as a biologic leash for the increasingly mobile infant who continually checks for the mother’s presence, becomes acutely distressed on discovering her absence, and immediately attempts to elicit retrieval by crying. In humans, separation anxiety usually wanes around age four when the now verbally skilled child can successfully elicit care even from non-relatives.

Using electrical brain stimulation (ESB), DVs have been elicited in many species from homologous areas, including the midbrain, dorsomedial thalamus, ventral septum, preoptic area, and the bed nucleus striae terminalis (BNST). In some higher species, one can obtain separation calls by stimulating the central amygdala and dorsomedial hypothalamus. All these sites have high opioid receptor densities and figure heavily in sexual and maternal behaviors (Panksepp, 1998). Cortically, electrical stimulation of the rostral cingulate gyrus in monkeys consistently elicits distress calls (Jürgens and Ploog, 1970; Ploog, 1981). The cingulate cortex, found exclusively in mammals, is particularly well developed in humans and contains high densities of opioid receptors (Wise and Herkenham, 1982).

Naloxone-blockable opioid agonists reduce isolation-induced distress vocalizations (DV) across mammalian species (Hofer and Shair, 1978; Kalin et al., 1988; Kehoe and Blass, 1986; Panksepp et al., 1978). In beagles (Scott, 1974), the only psychotropic drug that yielded specific DV reduction at nonsedating doses, was imipramine, the classic antipanic agent. Naloxone given to guinea pigs and young chicks (Panksepp et al., 1978) increased baseline vocalizations (by 600%), but only when the animals were in a group, since isolates already emitted maximum DVs.

Kalin et al. (1988) studied opioid modulation of separation distress in primates, showing morphine (0.1 mg/kg) significantly decreased separation distress vocalizations without changes in autonomic and hormonal activation. Naloxone (0.1 mg/kg) blocked this effect. Sympathetic blockade using the $\alpha(2)$ agonist, clonidine, and the β adrenergic antagonist, propranolol, had no specific effect on separation-induced “coos” in infant rhesus monkeys (Kalin and Shelton, 1988).

Sighing, Yawning and Respiratory Chaos

The neural structures necessary for yawning are located near (or identical with) other phylogenetically old, respiratory and vasomotor centers. Anencephalic infants, born with only the medulla oblongata, yawn. Hypothalamic neurons originating in the paraventricular nucleus facilitate yawning by releasing oxytocin at distant sites, such as the hippocampus and ponto-

medullary structures. Opioids inhibit the yawning response at the level of the paraventricular nucleus by decreasing central oxytocinergic transmission. Conversely, naloxone may increase yawning, a classic sign of opiate withdrawal (Argiolas and Melis, 1998).

Both yawning and sighing are contagious. Observed acute inspirations may be interpreted as tests of increased ambient carbon dioxide or efforts to overcome breathlessness. Thus, observing another's yawn may incite one's own yawning test, without any relevant cognition, by activation of a phylogenetically fixed action pattern.

Venerable features of "neurosis" are frequent sighs and yawns. A feeling of respiratory oppression precedes sighing. The deep inspiration of a sigh doubles the normal tidal volume, abruptly lowers pCO₂, and relieves respiratory distress. Although PD and generalized anxiety disorder (GAD) patients were equivalent on baseline anxiety levels, Hegel and Ferguson (1997) demonstrated significantly lower baseline end-tidal CO₂ levels (EtCO₂) in PD compared to GAD and normal controls. Moreover, eight of sixteen panic patients who reported a high level of respiratory symptoms during attacks had the lowest baseline end-tidal CO₂ levels.

Comparing, at rest, panic disorder with generalized anxiety disorder (GAD) and normal controls, Wilhelm et al. (2001a) showed marked differences between PD and normals: respiratory rate was lower, tidal volume was higher, end-tidal CO₂ (EtCO₂) was lower, and the number of sighs was higher. In GAD some of these respiratory abnormalities were present in attenuated form.

Wilhelm et al. (2001b) found that panic patients at rest for 30 minutes sighed more frequently than normals. Episodic sighing, rather than sustained increases in ventilation, accounted for the decreased EtCO₂ in PD. In normals, the precipitous drop in EtCO₂ after a sigh was nullified by an immediate tidal volume decrease, thus raising EtCO₂ levels to baseline. However, panic patients continued to ventilate at an increased tidal volume for a number of post-sigh breaths, maintaining EtCO₂ at a lower level before returning to baseline. This may indicate a defense against a swift EtCO₂ increment that could trigger the suffocation alarm.

Abelson et al. (2001) studied breath-by-breath tidal volume and respiratory rate responses to a doxapram challenge in PD and a normal control group. Half of each group received a cognitive intervention designed to reduce doxapram induced anxiety/distress responses. Compared to normals, PD patients had a characteristic sighing pattern of breathing, thus producing significantly greater tidal volume irregularity. Of note, the cognitive intervention attenuated fearful response, but did not significantly influence doxapram-induced hyperventilation.

Opioids and the Control of Separation, and Social-affiliative Behavior

The first neurochemical system found to inhibit separation distress was the endogenous opioid system. Originally formulated by Panksepp, the brain opioid theory of social attachment was based on phenomenological similarities between social and narcotic dependence, including the stages of euphoria, tolerance and withdrawal. It predicted that opioid release would result in feelings of comfort and alleviation of emotional distress arising from loss and social isolation (Panksepp, 2003, 2005; Panksepp et al., 1978; 1980). Opiates, mimicking endogenous opioids, artificially create feelings of social comfort but decrease motivation to seek out social contact. Opiate antagonists increase social motivation, but reduce the reward afforded by endogenous opioid release.

This evolutionary, neurobiologic attachment theory has received much empirical support (Nelson and Panksepp, 1998). It is now appears that:

1. the endogenous opioid system is activated by several positive social interactions, ranging from mutual grooming in young animals (Keverne et al., 1989; Knowles et al., 1989) to sexual gratification;
2. opioids attenuate the reaction to social separation;
3. a low (but not a high) basal level of opioids increases motivation to seek social contact.

Panksepp (1998) hypothesizes that certain people become addicted to external opiates because they artificially induce feelings of gratification similar to - and probably above and beyond - those achieved by the release of endogenous opioids in social interactions.

Opioid Antagonists in Panic

An open pilot study (Sinha et al., 2007) showed preliminary results supporting our hypothesis of a functional endogenous opioid deficit in PD (amplified SFA theory). In 8 of 12 normal subjects, naloxone infusion (2mg/kg) followed by lactate showed significant tidal volume responses similar to those observed during clinical panic attacks (Martinez et al., 1996). Four of these subjects then received placebo-lactate and did not show this tidal volume increment. The naloxone-lactate effect resembled the non-fearful panic described by Beitman et al. (1990) in patients with attacks of cardiorespiratory distress whose cardiac catheterization was normal. DSM-4 (American Psychiatric Association; 1994) requires either acute fear or distress as necessary panic criteria. The panic patients originally studied were psychiatric inpatients whereas Beitman's patients were found in cardiac and neurological practices.

Following Sinha et al.'s report, we designed a properly controlled experimental study of the interaction of naloxone with lactate in normal subjects (Preter et al., 2007, in prep.), using the LifeShirt as a recording device (Wilhelm et al., 2003). Our initial analyses confirm Sinha et al.'s preliminary findings, lending support to our hypothesis that the endogenous opioid system serves to buffer normal subjects from the behavioral and physiological effects of lactate (Lee et al., 2007). Conversely, an episodic opioidergic deficit may underlie the suffocation sensitivity and separation anxiety of panic patients. Independently, Facchinetti et al. (1994; 1998) attributed premenstrual syndrome to periodic opioidergic deficit. Remarkably, this syndrome is also vulnerable to CO₂ and lactate induced panic attacks.

Thus, the naloxone-lactate (N-L) interaction may be an experimental model of the clinical panic attack. However, unless elaborated by a confirmatory, double-blind experiment contrasting the N-L effect in the context of a variety of anti-panic and panic irrelevant drugs, it cannot be considered definitively related to the clinical episode. If this were found to be the case, the N-L probe would afford two useful advances: First, there is currently no specific screening method for testing putative anti-panic drugs except by experimental treatment of panic disorder patients. Second, and probably of more ultimate importance, finding support for our theory that an opioidergic dysfunction is the pathophysiological mechanism underlying panic disorder allows new theoretical and practical approaches to a range of related illnesses. If opioidergic dysfunction underlies panic pathophysiology, the appropriateness of a new class of therapeutic agents comes into question. Recent work with mixed agonist-antagonists, e.g. buprenorphine (Gerra et al., 2006; Wallen et al., 2006) may be relevant.

Positive results would foster investigations into basic molecular mechanisms. For instance, we note that the dose of naloxone used in our study (2mg/kg) exceeds that needed for μ opioid receptor (MOR) blockade (Sluka et al., 1999), suggesting a role for the δ opioid receptor (DOR). Investigations of μ knockout mice and δ knockout mice (Gaveriaux-Ruff and Kieffer, 2002; Nadal et al., 2006) also indicate that the DOR is distinctively related to emotionality. A possible approach is to study lactate and CO₂ sensitivity in DOR knockout mice as compared to other mouse strains and preparations. The expectation would be that lactate and CO₂ would

distinctively elicit emotional/respiratory responses in DOR knockout as compared to other mice strains or knockout preparations.

Conclusion

Our current model provides a framework connecting PD data to endogenous opioidergic dysfunction, separation anxiety, and respiratory vulnerabilities, thus amplifying the suffocation false alarm theory (Klein, 1993). We propose that panic, separation anxiety and opioid dysfunction-related conditions, such as premenstrual dysphoria, may be due to a disturbance of endogenous opioid systems that adaptively regulate respiration, separation anxiety, consummatory pleasures, and social-affiliative rewards, in addition to pain.

The present review has focused on the possible central role of the opioid receptor in pathological panic as it occurs in PD. One of the necessary limitations of this article is that we have not critically discussed alternative views considering the relevance of other neural systems, e.g. cholinergic (Battaglia, 2002), adrenergic (Charney et al., 1990), amygdalocentric (LeDoux, 1998) etc.

Further SFA refinements are necessary to address the gastrointestinal (Fleisher et al., 2005; Lydiard 2005) and headache symptomatology of separation anxious children and some adult panic patients. That migraine headaches are highly comorbid with PD (Harter et al., 2003), and bi-directional risk factors for onset (Breslau et al., 2001) may provide clues.

Prospective, longitudinal psychobiological studies of genetic predisposition, separation, divorce, grief, bereavement, abortion, birth and adoption, in the context of challenge and therapeutic approaches offer pointed investigative opportunities. The neuroscience and evolutionary psychobiology frameworks serve as heuristic stimuli.

Our hypotheses are sufficiently concrete that falsifications, amplifications and modifications are possible (Klein, 1969). However, developing stable funding mechanisms to support such complex, longitudinal, person-oriented and physiologically sophisticated studies are a necessary precondition.

Reference List

- Abelson JL, Weg JG, Nesse RM, Curtis GC. Persistent respiratory irregularity in patients with panic disorder. *Biol Psychiatry* 2001;49:588–95. [PubMed: 11297716]
- Adamson U, Lins PE, Grill V. Fasting for 72 h decreases the responses of counterregulatory hormones to insulin-induced hypoglycaemia in normal man. *Scand J Clin Lab Invest* 1989;49:751–6. [PubMed: 2694325]
- Akil H, Owens C, Gutstein H, Taylor L, Curran E, Watson S. Endogenous opioids: overview and current issues. *Drug Alcohol Depend* 1998;51:127–40. [PubMed: 9716935]
- Akiyama Y, Nishimura M, Suzuki A, Yamamoto M, Kishi F, Kawakami Y. Naloxone increases ventilatory response to hypercapnic hypoxia in healthy adult humans. *Am Rev Respir Dis* 1990;142:301–5. [PubMed: 2382893]
- American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders*. 4. Washington, DC: American Psychiatric Association; 1994.
- Amering M, Bankier B, Berger P, Griengl H, Windhaber J, Katschnig H. Panic disorder and cigarette smoking behavior. *Compr Psychiatry* 1999;40:35–8. [PubMed: 9924875]
- Argiolas A, Melis MR. The neuropharmacology of yawning. *Eur J Pharmacol* 1998;343:1–16. [PubMed: 9551709]
- Baron C, Lamarre A, Veilleux P, Ducharme G, Spier S, Lapierre JG. Psychomaintenance of childhood asthma: a study of 34 children. *J Asthma* 1986;23:69–79. [PubMed: 3745110]

- Battaglia M, et al. Age at onset of panic disorder: influence of familial liability to the disease and of childhood separation anxiety disorder. *Am J Psychiatry* 1995;152:1362–4. [PubMed: 7653694]
- Battaglia M. Beyond the usual suspects: a cholinergic route for panic attacks. *Mol Psychiatry* 2002;7:239–46. [PubMed: 11920151]
- Baydur A. Nebulized morphine: a convenient and safe alternative to dyspnea relief? *Chest* 2004;125:363–5. [PubMed: 14769709]
- Beck JG, Ohtake PJ, Shipherd JC. Exaggerated anxiety is not unique to CO₂ in panic disorder: a comparison of hypercapnic and hypoxic challenges. *J Abnorm Psychol* 1999;108:473–82. [PubMed: 10466271]
- Beck JG, Shipherd JC, Ohtake P. Do panic symptom profiles influence response to a hypoxic challenge in patients with panic disorder? A preliminary report *Psychosom Med* 2000;62:678–83.
- Beitman BD, Kushner M, Lamberti JW, Mukerji V. Panic disorder without fear in patients with angiographically normal coronary arteries. *J Nerv Ment Dis* 1990;178:307–12. [PubMed: 2338538]
- Bellodi L, Perna G, editors. *The Panic Respiration Connection*. Milan: MDM Medical Media; 1998.
- Bernstein GA, et al. Imipramine plus cognitive-behavioral therapy in the treatment of school refusal. *J Am Acad Child Adolesc Psychiatry* 2000;39:276–83. [PubMed: 10714046]
- Biber B, Alkin T. Panic disorder subtypes: differential responses to CO₂ challenge. *Am J Psychiatry* 1999;156:739–44. [PubMed: 10327907]
- Bouwer C, Stein D. Panic disorder following torture by suffocation is associated with predominantly respiratory symptoms. *Psychol Med* 1999;29:233–6. [PubMed: 10077312]
- Bowlby, J. *Attachment and Loss*. II. New York: Basic Books; 1973.
- Brannan S, et al. Neuroimaging of cerebral activations and deactivations associated with hypercapnia and hunger for air. *Proc Natl Acad Sci USA* 2001;98:2029–34. [PubMed: 11172070]
- Breslau N, Klein DF. Smoking and panic attacks: an epidemiologic investigation. *Arch Gen Psychiatry* 1999;56:1141–7. [PubMed: 10591292]
- Breslau N, Novak SP, Kessler RC. Daily smoking and the subsequent onset of psychiatric disorders. *Psychol Med* 2004;34:323–33. [PubMed: 14982138]
- Breslau N, Schultz LR, Stewart WF, Lipton R, Welch KM. Headache types and panic disorder: directionality and specificity. *Neurology* 2001;56:350–4. [PubMed: 11171900]
- Brooks CM, Richards JM Jr, Bailey WC, Martin B, Windsor RA, Soong SJ. Subjective symptomatology of asthma in an outpatient population. *Psychosom Med* 1989;51:102–8. [PubMed: 2928458]
- Bruera E, Sala R, Spruyt O, Palmer JL, Zhang T, Willey J. Nebulized versus subcutaneous morphine for patients with cancer dyspnea: a preliminary study. *J Pain Symptom Manage* 2005;29:613–8. [PubMed: 15963870]
- Cannon, WB. *Bodily changes in pain, fear, hunger, and rage*. New York: Appleton and Co; 1920.
- Cevese A, Verlato G. Haemodynamic effects of withdrawal of efferent cervical vagal stimulation on anesthetized dogs—relative importance of chronotropic and non-chronotropic mechanisms. *J Auton Nerv Syst* 1985;14:125–36. [PubMed: 4067179]
- Charney DS, Woods SW, Nagy LM, Southwick SM, Krystal JH, Heninger GR. Noradrenergic function in panic disorder. *J Clin Psychiatry* 1990;51(Suppl A):5–11. [PubMed: 2258377]
- Connor M, Christie MD. Opioid receptor signalling mechanisms. *Clin Exp Pharmacol Physiol* 1999;26:493–9. [PubMed: 10405772]
- Corfield DR, et al. Evidence for limbic system activation during CO₂-stimulated breathing in man. *J Physiol* 1995;488(Pt 1):77–84. [PubMed: 8568667]
- Craske MG, Poulton R, Tsao JC, Plotkin D. Paths to panic disorder/agoraphobia: an exploratory analysis from age 3 to 21 in an unselected birth cohort. *J Am Acad Child Adolesc Psychiatry* 2001;40:556–63. [PubMed: 11349700]
- De Souza, EB.; Nemeroff, CB., editors. *Corticotropin-Releasing Factor: Basic and Clinical Studies of Neuropeptides*. Boca Raton: CRC Press; 1989.
- Dunn AJ, Berridge CW. Physiological and behavioral responses to corticotropin-releasing factor administration: is CRF a mediator of anxiety or stress responses? *Brain Res Brain Res Rev* 1990;15:71–100. [PubMed: 1980834]

- Edlund MJ, McNamara ME, Millman RP. Sleep apnea and panic attacks. *Compr Psychiatry* 1991;32:130–2. [PubMed: 2022111]
- Eldridge FL, Millhorn DE. Central regulation of respiration by endogenous neurotransmitters and neuromodulators. *Annu Rev Physiol* 1981;43:121–35. [PubMed: 6111289]
- Evans KC, Banzett RB, Adams L, McKay L, Frackowiak RS, Corfield DR. BOLD fMRI identifies limbic, paralimbic, and cerebellar activation during air hunger. *J Neurophysiol* 2002;88:1500–11. [PubMed: 12205170]
- Facchinetti F, Fioroni L, Martignoni E, Sances G, Costa A, Genazzani AR. Changes of opioid modulation of the hypothalamo-pituitary-adrenal axis in patients with severe premenstrual syndrome. *Psychosom Med* 1994;56:418–22. [PubMed: 7809341]
- Facchinetti F, Tarabusi M, Nappi G. Premenstrual syndrome and anxiety disorders: a psychobiological link. *Psychother Psychosom* 1998;67:57–60. [PubMed: 9556195]
- Faravelli C, Pallanti S. Recent life events and panic disorder. *Am J Psychiatry* 1989;146:622–6. [PubMed: 2712167]
- Fleisher DR, Gornowicz B, Adams K, Burch R, Feldman EJ. Cyclic Vomiting Syndrome in 41 adults: the illness, the patients, and problems of management. *BMC Med* 2005;3:20. [PubMed: 16368014]
- Gater R, Tansella M, Korten A, Tiemens BG, Mavreas VG, Olatawura MO. Sex differences in the prevalence and detection of depressive and anxiety disorders in general health care settings: report from the World Health Organization Collaborative Study on Psychological Problems in General Health Care. *Arch Gen Psychiatry* 1998;55:405–13. [PubMed: 9596043]
- Gautier H. Interactions among metabolic rate, hypoxia, and control of breathing. *J Appl Physiol* 1996;81:521–7. [PubMed: 8872614]
- Gaveriaux-Ruff C, Kieffer BL. Opioid receptor genes inactivated in mice: the highlights. *Neuropeptides* 2002;36:62–71. [PubMed: 12359497]
- Gerra G, et al. Buprenorphine treatment outcome in dually diagnosed heroin dependent patients: A retrospective study. *Prog Neuropsychopharmacol Biol Psychiatry* 2006;30:265–72. [PubMed: 16309810]
- Gittelman-Klein R, Klein DF. School phobia: diagnostic considerations in the light of imipramine effects. *J Nerv Ment Dis* 1973;156:199–215. [PubMed: 4698665]
- Glattard E, Muller A, Aunis D, Metz-Boutigue MH, Stefano GB, Goumon Y. Rethinking the opiate system? Morphine and morphine-6-glucuronide as new endocrine and neuroendocrine mediators. *Med Sci Monit* 2006;12:SR25–7. [PubMed: 16733499]
- Goodwin RD, Eaton WW. Asthma and the risk of panic attacks among adults in the community. *Psychol Med* 2003;33:879–85. [PubMed: 12877402]
- Goodwin RD, Fergusson DM, Horwood LJ. Asthma and depressive and anxiety disorders among young persons in the community. *Psychol Med* 2004;34:1465–74. [PubMed: 15724877]
- Goodwin RD, Messineo K, Bregante A, Hoven CW, Kairam R. Prevalence of probable mental disorders among pediatric asthma patients in an inner-city clinic. *J Asthma* 2005;42:643–7. [PubMed: 16266954]
- Grunstein RR, Stewart DA, Lloyd H, Akinci M, Cheng N, Sullivan CE. Acute withdrawal of nasal CPAP in obstructive sleep apnea does not cause a rise in stress hormones. *Sleep* 1996;19:774–82. [PubMed: 9085485]
- Harter MC, Conway KP, Merikangas KR. Associations between anxiety disorders and physical illness. *Eur Arch Psychiatry Clin Neurosci* 2003;253:313–20. [PubMed: 14714121]
- Hasler G, et al. Asthma and panic in young adults: a 20-year prospective community study. *Am J Respir Crit Care Med* 2005;171:1224–30. [PubMed: 15764721]
- Hegel MT, Ferguson RJ. Psychophysiological assessment of respiratory function in panic disorder: evidence for a hyperventilation subtype. *Psychosom Med* 1997;59:224–30. [PubMed: 9178332]
- Hiramatsu K, Yamada T, Katakura M. Acute effects of cold on blood pressure, renin-angiotensin-aldosterone system, catecholamines and adrenal steroids in man. *Clin Exp Pharmacol Physiol* 1984;11:171–9. [PubMed: 6378465]
- Hjalmarsen A, Aasebo U, Birkeland K, Sager G, Jorde R. Impaired glucose tolerance in patients with chronic hypoxic pulmonary disease. *Diabetes Metab* 1996;22:37–42. [PubMed: 8697294]

- Hofer MA, Shair H. Ultrasonic vocalization during social interaction and isolation in 2-week-old rats. *Dev Psychobiol* 1978;11:495–504. [PubMed: 689298]
- Iasnetsov VV, Pravdivtsev VA, Motin VG. Effect of beta-endorphin, enkephalins and their synthetic analogs on the neuronal electrical activity of the respiratory center in the medulla oblongata. *Biull Eksp Biol Med* 1984;98:687–90. [PubMed: 6095954]
- Isensee B, Wittchen HU, Stein MB, Hofler M, Lieb R. Smoking increases the risk of panic: findings from a prospective community study. *Arch Gen Psychiatry* 2003;60:692–700. [PubMed: 12860773]
- Johnson JG, Cohen P, Pine DS, Klein DF, Kasen S, Brook JS. Association between cigarette smoking and anxiety disorders during adolescence and early adulthood. *JAMA* 2000;284:2348–51. [PubMed: 11066185]
- Jokinen E, Valimäki I, Marniemi J, Seppanen A, Irjala K, Simell O. Children in sauna: hormonal adjustments to intensive short thermal stress. *Acta Physiol Scand* 1991;142:437–42. [PubMed: 1927556]
- Jurenc GS. Identification of subgroups of childhood asthmatics: a review. *J Asthma* 1988;25:15–25. [PubMed: 3053600]
- Jurgens U, Ploog D. Cerebral representation of vocalization in the squirrel monkey. *Exp Brain Res* 1970;10:532–54. [PubMed: 4988409]
- Kalin NH, Shelton SE. Effects of clonidine and propranolol on separation-induced distress in infant rhesus monkeys. *Brain Res* 1988;470:289–95. [PubMed: 3219586]
- Kalin NH, Shelton SE, Barksdale CM. Separation distress in infant rhesus monkeys: effects of diazepam and Ro 15-1788. *Brain Res* 1987;408:192–8. [PubMed: 3036301]
- Kalin NH, Shelton SE, Barksdale CM. Opiate modulation of separation-induced distress in non-human primates. *Brain Res* 1988;440:285–92. [PubMed: 3359215]
- Karajgi B, Rifkin A, Doddi S, Kolli R. The prevalence of anxiety disorders in patients with chronic obstructive pulmonary disease. *Am J Psychiatry* 1990;147:200–1. [PubMed: 2301659]
- Katon WJ, Richardson L, Lozano P, McCauley E. The relationship of asthma and anxiety disorders. *Psychosom Med* 2004;66:349–55. [PubMed: 15184694]
- Kaunonen M, Paivi AK, Paunonen M, Erjanti H. Death in the Finnish family: experiences of spousal bereavement. *Int J Nurs Pract* 2000;6:127–34. [PubMed: 11249410]
- Kehoe P, Blass EM. Opioid-mediation of separation distress in 10-day-old rats: reversal of stress with maternal stimuli. *Dev Psychobiol* 1986;19:385–98. [PubMed: 3732628]
- Kessler RC, Chiu WT, Jin R, Ruscio AM, Shear K, Walters EE. The epidemiology of panic attacks, panic disorder, and agoraphobia in the National Comorbidity Survey Replication. *Arch Gen Psychiatry* 2006;63:415–24. [PubMed: 16585471]
- Keverne EB, Martensz ND, Tuite B. Beta-endorphin concentrations in cerebrospinal fluid of monkeys are influenced by grooming relationships. *Psychoneuroendocrinology* 1989;14:155–61. [PubMed: 2525263]
- Klein DF. The prospective study of bereavement. *Arch Thanatol* 1969;1:13.
- Klein DF. False suffocation alarms, spontaneous panics, and related conditions. An integrative hypothesis. *Arch Gen Psychiatry* 1993;50:306–17. [PubMed: 8466392]
- Klein DF. Asthma and psychiatric illness. *JAMA* 2001;285:881–2. [PubMed: 11180724]
- Klein DF, Fink M. Psychiatric reaction patterns to imipramine. *Am J Psychiatry* 1962;119:432–8. [PubMed: 14033346]
- Klein RG. Is Panic Disorder Associated with Childhood Separation Anxiety Disorder? *Clin Neuropsychopharmacology* 1995;18(Suppl 2):7–14.
- Knowles PA, Conner RL, Panksepp J. Opiate effects on social behavior of juvenile dogs as a function of social deprivation. *Pharmacol Biochem Behav* 1989;33:533–7. [PubMed: 2587594]
- Kukkonen-Harjula K, Kauppinen K. How the sauna affects the endocrine system. *Ann Clin Res* 1988;20:262–6. [PubMed: 3218898]
- Larsen JJ, Hansen JM, Olsen NV, Galbo H, Dela F. The effect of altitude hypoxia on glucose homeostasis in men. *J Physiol* 1997;504(Pt 1):241–9. [PubMed: 9350634]
- LeDoux J. Fear and the brain: where have we been, and where are we going? *Biol Psychiatry* 1998;44:1229–38. [PubMed: 9861466]

- Lee YJ, Curtis GC, Weg JG, Abelson JL, Modell JG, Campbell KM. Panic attacks induced by doxapram. *Biol Psychiatry* 1993;33:295–7. [PubMed: 8471685]
- Liotti M, et al. Brain responses associated with consciousness of breathlessness (air hunger). *Proc Natl Acad Sci USA* 2001;98:2035–40. [PubMed: 11172071]
- Lipsitz JD, et al. Childhood separation anxiety disorder in patients with adult anxiety disorders. *Am J Psychiatry* 1994;151:927–9. [PubMed: 8185008]
- Lydiard RB. Increased prevalence of functional gastrointestinal disorders in panic disorder: clinical and theoretical implications. *CNS Spectr* 2005;10:899–908. [PubMed: 16273019]
- Manicavasagar V, Silove D, Curtis J, Wagner R. Continuities of separation anxiety from early life into adulthood. *J Anxiety Disord* 2000;14:1–18. [PubMed: 10770232]
- Manicavasagar V, Silove D, Rapee R, Waters F, Momartin S. Parent-child concordance for separation anxiety: a clinical study. *J Affect Disord* 2001;65:81–4. [PubMed: 11426514]
- Manicavasagar V, Silove D, Wagner R, Drobny J. A self-report questionnaire for measuring separation anxiety in adulthood. *Compr Psychiatry* 2003;44:146–53. [PubMed: 12658624]
- Martinez JM, et al. Ambulatory monitoring of respiration in anxiety. *Anxiety* 1996;2:296–302. [PubMed: 9160637]
- Mayer DJ, Wolfle TL, Akil H, Carder B, Liebeskind JC. Analgesia from electrical stimulation in the brainstem of the rat. *Science* 1971;174:1351–4. [PubMed: 5167502]
- Mayfield KP, D'Alecy LG. Role of endogenous opioid peptides in the acute adaptation to hypoxia. *Brain Res* 1992;582:226–31. [PubMed: 1393544]
- Milrod B, Leon AC, Shear MK. Can interpersonal loss precipitate panic disorder? *Am J Psychiatry* 2004;161:758–9. [PubMed: 15056531]
- Nadal X, Banos JE, Kieffer BL, Maldonado R. Neuropathic pain is enhanced in delta-opioid receptor knockout mice. *Eur J Neurosci* 2006;23:830–4. [PubMed: 16487163]
- Nascimento I, et al. Psychiatric disorders in asthmatic outpatients. *Psychiatry Res* 2002;110:73–80. [PubMed: 12007595]
- Nelson EE, Panksepp J. Brain substrates of infant-mother attachment: contributions of opioids, oxytocin, and norepinephrine. *Neurosci Biobehav Rev* 1998;22:437–52. [PubMed: 9579331]
- Olson GA, Olson RD, Kastin AJ. Endogenous opiates: 1996. *Peptides* 1997;18:1651–88. [PubMed: 9437729]
- Panksepp, J. *Affective neuroscience: The foundations of human and animal emotions*. New York: Oxford University Press; 1998.
- Panksepp J. Neuroscience. Feeling the pain of social loss. *Science* 2003;302:237–9. [PubMed: 14551424]
- Panksepp J. Why does separation distress hurt? Comment on MacDonald and Leary (2005). *Psychol Bull* 2005;131:224–30. [PubMed: 15740418]author reply 237-40
- Panksepp, J.; Newman, JD.; Insel, TR. Critical conceptual issues in the analysis of separation-distress systems of the brain. In: Strongman, KT., editor. *International review of studies on emotion*. 2. New York: Wiley; 1992.
- Panksepp J, Herman B, Conner R, Bishop P, Scott JP. The biology of social attachments: opiates alleviate separation distress. *Biol Psychiatry* 1978;13:607–18. [PubMed: 83167]
- Panksepp J, Herman BH, Vilberg T, Bishop P, DeEsquinazi FG. Endogenous opioids and social behavior. *Neurosci Biobehav Rev* 1980;4:473–87. [PubMed: 6258111]
- Papp LA, et al. Sertraline for chronic obstructive pulmonary disease and comorbid anxiety and mood disorders. *Am J Psychiatry* 1995;152:1531. [PubMed: 7573598]
- Parsons LM, et al. Neuroimaging evidence implicating cerebellum in the experience of hypercapnia and hunger for air. *Proc Natl Acad Sci USA* 2001;98:2041–6. [PubMed: 11172072]
- Perna G, Bertani A, Politi E, Colombo G, Bellodi L. Asthma and panic attacks. *Biol Psychiatry* 1997;42:625–30. [PubMed: 9376459]
- Pine DS, et al. Differential carbon dioxide sensitivity in childhood anxiety disorders and nonill comparison group. *Arch Gen Psychiatry* 2000;57:960–7. [PubMed: 11015814]
- Pine DS, et al. Response to 5% carbon dioxide in children and adolescents: relationship to panic disorder in parents and anxiety disorders in subjects. *Arch Gen Psychiatry* 2005;62:73–80. [PubMed: 15630075]

- Ploog D. Neurobiology of primate audio-vocal behavior. *Brain Res* 1981;228:35–61. [PubMed: 7023614]
- Pohl R, Yeragani VK, Balon R, Lycaki H, McBride R. Smoking in patients with panic disorder. *Psychiatry Res* 1992;43:253–62. [PubMed: 1438623]
- Porges SW. Orienting in a defensive world: mammalian modifications of our evolutionary heritage. A Polyvagal Theory. *Psychophysiology* 1995;32:301–18. [PubMed: 7652107]
- Porges SW. Emotion: an evolutionary by-product of the neural regulation of the autonomic nervous system. *Ann N Y Acad Sci* 1997;807:62–77. [PubMed: 9071344]
- Porges SW. The Polyvagal Theory: phylogenetic contributions to social behavior. *Physiol Behav* 2003;79:503–13. [PubMed: 12954445]
- Porges SW. The polyvagal perspective. *Biol Psychol* 2007;74:116–43. [PubMed: 17049418]
- Porges SW, Doussard-Roosevelt JA, Portales AL, Greenspan SI. Infant regulation of the vagal “brake” predicts child behavior problems: a psychobiological model of social behavior. *Dev Psychobiol* 1996;29:697–712. [PubMed: 8958482]
- Preter, M.; Klein, DF. Panic disorder and the suffocation false alarm theory: current state of knowledge and further implications for neurobiologic theory testing. In: Bellodi, L.; Perna, G., editors. *The Panic Respiration Connection*. Milan: MDM Medical Media; 1998.
- Reichborn-Kjennerud T, et al. Genetic and environmental influences on the association between smoking and panic attacks in females: a population-based twin study. *Psychol Med* 2004;34:1271–7. [PubMed: 15697053]
- Reisine T. Opiate receptors. *Neuropharmacology* 1995;34:463–72. [PubMed: 7566479]
- Roy-Byrne PP, Craske MG, Stein MB. Panic disorder. *Lancet* 2006;368:1023–32. [PubMed: 16980119]
- Santiago TV, Edelman NH. Opioids and breathing. *J Appl Physiol* 1985;59:1675–85. [PubMed: 2934364]
- Scott, JP. Effects of psychotropic drugs in separation distress in dogs. Amsterdam. *Proc IX Congress ECNP Exc Med*; 1974.
- Selye, H. *The stress of life*. New York: Mcgraw-Hill International Book; 1956.
- Shavitt RG, Gentil V, Mandetta R. The association of panic/agoraphobia and asthma. Contributing factors and clinical implications. *Gen Hosp Psychiatry* 1992;14:420–3. [PubMed: 1473713]
- Sheikh JI, Leskin GA, Klein DF. Gender differences in panic disorder: findings from the National Comorbidity Survey. *Am J Psychiatry* 2002;159:55–8. [PubMed: 11772690]
- Sietsema KE, Simon JI, Wasserman K. Pulmonary hypertension presenting as a panic disorder. *Chest* 1987;91:910–2. [PubMed: 3581939]
- Silove D, Manicavasagar V, Curtis J, Blaszczyński A. Is early separation anxiety a risk factor for adult panic disorder?: a critical review. *Compr Psychiatry* 1996;37:167–79. [PubMed: 8732584]
- Sinha SS, Coplan JD, Pine DS, Martinez JA, Klein DF, Gorman JM. Panic induced by carbon dioxide inhalation and lack of hypothalamic-pituitary-adrenal axis activation. *Psychiatry Res* 1999;86:93–8. [PubMed: 10397411]
- Sinha SS, Goetz RR, Klein DF. Physiological and behavioral effects of naloxone and lactate in normal volunteers with relevance to the pathophysiology of panic disorder. *Psychiatry Res* 2007;149:309–14. [PubMed: 17156855]
- Sluka KA, Deacon M, Stibal A, Strissel S, Terpstra A. Spinal blockade of opioid receptors prevents the analgesia produced by TENS in arthritic rats. *J Pharmacol Exp Ther* 1999;289:840–6. [PubMed: 10215661]
- Smoller JW, Pollack MH, Systrom D, Kradin RL. Sertraline effects on dyspnea in patients with obstructive airways disease. *Psychosomatics* 1998;39:24–9. [PubMed: 9538672]
- Stark RD, Morton PB, Sharman P, Percival PG, Lewis JA. Effects of codeine on the respiratory responses to exercise in healthy subjects. *Br J Clin Pharmacol* 1983;15:355–9. [PubMed: 6405772]
- Stefano GB, et al. Opioid and opiate immunoregulatory processes. *Crit Rev Immunol* 1996;16:109–44. [PubMed: 8879941]
- Stefano GB, et al. Endogenous morphine. *Trends Neurosci* 2000;23:436–42. [PubMed: 10941194]
- Stefano GB, Scharrer B. Endogenous morphine and related opiates, a new class of chemical messengers. *Adv Neuroimmunol* 1994;4:57–67. [PubMed: 7952829]
- Su YF, McNutt RW, Chang KJ. Delta-opioid ligands reverse alfentanil-induced respiratory depression but not antinociception. *J Pharmacol Exp Ther* 1998;287:815–23. [PubMed: 9864259]

- Sugihara H, Ishihara K, Noguchi H. Clinical experience with amitriptyline (tryptanol). in the treatment of bronchial asthma. *Ann Allergy* 1965;23:422–9. [PubMed: 5830238]
- Valenca AM, et al. The relationship between the severity of asthma and comorbidities with anxiety and depressive disorders. *Rev Bras Psiquiatr* 2006;28:206–8. [PubMed: 16924350]
- van der Molen GM, van den Hout MA, van Dieren AC, Griez E. Childhood separation anxiety and adult-onset panic disorders. *J Anxiety Disord* 1989;3:97–106.
- Verborgh C, Meert TF. The effects of intravenous naltrindole and naltrindole 5'-isothiocyanate on sufentanil-induced respiratory depression and antinociception in rats. *Pharmacol Biochem Behav* 1999;63:175–83. [PubMed: 10340539]
- Verburg K, Griez E, Meijer J, Pols H. Respiratory disorders as a possible predisposing factor for panic disorder. *J Affect Disord* 1995;33:129–34. [PubMed: 7759661]
- Wallen MC, Lorman WJ, Gosciniak JL. Combined buprenorphine and chlonidine for short-term opiate detoxification: patient perspectives. *J Addict Dis* 2006;25:23–31. [PubMed: 16597570]
- Weissman MM. The epidemiology of anxiety disorders: rates, risks and familial patterns. *J Psychiatr Res* 1988;22(Suppl 1):99–114. [PubMed: 3050062]
- West MJ, King AP. Settling nature and nurture into an ontogenetic niche. *Dev Psychobiol* 1987;20:549–62. [PubMed: 3678619]
- Wilhelm FH, Gevirtz R, Roth WT. Respiratory dysregulation in anxiety, functional cardiac, and pain disorders. Assessment, phenomenology, and treatment. *Behav Modif* 2001;25:513–45. [PubMed: 11530714]
- Wilhelm FH, Roth WT, Sackner MA. The lifeShirt. An advanced system for ambulatory measurement of respiratory and cardiac function. *Behav Modif* 2003;27:671–91. [PubMed: 14531161]
- Wilhelm FH, Trabert W, Roth WT. Characteristics of sighing in panic disorder. *Biol Psychiatry* 2001;49:606–14. [PubMed: 11297718]
- Wingate BJ, Hansen-Flaschen J. Anxiety and depression in advanced lung disease. *Clin Chest Med* 1997;18:495–505. [PubMed: 9329872]
- Wise SP, Herkenham M. Opiate receptor distribution in the cerebral cortex of the Rhesus monkey. *Science* 1982;218:387–9. [PubMed: 6289441]
- Yellowlees PM, Haynes S, Potts N, Ruffin RE. Psychiatric morbidity in patients with life-threatening asthma: initial report of a controlled study. *Med J Aust* 1988;149:246–9. [PubMed: 3412214]
- Yellowlees PM, Kalucy RS. Psychobiological aspects of asthma and the consequent research implications. *Chest* 1990;97:628–34. [PubMed: 2407451]
- Yeragani VK, et al. Decreased R-R variance in panic disorder patients. *Acta Psychiatr Scand* 1990;81:554–9. [PubMed: 2378249]
- Yeragani VK, et al. Heart rate variability in patients with major depression. *Psychiatry Res* 1991;37:35–46. [PubMed: 1862160]
- Yeragani VK, Srinivasan K, Balon R, Ramesh C, Berchou R. Lactate sensitivity and cardiac cholinergic function in panic disorder. *Am J Psychiatry* 1994;151:1226–8. [PubMed: 8037261]
- Yonkers KA, Zlotnick C, Allsworth J, Warshaw M, Shea T, Keller MB. Is the course of panic disorder the same in women and men? *Am J Psychiatry* 1998;155:596–602. [PubMed: 9585708]
- Zebraski SE, Kochenash SM, Raffa RB. Lung opioid receptors: pharmacology and possible target for nebulized morphine in dyspnea. *Life Sci* 2000;66:2221–31. [PubMed: 10855942]
- Zhu W, Bilfinger TV, Baggerman G, Goumon Y, Stefano GB. Presence of endogenous morphine and morphine 6 glucuronide in human heart tissue. *Int J Mol Med* 2001;7:419–22. [PubMed: 11254885]

Abbreviations

ACTH	adreno-cortico-tropic hormone
BNST	bed nucleus striae terminalis
COPD	

	chronic obstructive pulmonary disease
CPAP	
	continuous positive airway pressure
DVs	
	distress vocalizations
ESB	
	electrical brain stimulation
EtCO₂	
	end-tidal CO ₂
GAD	
	Generalized Anxiety Disorder
HPA	
	hypothalamic-pituitary-adrenal
N-L	
	naloxone-lactate
PCO₂	
	the partial pressure of CO ₂ (the amount of carbon dioxide gas dissolved in the blood)
PD	
	panic disorder
PET	
	positron emission tomography
POMC	
	pro-opio-melano-cortin
SAD	
	separation anxiety disorder
SFA	
	suffocation false alarm theory