Published in final edited form as:

Tetrahedron Lett. 2007 November 5; 48(45): 8029–8033.

Synthetic methodology for the preparation of *N*-hydroxysulfamides†

Krishnaswamy Devanathan, Jennifer A. Bell, Patricia C. Wilkins, Hollie K. Jacobs, and Aravamudan S. Gopalan

Department of Chemistry and Biochemistry, MSC 3C, New Mexico State University, Las Cruces, NM 88003-8001

Abstract

A convenient synthesis of a variety of substituted *N*-hydroxysulfamides from chlorosulfonyl isocyanate is reported. Alkyl groups can be introduced selectively on the *N*-Boc nitrogen of key intermediates **1a** or **1b** using the Mitsunobu reaction with alcohols. Subsequently the nitrogen carrying the silyloxy group can be alkylated under traditional conditions. Deprotection to the desired *N*-hydroxysulfamide can be achieved in high yields. Using this method, a number of structurally diverse *N*-hydroxysulfamides have been prepared. The usefulness of this methodology has further been demonstrated by the synthesis of more complex targets such as *bis*-hydroxysulfamide, **5**, and cyclic hydroxysulfamides **7** and **8**.

N-Hydroxysulfamides are a class of compounds that are structurally similar to *N*-hydroxyureasⁱ, *N*-hydroxysulfonamidesⁱⁱ, and sulfamidesⁱⁱⁱ, which exhibit a wide range of biological activity. However, little research has focused on the synthesis of *N*-hydroxysulfamides or their biological activity. Recently, the synthesis of a few *N*-alkyl, *N'*-hydroxysulfamides was published.^{iv} These compounds were shown to be potent inhibitors of several isozymes of carbonic anhydrase establishing the potential of *N*-hydroxysulfamides as pharmacophores.^v Supuran et al^{vi} have recently obtained the x-ray crystal structure of *N*-hydroxysulfamide (NH₂SO₂NHOH) with an isozyme of CA, which shows it is bound to the Zn through the ionized primary amino group. They propose that incorporation of R groups at the *N*-hydroxy position could lead to more effective enzyme inhibitors. Also, some phenylalanine-based *N*-hydroxysulfamides have been shown to be inhibitors of carboxypeptidase A.^{vii} These results prompt us to report our own synthetic efforts to develop a convenient synthetic methodology for this class of molecules.

N-hydroxysulfamides have been prepared in low yields by the direct coupling of a sulfamoyl chloride^{viii} or fluoride^{ix} with hydroxylamine. This method is limited by poor availability of sulfamoyl chlorides and hydroxylamines. The first attempt to prepare some simple *N*-hydroxysulfamides from chlorosulfonylisocyanate (CSI) gave some surprising results. CSI was reacted with *t*-BuOH to give the *N*-Boc sulfamoylchloride, which was then reacted with *O*-benzylhydroxylamine. However, attempts to deprotect the *N*-benzyloxy group using hydrogenolysis resulted in cleavage of the N-O bond yielding the corresponding sulfamide and not the desired *N*-hydroxysulfamide.^x In one example, direct coupling with unprotected

†Portions of this work were presented at the 229th American Chemical Society National Meeting, San Diego, CA, March 13–17, 2005. ORGN 525.

*Corresponding author. Tel: +1 505 646 2589; Fax: +1 505 646 2649; e-mail: agopalan@nmsu.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

hydroxylamine did lead to the unsubstituted *N*-hydroxysulfamide. More recently, a limited study was published that involved the coupling of *O*-(*t*-butyldimethylsilyl)-hydroxylamine with *N*-Boc sulfamoylchloride.⁴ Reaction of the resulting sulfamide with aliphatic alcohols under Mitsunobu conditions gave the corresponding *N*-alkyl-*N'*-hydroxysulfamides after deprotection with 5% water in TFA. However, this study was of limited synthetic scope and needed further expansion. We desired a method for the selective preparation of disubstituted *N*-hydroxysulfamides (including cyclic examples) and *bis*-*N*-hydroxysulfamides.

The known *N*-Boc-sulfamoyl chloride was prepared by the reaction of *t*-butanol with CSI and its reaction with several *O*-TBDMS protected hydroxylamines was investigated (Scheme 1). In general, a dichloromethane solution of the *N*-Boc sulfamoyl chloride (1 eq, prepared in situ) was added to a dichloromethane solution of *O*-TBDMS protected hydroxylamine (1–1.2 eq, prepared in situ) in the presence of triethylamine. After stirring the mixture overnight, the product was isolated and purified by flash chromatography. Using this procedure it was possible to prepare protected *N*-hydroxysulfamides, **1a–d**, in good yields (Scheme 1).^{xi,xii} However, the yield was lower in the coupling of the hindered *t*-butyl hydroxylamine to give **1e**.

The protected sulfamides **1** provide an excellent starting material for preparation of derivatives by alkylation. We first examined the alkylation of the Boc protected nitrogen of **1a** using Mitsunobu conditions (R¹OH, PPh₃, DEAD - Method A^{xiii}). The desired products were obtained in good yields after isolation and purification by silica gel chromatography. The Mitsunobu reaction proceeded smoothly with selectivity in the case of the unsubstituted sulfamide **1b**, clearly demonstrating the nitrogen of higher nucleophilicity. The generality of this method is clearly shown by the examples listed in Table 1.

Because of the time consuming and more difficult isolation of the products from the Mitsunobu reaction, we also examined the reactions of *N*-hydroxysulfamides **1** with alkyl halides using standard alkylation conditions (R¹X, K₂CO₃, CH₃CN, reflux-method B^{xiv}). In the case of sulfamide **1a**, *N*-alkylation using method B gave good yields of the desired products, **2a–d**, (Table 1, entries 1–4) after isolation and purification by silica gel chromatography. Alkylation using the more hindered 2-iodopropane (Table 1, entry 5) gave a lower yield of the desired product **2e**.

Alkylation of **1b** (R = H) with 1 equivalent of ethyl iodide or benzyl bromide using method B, was more complex and gave mixtures of mono (alkylation at the Boc protected nitrogen) and dialkylated products (Table 1, entries 6 & 7). It was not possible to selectively effect this alkylation by variation of the reaction conditions. Though *N*-alkylation of the more acidic Boc protected nitrogen was the faster process, it was somewhat surprising that alkylation on the hydroxylamine nitrogen also occurred readily. This led us to take advantage of this observation and exploit it for the one pot preparation of symmetrical dialkylated sulfamides. For example **2j** & **2k** were prepared in good yields using method B with 3 equivalents of butyl iodide or allyl bromide.

Introduction of different alkyl groups on the two nitrogens of **1b** was accomplished by a two-step procedure (Table 1, entries 12 & 13).^{xv} In one example, the sulfamide **1b** was first reacted with benzyl alcohol under Mitsunobu conditions (method A) to give the monoalkylated product **2g**. Subsequent alkylation of **2g** using butyl iodide (method B) gave the unsymmetrical dialkylated sulfamide **2l**. The two-step approach clearly demonstrates the potential to incorporate the desired alkyl group at the appropriate nitrogen of **1b**.

Representative sulfamides were deprotected to yield the corresponding *N*-hydroxysulfamides **3a–d** (Table 2). The Boc group was first removed by treatment of the sulfamide with 10%

trifluoroacetic acid in dichloromethane at room temperature. Subsequent removal of the TBDMS group was accomplished using 15% HCl in methanol.^{xvi}

It is exciting to note that this methodology can be extended to prepare new classes of previously unreported *N*-hydroxysulfamide ligand systems. Treatment of α , α' -dibromo-*m*-xylene with two equivalents of sulfamide **1a** in the presence of potassium carbonate in refluxing acetonitrile gave the protected sulfamide **4** in 79% yield after purification. Removal of the Boc protecting groups with 10% TFA in dichloromethane followed by treatment with 15% HCl in methanol gave the *bis-N*-hydroxysulfamide **5** in 78% yield (Scheme 2).

This methodology is also useful for the synthesis of cyclic *N*-hydroxysulfamides (Scheme 3). Alkylation of 1,3-dibromopropane with 1 equivalent of **1b** in the presence of potassium carbonate in refluxing acetonitrile followed by removal of the Boc protecting group using 10% TFA in dichloromethane gave the cyclic sulfamide **6**. Removal of the TBDMS protecting group using standard methods gave the cyclic *N*-hydroxysulfamide **7** in excellent yield. Alternatively, the sulfamide **6** could be alkylated with methyl iodide followed by removal of the TBDMS protecting group to give the cyclic *N*-methyl, *N'*-hydroxysulfamide **8**. Cyclic sulfamides have been shown to inhibit a variety of enzymes.^{1b}

In conclusion, we have developed a versatile synthetic method for the preparation of a wide array of *N*-hydroxysulfamides. Further, the methodology has been shown to be useful for the synthesis of *bis-N*-hydroxysulfamides as well as cyclic *N*-hydroxysulfamides. The present study increases the availability of diverse substrates for evaluation by researchers in the field of carbonic anhydrase inhibition, an area of much biomedical interest.

Acknowledgments

This work was supported by a grant from the National Institutes of Health under PHS Grant no. S06 GM08136.

References

- i. a) Lori F, Kelly LM, Folli A, Lisiewicz J. Expert Opinion of Drug Safety 2004;3(4):279. b) Huang J, Zou Z, Kim-Shapiro DB, Ballas SK, King SB. J. Med. Chem 2003;46:3748. [PubMed: 12904079] c) King SB. Curr. Med. Chem 2003;10:437. [PubMed: 12570692] d) Granik VG, Grigor'ev NB. Russ. Chem. Bull. Int. Ed 2002;51:1375. Paz-Ares, L.; Donehower, RC. Cancer Chemotherapy and Biotherapy-Principles and Practice, ch. 11. 3rd edn.. Chabner, BA.; Longo, DL., editors. Philadelphia: Lippincott, Williams, and Wilkins; 2001. p. 315-328. f) Frank I. J. Biol. Regulators and Homeostatic Agents 1999;13(3):186. g) Charache S, Barton FB, Moore RD, Terrin ML, Steinberg MH, Dover GJ, Ballas SK, McMahon RP, Castro O, Orringer EP. Medicine 1996;75:300. [PubMed: 8982148]
- ii. a) Nuti E, Orlandini E, Nencetti S, Rossello A, Innocenti A, Scozzafava A, Supuran CT. Bioorg. Med. Chem 2007;15:2298. [PubMed: 17276072] b) Wilkins PC, Jacobs HK, Johnson MD, Gopalan AS. Inorg. Chem 2004;43(24):7877-7881. [PubMed: 15554653] c) Lee MJC, Shoeman DW, Goon DJW, Nagasawa HT. Nitric Oxide: Biology and Chemistry 2001;5:278. d) Scozzafava A, Supuran CT. J. Med. Chem 2000;43:3677. [PubMed: 11020282] e) Mincione F, Menabuoni L, Briganti F, Mincione G, Scozzafava A, Supuran CT. J. Enzyme Inhib 1998;13:267. [PubMed: 9795865]
- iii. a) Winum J-Y, Scozzafava A, Montero J-L, Supuran CT. Med. Res. Rev 2006;26:767. [PubMed: 16710859] b) Winum J-Y, Scozzafava A, Montero J-L, Supuran CT. Expert Opinion on Therapeutic Patents 2006;16(1):27. c) Casini A, Winum J-Y, Montero J-L, Scozzafava A, Supuran CT. Bioorg. Med. Chem. Lett 2003;13:837. [PubMed: 12617903] d) Park JD, Kim DH, Kim S-J, Woo J-R, Ryu SE. J. Med. Chem 2002;45:5295. [PubMed: 12431056] e) Tozer MJ, Buck IM, Cooke T, Kalindjian SB, McDonald JM, Pether MJ, Steel KIM. Bioorg. Med. Chem. Lett 1999;9:3103. [PubMed: 10560733]
- iv. Winum J-Y, Innocenti A, Nasr J, Montero J-L, Scozzafava A, Vullo D, Supuran CT. Bioorg. Med. Chem. Lett 2005;15:2353. [PubMed: 15837324]

- v. For some recent reviews on design of carbonic anhydrase inhibitors see: a) Winum J-Y, Scozzafava A, Montero J-L, Supuran CT. *Curr. Topics Med. Chem* 2007;7:835. b) Supuran CT, Vullo D, Manole G, Casini A, Scozzafava A. *Curr. Med. Chem. Cardiovascular and Hematological Agents* 2004;2:49. c) Supuran CT, Scozzafava A, Casini A. *Med. Res. Rev* 2003;23:146. [PubMed: 12500287]
- vi. Temperini C, Winum J-Y, Montero J-L, Scozzafava A, Supuran CT. *Bioorg. Med. Chem. Lett* 2007;17:2795. [PubMed: 17346964]
- vii. Park JD, Kim DH. *Bioorg. Med. Chem* 2004;12:2349. [PubMed: 15080932]
- viii. Oettle J, Brink K, Morawietz G, Backhaus M, Boldhaus M, Bliefert CZ. *Naturforsch* 1978;33b:1193.
- ix. Boldhaus M, Brink K, Bliefert C. *Angew. Chem. Int. Ed. Engl* 1980;19(11):943.
- x. Hajri A-H, Dewynter G, Criton M, Dilda P, Montero J-L. *Heteroatom Chemistry* 2001;12(1):1.
- xi. Spectroscopic (^1H , ^{13}C , and IR) and elemental analyses of all compounds are in agreement with the assigned structures.
- xii. Representative procedure for the preparation of sulfamides **1**. Preparation of *N*-(*tert*-butoxycarbonyl)-*N'*-methyl-*N'*-[(*tert*-butyldimethyl)silyloxy]-sulfamide (**1a**): Triethylamine (2.5 mL, 17.9 mmol) was added dropwise to a suspension of *N*-methylhydroxylamine hydrochloride (0.50 g, 6.0 mmol) in anhydrous CH_2Cl_2 (10 mL) under N_2 at 0°C and stirred at rt for 2 h. The reaction mixture was cooled to 0°C and a solution of TBDMSCl (0.90 g, 6.0 mmol) in anhydrous CH_2Cl_2 (5 mL) was slowly added at 0°C . The reaction mixture was then stirred at rt for 20 h. In a separate flask, *t*-butanol (0.57 mL, 6.0 mmol) was added dropwise to a solution of chlorosulfonylisocyanate (0.51 mL, 6.0 mmol) in anhydrous CH_2Cl_2 (15 mL) under N_2 at 0°C and stirred at 0°C for 40 min to give the *t*-butoxycarbamoylsulfamoyl chloride. Triethylamine (2.5 mL, 17.9 mmol) was added to the solution of protected hydroxylamine at 0°C . The solution containing the sulfamoyl chloride was added to the protected hydroxylamine via syringe. The reaction mixture was warmed to rt and stirred for 18 h. The solvent was removed under reduced pressure. The residue was dissolved in EtOAc (100 mL) and washed with water (10 mL), 0.1 M HCl (3×10 mL), satd NaHCO_3 (3×10 mL), brine (15 mL) and dried (Na_2SO_4). The crude product was purified by flash silica gel chromatography to give **1a** (1.44 g, 71%) as a white solid. ^1H NMR (CDCl_3 , 200 MHz) δ 0.22(s, 6H), 0.94(s, 9H), 1.51(s, 9H), 3.09(s, 3H), 6.99 (br s, 1H); ^{13}C NMR (CDCl_3 , 50 MHz) δ -5.1, 17.7, 25.7, 27.8, 42.8, 84.0, 149.1; IR (KBr) 3192, 2990, 2932, 2859, 1713 cm^{-1} ; Anal calcd for $\text{C}_{12}\text{H}_{28}\text{O}_5\text{N}_2\text{SSi}$: C, 42.33; H, 8.29; N, 8.23. Found: C, 42.09; H, 7.98; N, 8.34.
- xiii. Method A-Representative procedure: Preparation of *N*-[(*tert*-butoxycarbonyl)-*N*-(ethyl)-*N'*-(methyl)-*N'*-[(*tert*-butyldimethyl)silyloxy]-sulfamide (**2a**): To a solution of **1a** (0.350 g, 1.03 mmol) and DEAD (40% in toluene, 0.448 mL, 1.03 mmol) in anhydrous THF (5 mL) under N_2 , was added a solution of PPh_3 (0.269 g, 1.03 mmol) and EtOH (0.060 g, 1.24 mmol) in anhydrous THF (3 mL) and the reaction mixture was stirred at rt for 16 h. The solvent was removed under reduced pressure and the crude product was purified by flash silica gel chromatography to give **2a** (0.281 g, 74%). ^1H NMR (CDCl_3 , 200 MHz) δ 0.22(s, 6H), 0.93(s, 9H), 1.30(t, 3H), 1.52(s, 9H), 3.08(s, 3H), 3.75 (q, 2H); ^{13}C NMR (CDCl_3 , 50 MHz) δ -5.2, 15.0, 17.7, 25.7, 27.9, 43.0, 46.07, 83.8, 151.0; IR (Neat) 2935, 2860, 1724 cm^{-1} ; Anal calcd for $\text{C}_{14}\text{H}_{32}\text{O}_5\text{N}_2\text{SSi}$: C, 45.62; H, 8.75; N, 7.60. Found: C, 45.91; H, 8.40; N, 7.41.
- xiv. Method B: Representative procedure for **2a**: To a solution of **1a** (0.150 g, 0.44 mmol) in anhydrous CH_3CN (5 mL) was added anhydrous K_2CO_3 (0.073 g, 0.53 mmol) and $\text{CH}_3\text{CH}_2\text{I}$ (0.082 g, 0.53 mmol) and the reaction mixture was heated at reflux for 4 h. The solvent was removed under reduced pressure. The residue was dissolved in EtOAc (25 mL) and washed with water (3×10 mL), brine (10 mL) and dried (Na_2SO_4). The crude product was purified by flash silica gel chromatography to give **2a** (0.122g, 75%).
- xv. For sequential alkylation on *N*-Boc protected sulfamides see: a) Montero J-L, Dewynter G-F, Agoh B, Delaunay B, Imbach J-L. *Tetrahedron Lett* 1983;24:3091. b) Regainia Z, Abdaoui M, Aouf N, Dewynter G, Montero J-L. *Tetrahedron* 2000;56:381.
- xvi. General procedure for deprotection of sulfamides: Preparation of *N*-(ethyl)-*N'*-(methyl)-*N'*-(hydroxy)sulfamide (**3c**): To a solution of **2a** (0.29 g, 0.79 mmol) in anhydrous CH_2Cl_2 (10 mL) was added trifluoroacetic acid (0.9 mL, 3.60 mmol) at rt and the reaction mixture was stirred for 1.5 h. The solvent was removed under reduced pressure to give the Boc deprotected compound (0.21 g, 99%) which was used without purification. ^1H NMR (CDCl_3 , 200 MHz) δ 0.22(s, 6H), 0.93

(s, 9H), 1.24(t, 3H), 2.93(s, 3H), 3.30–3.40(m, 2H), 4.26(br s, 1H). The Boc deprotected compound (0.177 g) was stirred with 15% HCl in methanol (3 mL) at rt. After 3 h, the solvent was removed under reduced pressure. The residue was washed with hexane and dried under vacuum to give *N*-hydroxysulfamide **3c** as a pale yellow oil (0.099 g, 97% two steps). ¹H NMR (CDCl₃, 200 MHz) δ 1.25(t, 3H), 3.02(s, 3H), 3.25–3.40(m, 2H), 4.66(br s, 1H), 6.26(br s, 1H); ¹³C NMR (CDCl₃, 50 MHz) δ 15.7, 39.8, 40.4; IR (neat) 3313, 2984, 1435 cm⁻¹; Anal calcd for C₃H₁₀O₃N₂S: C, 23.37; H, 6.54; N, 18.17. Found: C, 23.70; H, 6.93; N, 17.78.

Scheme 1.

Scheme 2.

Scheme 3.

Table 1

Alkylation of sulfamides using Method A or B

Entry	Substrate	Product	1 or 2	Method A Method B	R	R'	Method	Yield (%)
1	1a	2a	1	Method A	CH ₃	CH ₃ CH ₂	A ^d	74
2	1a	2b	1	Method B	CH ₃	Bn	B ^b	75
3	1a	2c	1	Method B	CH ₃	allyl	A	77
4	1a	2d	1	Method B	CH ₃	allyl	B	74
5	1a	2e	1	Method B	CH ₃		A	66
6	1b	2f	1	Method B	H	CH ₃ CH ₂	B	77
7	1b	2g	1	Method B	H	Bn	A	74
8	1b	2h	1	Method B	H	allyl	B	29
9	1b	2i	1	Method B	H	<i>i</i> -Pr	A	78
10	1b	2j	1	Method B	<i>n</i> Bu	<i>i</i> -Pr	A	mixture ^c
11	1b	2k	1	Method B	allyl	allyl	B	75
12	2g	2l	1	Method B	<i>n</i> Bu	Bn	A	mixture ^d
13	2i	2m	1	Method B	Bn	<i>i</i> -Pr	B	68

^a) Method A-1, DEAD, PPh₃, R'OH, THF, rt.

^b) Method B-1, R'X (1.2 eq), K₂CO₃, CH₃CN, reflux: X = I, except in 2b, 2c, 2k and 2g where X = Br

^c) 50:50 mixture of mono and dialkylated product (based on ¹H NMR)

^d) 40:60 mixture of mono and dialkylated product (based on ¹H NMR)

^e) 3 eq of RX.

Table 2

Preparation of *N*-hydroxysulfamides **3**

Entry	Substrate	Product	$ \begin{array}{c} \text{OH} \\ \\ \text{H} - \text{N} - \text{S} - \text{N} - \text{R} \\ \quad \quad \\ \text{H} \quad \text{R}^1 \quad \text{O} \\ \text{3} \end{array} $	R ¹	Yield (%) ^a
1	1c	3a	Bn	H	93
2	1d	3b	<i>n</i> -hexyl	H	90
3	2a	3c	CH ₃	CH ₃ CH ₂	97
4	2d	3d	CH ₃		85

^a) two-step yield