Published in final edited form as:

Tetrahedron. 2008 February 4; 64(6): 988–1001. doi:10.1016/j.tet.2007.10.038.

The Rhodium(II) Carbenoid Cyclization-Cycloaddition Cascade of α -Diazo Dihydroindolinones for the Synthesis of Novel Azapolycyclic Ring Systems[‡]

Dylan B. England, James M. Eagan, Gokce Meray[#], Olcay Anac[#], and Albert Padwa^{*}

Department of Chemistry, Emory University, Atlanta, GA 30322, USA

Abstract

Tandem carbonyl ylide formation-1,3-dipolar cycloaddition of α -diazo *N*-acetyl-tetrahydro- β -carbolin-1-one derivatives occur efficiently in the presence of a dirhodium catalyst to afford bimolecular cycloadducts in high yield. The Rh(II)-catalyzed reaction also takes place intramolecularly to give products derived from trapping of the carbonyl ylide dipole with a tethered alkene. The power of the intramolecular cascade sequence is that it rapidly assembles a pentacyclic ring system containing three new stereocenters and two adjacent quaternary centers stereospecifically in a single step and in high yield.

Keywords

α -diazo dihydroindolinones; rhodium(II); synthesis; carbenoid; intramolecular; dipolar cycloaddition; azapentacycles

1. Introduction

Novel strategies for the stereoselective synthesis of oxy-polycyclic ring systems continue to receive considerable attention in the field of synthetic organic chemistry.^{1–7} Construction of complex oxygen heterocycles through tandem cascade chemistry^{8–10} has been a particularly fruitful area of investigation, and the synthesis of various natural products by the use of domino reactions has been carried out by numerous investigators.^{11–14} One of the more interesting examples of this approach is the cyclization of a metallocarbenoid intermediate generated from the diazo precursor **1** onto a neighboring carbonyl group to furnish a carbonyl ylide dipole **2**.^{15,16} The cycloaddition chemistry of the resulting 1,3-dipole with various dipolarophiles has been extensively studied and also applied to the synthesis of numerous natural products.¹⁷ Highly substituted tetrahydrofuran structural units of such natural compounds as the ionophores, brevetoxins, and other marine products can be obtained by this route.¹⁸ The synthesis of illudins,¹⁹ phorbol ester derivatives,²⁰ and different sesquiterpenes²¹ has also been achieved using the carbenoid cyclization-carbonyl ylide cycloaddition process.²² In most cases, 1,3-dipolar cycloaddition reactions of carbonyl ylides proceed with varying *exo/endo*,

[‡]Dedicated to Csaba Szántay on the occasion of his 80th birthday and for his many contributions to the field of alkaloid chemistry.

^{*}Corresponding Author: Tel.: 404-727-0283; Fax: 404-727-6629; e-mail: chemap@emory.edu.

[#]Chemistry Department, Istanbul Technical University, Ayazaga-Istanbul, 34469, Turkey.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

regio-, and facial selectivity depending on the properties of the substrates and reaction conditions.¹⁵ With limited exceptions, alkyl and aryl ketones were usually the source of the interacting carbonyl group and the resulting dipole **2** was generated by the transition metal-catalyzed decomposition of the diazo alkanedione **1** in benzene at 80 °C.^{15–17}

Our group has also been interested in the formation of push-pull dipoles from the Rh(II)-catalyzed reaction of α -diazoamides²³ and noted that a smooth intramolecular 1,3-dipolar cycloaddition occurred across both alkenyl and heteroaromatic π -bonds to provide novel azapentacyclic compounds in good yield and in a stereocontrolled fashion.^{24,25} The recent synthesis of ($\pm$)-aspidophytine (**7**) nicely demonstrated the utility of this cascade methodology for the construction of complex *aspidosperma* alkaloids.²⁶ Thus, the Rh(II)-catalyzed reaction of indole **4** produced cycloadduct **6** in 97% yield *via* the intermediacy of the carbonyl ylide dipole **5**. The acid lability of cycloadduct **6** was exploited to provide the complete skeleton of aspidophytine in several additional steps (Scheme 2).

In more recent studies directed toward the total synthesis of *vinca* alkaloids,²⁷ our retrosynthetic analysis envisaged the azapentacyclic framework of 3*H*-epivincamine (**8**) to arise from a metal carbene cyclization-cycloaddition cascade (Scheme 3). Using this metal-catalyzed domino reaction as the key step, the core skeleton of 3*H*-epivincamine (**8**) retroanalyzes most straightforwardly to diazo amidoester **10**.²⁸

2. Results and Discussion

For the initial investigation into establishing the feasibility of this particular approach towards the synthesis of various members of the *vinca* alkaloid family, we chose to examine the Rh(II)-catalyzed behavior of the simpler α -diazo indolo-amide **13** as a test substrate. Compound **13** was easily prepared by treating carboline **11** with acetic anhydride under refluxing conditions followed by reaction of the resulting *N*-acetyl carboline **12** with sodium hydride and ethyl 2-diazomalonate²⁹ (Scheme 4). Heating compound **13** at 80 °C in the presence of catalytic Rh₂(OAc)₄ generates a rhodium carbenoid intermediate which undergoes cyclization with the neighboring amido group to form a transient carbonyl ylide dipole **14**. Subsequent bimolecular trapping of the dipole with various common dipolarophiles led to the expected [3+2]-cycloadducts (*i.e.* **15**). Table 1 illustrates the scope of the cycloaddition by showcasing the reaction with a variety of commercially available dipolarophiles. In a typical experiment, refluxing a solution of **13** with Rh₂(OAc)₄ in benzene at 80 °C in the presence of an equivalent of a trapping agent affords cycloadducts **16–18** in *ca* 75% yield with complete regio- and diastereoselectivity. In general, the cascade reaction sequence produced only the *exo*-cycloaddition product isolated as a single diastereomer. When benzaldehyde was used as the trapping agent, the cycloaddition also proceeded with high selectivity and afforded cycloadduct **19** in 85% yield as the only isolable product. The exclusive formation of cycloadducts **17–19** is consistent with the regiochemistry predicted by FMO theory. The most favorable interaction is between the HOMO of the carbonyl ylide dipole and the LUMO of the dipolarophile. The preferred *exo*-selectivity is associated with fewer non-bonded interactions in the transition state for the cycloaddition process.

Push-pull carbonyl ylide dipoles represent useful intermediates for the synthesis of complex azapolycyclic frameworks as found in different classes of alkaloids. Since only limited reports are available describing the bimolecular cycloaddition behavior of these dipoles,²⁵ we thought it would be worthwhile to further probe the facility of their reactivity with various trapping agents. With this in mind, the Rh(II)-induced carbenoid cyclization-cycloaddition sequence was extended to the α -diazo oxindole system. Our initial studies centered on the bimolecular cycloaddition chemistry of α -diazo dihydroindolinone **20a** (R₁ = Me; R₂ = OMe). Our intention

was to examine the regio and stereoselective aspects of the reaction (Scheme 5) using α -diazo oxindolo-amide **20a** as a model system.

Toward this goal, α -diazo dihydroindolinone **20a** was treated with dimethylacetylene dicarboxylate (DMAD) in the presence of catalytic $\text{Rh}_2(\text{OAc})_4$ at 80 °C in benzene and this resulted in the formation of cycloadduct **23** in 98% yield. A similar cycloaddition cascade occurred when the reaction was carried out using methyl propiolate. In this case, a 1:1-mixture of both regioisomeric adducts **24a** and **25a** were obtained in 80% overall yield. The observed regioselectivity of the cycloaddition is significantly lower with this particular α -diazo amide than that previously encountered with the related diazo indolo amide **13**. The exact reason for this difference in product distribution is unclear but is speculated to be related to steric interactions in the transition state leading to the type I cycloadduct **24a** (HOMO dipole controlled). These steric interactions are more severe in the transition state when the carbomethoxy group of the dipolarophile and the substituent groups on the oxindole are adjacent to each other (*i.e.*, **24a**). Consequently, the electronically less preferred FMO product **25a** becomes more significant with this system and stands in contrast with the results encountered with the related diazo indolo amide **13**. In support of this suggestion we have also studied the Rh(II)-catalyzed reaction of α -diazo dihydroindolinone **20b** which contains the much larger TBS group attached to the oxindole ring.

Reaction of this system with methyl propiolate afforded a 1:3-mixture of regioisomeric cycloadducts in 70% overall yield. The major adduct formed (**25b**) corresponds to the sterically less crowded product and not the one expected on the basis of FMO considerations.

The Rh(II)-catalyzed reaction was also carried out using **20a** with *N*-phenylmaleimide as the added dipolarophile. The ^1H -NMR of the crude reaction mixture showed a 3:2-mixture of *exo/endo*-diastereomers in 95% yield. We did not observe any product resulting from the cycloaddition of the transient carbonyl ylide across the C=O group of *N*-phenylmaleimide. Another experiment that was performed involved the use of **20a** with methyl acrylate in the presence of $\text{Rh}_2(\text{OAc})_4$ as described above. The ^1H -NMR of the crude reaction showed a mixture of four diastereomers in the ratio of 4.5:2:1.5:1. Chromatographic purification of the reaction mixture did not lead to the complete separation of the various isomers and thus the determination of the stereochemistry of each cycloadduct remains undefined. Nevertheless, on the basis of similarity in spectral data, it is clear that the cycloaddition produced a mixture of regioisomers in both the *exo* and *endo* orientations and without any significant selectivity differences.

After performing the cycloaddition of **20a** with various C=C bonds, we decided to extend the cascade reaction using benzaldehyde as the trapping agent. A survey of the literature revealed that only a few reports are available dealing with the reactions of carbonyl ylides with carbonyl groups as hetero-dipolarophiles.^{30,31} The Rh(II)-catalyzed reaction of diazo amide **20a** was carried out in the presence of a slight excess of benzaldehyde in benzene at 80 °C. The reaction was monitored by TLC and column chromatographic purification of the crude reaction mixture afforded cycloadduct **28** as the exclusive cycloadduct in 98% yield (see Figure 2). The assignment of the *exo* orientation of the phenyl group on the dioxo bicyclic framework is based on extensive precedent for related transformations.³¹ The regiochemistry observed is readily rationalized in terms of maximum overlay of the dipole HOMO-dipolarophile LUMO and the high selectivity observed may be due to the much larger coefficient on the oxygen atom of the carbonyl group.

The cascade reactions discussed above are of interest due to the potential for rapid generation of molecular complexity from a readily available diazo substrate. Earlier studies from our group established that the Rh(II)-catalyzed tandem carbonyl ylide formation-1,3-dipolar

cycloaddition also occurred intramolecularly when non-activated C=C bonds were used to trap the dipole.³² At this point, we decided to study the intramolecular cycloaddition behavior of several related α -diazo dihydroindolinones. In this spirit we first prepared α -diazo dihydroindolinone **31** starting from isatin and the Grignard reagent derived from 4-bromo-1-butene (Scheme 6). The resulting alcohol **29** was converted to the corresponding TBS ether **30** using TBSOTf and the required diazo functionality was then introduced using 2-diazo malonyl chloride.²⁹ Exposure of **31** to the standard Rh(II) reaction conditions afforded the intramolecular cycloadduct **32** as the only product in 70% isolated yield. We also carried out a similar cyclization-cycloaddition cascade with the homologous α -diazo dihydroindolinone **35** which contains a five-carbon tether, $n=2$ (to form a six-membered ring). The resulting cycloadduct **36** was isolated in comparable yield (*i.e.*, 69%). Even in the presence of DMAD, the same two cycloadducts (**32** and **36**) were obtained as the only products of the cycloaddition reaction. Thus, the rapidity of the intramolecular cycloaddition of both α -diazo dihydroindolinones **31** and **35** significantly overshadows any bimolecular processes. The power of this intramolecular cascade sequence is that it rapidly assembles a pentacyclic ring system containing three new stereocenters and two adjacent quaternary centers stereospecifically in a single step and in high yield. What is also noteworthy about the intramolecular cycloaddition is that it creates a ring fused to the cycloadduct, the size of which is dictated by the length of the tether connected to the dipolarophile.

Bolstered by this positive result, we next examined the Rh(II)-catalyzed chemistry of several related alkenyl ethers (*i.e.*, **37–39**). Exposure of **37** to Rh₂(OAc)₄ in benzene produced cycloadduct **40** in 72% yield (Scheme 7). A related set of results was encountered with α -diazo dihydroindolinones **38** and **39**. The intramolecular cycloaddition reaction occurred smoothly giving rise to cycloadducts **41** and **42** in 82% and 80%, respectively.

Of the many questions concerning the factors that govern the outcome of intramolecular carbonyl ylide trapping reactions, one that is easily formulated focuses upon the course of the reaction as a function of the length of the tether connecting the reacting groups. The primary spatial requirement for intramolecular cycloaddition is that the distance between the dipole and alkene should be sufficiently close so that effective overlap of the π -orbitals can occur. Our earlier studies showed that the ring size of the resulting dipole also played a significant role in the efficiency of the tandem cyclization-cycloaddition process.³³ After studying the successful intramolecular cycloaddition reactions of the α -diazo 3-alkoxy substituted dihydroindolinones **37–39**, we embarked on a study whose primary focus was concerned with exploration of the chemoselectivity of the cycloaddition across different π -bonds contained within the same substrate. To perform the Rh(II)-catalyzed tandem ylide generation and competitive cycloaddition reactions, α -diazo dihydroindolinones **43–45** were selected as prototypical substrates. Treatment of the α -diazo substrate **43** with catalytic Rh₂(OAc)₄ afforded a 5:1-mixture of two cycloadducts. The major product isolated (**46**, 55%) corresponded to internal trapping of the carbonyl ylide intermediate across the ether substituted π -bond (Scheme 8). The minor product (**47**, 11%) was that derived from cycloaddition across the alkenyl π -bond contained within the all-carbon tether. Thus, there is a clear preference for cycloaddition across the ether tethered alkene even though the distance connecting the two reacting centers is essentially the same. The effect of changing the tether length was then probed by studying the Rh(II)-metal catalyzed reaction of α -diazo dihydroindolinones **44** and **45**. The Rh(II)-catalyzed reaction of **44**, which contains two unequal tether units, proceeded smoothly and afforded cycloadduct **48** in 79% yield as the exclusive product. Reaction of the isomeric α -diazo dihydroindolinone **45** also occurred readily in the presence of a catalytic amount of Rh₂(OAc)₄ to give a single intramolecular cycloadduct **49** in 75% yield. As was the case with the homologous system **44**, cycloaddition of the carbonyl ylide dipole took place exclusively across the ether substituted π -bond.

One explanation that could account for the preferred selectivity observed is that the intramolecular cycloaddition reactions of α -diazo hydroindolinones **43–45** are governed by FMO factors. Regiochemical control in [3+2] dipolar cycloadditions has generally been rationalized on the basis of FMO considerations.³⁴ For carbonyl ylides, the HOMO of the dipole is dominant for reactions with electron deficient dipolarophiles, while the LUMO becomes important for cycloaddition to more electron-rich species.³⁵ The presence of an electron withdrawing oxygen atom (inductive effect) might be expected to lower the LUMO level of the proximal π -bond and thereby promote the HOMO-controlled cycloaddition. Although this suggestion is not unreasonable, it seems to us as though the oxygen atom is simply too far removed from the π -bond to significantly influence its LUMO energy level. Rather, we suspect that the origin of the rate differences resides in either ground state conformational effects or in relative strain within the transition states. With the all carbon tether, the two methylene carbons adjacent to the quaternary center probably adopt an *anti*-conformation so as to minimize non-bonded interactions, and the olefinic π -bond is thus projected away from the carbonyl ylide dipole. A *gauche*-conformation about these two methylene carbons is required for the cycloaddition reaction to proceed. Placement of an oxygen atom within the tether will change the preferred conformation and this effectively places the olefinic π -bond in closer proximity to the dipole. Consequently, the minimum energy conformers within the ether linkage are closer to the reactive conformers necessary for the cycloaddition, thereby facilitating the reaction.

3. Conclusion

In conclusion, several trends have surfaced from our investigations in this area. First and foremost, these studies have demonstrated that the intramolecular tandem cyclization-cycloaddition reaction of α -diazo dihydroindolinones is a viable method for quickly assembling complex azapolycyclic ring systems from easily prepared precursors. Both alkenes and tethered alkenyl ethers readily undergo the intramolecular cycloaddition giving rise to five, six and seven-membered rings fused to a dihydroindolinone backbone. The distribution of regioisomeric products from the internal cycloaddition appears to correlate well with conformational effects in the transition state for the reaction. We are continuing to explore the scope, generality and synthetic applications of the Rh(II)-catalyzed tandem cyclization-cycloaddition reaction of α -diazo dihydroindolinones and will report additional findings at a later date.

4. Experimental

4.1 General

Melting points are uncorrected. Mass spectra were determined at an ionizing voltage of 70eV. Unless otherwise noted, all reactions were performed in flame dried glassware under an atmosphere of dry nitrogen. Solutions were evaporated under reduced pressure with a rotary evaporator and the residue was chromatographed on a silica gel column using an ethyl acetate-hexane mixture as the eluent unless specified otherwise.

4.1.1 2-Acetyl-2,3,4,9-tetrahydro- β -carbolin-1-one (12**)**—To a stirred solution of 2.4 g (12.9 mmol) of carboline **11**³⁶ in 100 mL of refluxing toluene was added 3.6 mL (26 mmol) of triethylamine and 6.1 mL (65 mmol) of acetic anhydride. The resulting mixture was heated at reflux for 8 h and the solvent was removed under reduced pressure. The residue was partitioned between EtOAc and water and the layers were separated. The aqueous layer was washed twice with EtOAc. The combined organic extracts were washed with a saturated sodium bicarbonate solution, brine, and dried over Na₂SO₄. The organic extracts were filtered and concentrated under reduced pressure. The crude residue was subjected to flash silica gel chromatography to give 1.9 g (65%) of carboline **12** as a white solid; mp 229–230 °C; IR (neat)

3301, 2934, 1687, 1663, and 1287 cm^{-1} ; $^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 9.10 (brs, 1H), 7.64 (d, 1H, $J = 8.3$ Hz), 7.45 (d, 1H, $J = 8.7$ Hz), 7.39 (t, 1H, $J = 8.3$ Hz), 7.19 (t, 1H, $J = 8.7$ Hz), 4.35 (t, 2H, $J = 6.4$ Hz), 3.07 (t, 2H, $J = 6.4$ Hz), and 2.67 (s, 3H); $^{13}\text{C-NMR}$ (100 MHz, CDCl_3) δ 20.8, 27.3, 43.9, 112.5, 120.8, 121.1, 124.1, 124.8, 126.4, 126.7, 138.6, 161.8, and 173.1

4.1.2 3-(2-Acetyl-1-oxo-1,2,3,4-tetrahydro- β -carbolin-9-yl)-2-diazo-3-oxo-propionic Acid Ethyl Ester (13**)**—To a suspension of 0.47 g (12 mmol) of NaH in 20 mL of THF cooled to 0 $^\circ\text{C}$ was added 1.8 g (8 mmol) of carboline **12** dissolved in 30 mL of THF *via* cannula. The solution was allowed to stir at 0 $^\circ\text{C}$ for 30 min, after which time 2.1 g (12 mmol) of ethyl 2-diazomalonyl chloride²⁹ dissolved in 20 mL of THF was added. The resulting mixture was stirred at 0 $^\circ\text{C}$ for 2 h, quenched with water, and extracted with EtOAc. The combined organic layers were washed with brine and dried over Na_2SO_4 . The organic extracts were filtered and concentrated under reduced pressure. The crude residue was subjected to flash silica gel chromatography to give 2.0 g (70%) of α -diazo indole **13** as a yellow oil: IR (neat) 2133, 1720, 1683, 1650, and 1307 cm^{-1} ; $^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 7.87 (d, 1H, $J = 8.6$ Hz), 7.60 (d, 1H, $J = 7.9$ Hz), 7.46 (t, 1H, $J = 8.6$ Hz), 7.30 (t, 1H, $J = 7.9$ Hz), 4.36–4.29 (m, 2H), 4.17 (q, 2H, $J = 7.0$ Hz), 3.00 (t, 2H, $J = 6.4$ Hz), 2.59 (s, 3H), and 1.22 (t, 3H, $J = 7.0$ Hz); $^{13}\text{C-NMR}$ (100 MHz, CDCl_3) δ 14.2, 21.0, 27.4, 42.0, 61.6, 114.1, 120.9, 123.5, 125.3, 128.5, 128.6, 129.3, 139.1, 159.9, 160.3, 161.7, and 172.8.

General Procedure for the Rh(II)-Catalyzed Dipolar Cycloaddition: α -Diazo-indole **13** (0.27 mmol) was stirred with rhodium(II) acetate (2 mg) along with the appropriate dipolarophile (0.4 mmol) in benzene (5 mL) and the mixture was heated to reflux for the indicated time. At the end of this time, the mixture was allowed to cool to rt and the solvent was removed under reduced pressure. The crude residue was purified as described to give cycloadducts **16–19**.

4.1.3 Rh(II)-Catalyzed Dipolar Cycloadduct 16—The general procedure described above was followed using 0.1 g (0.27 mmol) of α -diazo indole amide **13**, 0.4 g (0.4 mmol) of maleic anhydride, and 2 mg of rhodium(II) acetate in 5 mL of benzene. The resulting mixture was heated at reflux for 1 h and the solvent was removed under reduced pressure. The crude residue was subjected to trituration with Et_2O to give 0.09 g (75%) of cycloadduct **16** as a white solid: mp 246–247 $^\circ\text{C}$; IR (neat) 2975, 1788, 1755, 1731, and 1662 cm^{-1} ; $^1\text{H-NMR}$ (400 MHz, CD_3CN) δ 8.17 (d, 1H, $J = 7.0$ Hz), 7.65 (d, 1H, $J = 7.6$ Hz), 7.47–7.37 (m, 2H), 4.45–4.37 (m, 3H), 4.06–3.86 (m, 3H), 3.05–2.86 (m, 2H), 2.22 (s, 3H), and 1.38 (t, 3H, $J = 7.0$ Hz); HRMS Calcd. for $[\text{C}_{22}\text{H}_{18}\text{N}_2\text{O}_8 + \text{H}^+]$: 439.1136. Found 439.1137.

4.1.4 Rh(II)-Catalyzed Dipolar Cycloadduct 17—The general procedure described above was followed using 0.1 g (0.27 mmol) of α -diazo indole amide **13**, 0.04 mL (0.4 mmol) of methyl acrylate, and 2 mg of rhodium(II) acetate in 5 mL of benzene. The resulting mixture was heated at reflux for 1 h and the solvent was removed under reduced pressure. The crude residue was subjected to flash silica gel chromatography to give 0.09 g (77%) of cycloadduct **17** as a white solid: mp 176–177 $^\circ\text{C}$; IR (neat) 2987, 2953, 1755, 1736, 1655, 1455, and 1385 cm^{-1} ; $^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 8.20 (d, 1H, $J = 7.9$ Hz), 7.51 (d, 1H, $J = 7.9$ Hz), 7.43–7.34 (m, 2H), 5.30–5.20 (m, 1H), 4.46–4.39 (m, 2H), 3.75 (s, 4H), 3.45–3.40 (m, 2H), 2.90–2.72 (m, 3H), 2.26 (s, 3H), and 1.37 (t, 3H, $J = 7.0$ Hz); $^{13}\text{C-NMR}$ (100 MHz, CDCl_3) δ 14.0, 20.4, 25.1, 33.7, 40.9, 49.8, 53.0, 62.9, 85.0, 116.1, 119.6, 125.1, 126.3, 127.9, 134.6, 164.2, 164.9, and 170.3.

4.1.5 Rh(II)-Catalyzed Dipolar Cycloadduct 18—The general method described above was followed using 0.1 g (0.27 mmol) of α -diazo indole **13**, 0.04 mL (0.4 mmol) of methyl

propiolate, and 2 mg of rhodium(II) acetate in 5 mL of benzene. The resulting mixture was heated at reflux for 1 h and the solvent was removed under reduced pressure. The crude residue was subjected to flash silica gel chromatography to give 0.09 g (78%) of cycloadduct **18** as a white solid; mp 211–212 °C; IR (neat) 2950, 1753, 1716, 1675, and 1655 cm⁻¹; ¹H-NMR (400 MHz, CD₃CN) δ 8.04 (d, 1H, *J* = 7.9 Hz), 7.60 (d, 1H, *J* = 7.0 Hz), 7.42–7.33 (m, 2H), 7.15 (s, 1H), 4.40–4.35 (m, 2H), 4.27–4.24 (m, 1H), 3.66 (s, 3H), 3.55–3.48 (m, 1H), 2.93–2.91 (m, 2H), 2.24 (s, 3H), and 1.33 (t, 3H, *J* = 7.3 Hz).

4.1.6 Rh(II)-Catalyzed Dipolar Cycloadduct 19—The general method described above was followed using 0.1 g (0.27 mmol) of α-diazo indole **13**, 0.04 mL (0.4 mmol) of benzaldehyde, and 2 mg of rhodium(II) acetate in 5 mL of benzene. The resulting mixture was heated at reflux for 1 h and the solvent was removed under reduced pressure. The crude residue was subjected to flash silica gel chromatography to give 0.1 g (85%) of cycloadduct **19** as a white foamy solid; IR (neat) 2984, 1754, 1729, 1686, 1664, and 1373 cm⁻¹; ¹H-NMR (400 MHz, CDCl₃) δ 7.97–7.95 (m, 1H), 7.60–7.57 (m, 1H), 7.45–7.34 (m, 2H), 7.27–7.18 (m, 5H), 5.86 (s, 1H), 4.70 (ddd, 1H, *J* = 13.3, 9.2, and 4.4 Hz), 4.52–4.39 (m, 2H), 3.80 (ddd, 1H, *J* = 13.3, 9.2, and 4.4 Hz), 3.03–2.87 (m, 2H), 2.54 (s, 3H), and 1.40 (t, 3H, *J* = 7.3 Hz); ¹³C-NMR (100 MHz, CDCl₃) δ 13.9, 20.2, 24.9, 43.1, 63.4, 79.3, 88.7, 106.9, 116.0, 117.2, 120.0, 125.2, 126.2, 126.6, 127.6, 128.0, 128.5, 128.6, 129.1, 130.7, 133.0, 133.9, 160.3, 163.7, and 172.5; HRMS Calcd. for [C₂₅H₂₂N₂O₆+H⁺]: 447.1551. Found 447.1557.

4.1.7 2-Diazo-3-(3-methoxy-3-methyl-2-oxo-2,3-dihydro-indol-1-yl)-3-oxo-propionic Acid Ethyl Ester (20a)—To a solution of 0.07 g (0.4 mmol) of 3-methoxy-3-methyl-1,3-dihydro-indol-2-one³⁷ in 10 mL of THF at 0 °C was added to 0.03 g (0.8 mmol) of NaH (60% dispersion in mineral oil). After stirring for 30 min, a 0.12 g (0.8 mmol) sample of ethyl 2-diazomalonate²⁹ was added and the solution was stirred 3 h. The mixture was then quenched with water and extracted with EtOAc. The combined organic layers were washed with water, brine, and dried over Na₂SO₄. After removal of the solvent under reduced pressure, the residue was subjected to flash silica gel chromatography to give 0.1 g (85%) of **20a** as a pale yellow oil; IR (neat) 2986, 2144, 1770, 1729, 1666, 1608, and 1466 cm⁻¹; ¹H-NMR (400 MHz, CDCl₃) δ 1.31 (t, 3H, *J* = 7.3 Hz), 1.63 (s, 3H), 3.09 (s, 3H), 4.26–4.32 (m, 2H), 7.25 (t, 1H, *J* = 7.3 Hz), 7.36–7.40 (m, 2H), and 7.66 (d, 1H, *J* = 7.6 Hz); ¹³C-NMR (100 MHz, CDCl₃) δ 14.3, 24.8, 53.5, 62.0, 80.0, 114.4, 123.8, 125.3, 128.0, 130.0, 139.2, 160.0, 160.2, and 175.8.

4.1.8 Rh(II)-Catalyzed Dipolar Cycloadduct 23—To a solution containing 0.1 g (0.32 mmol) of diazo hydroindolinone **20a** in 15 mL of benzene was added 0.2 mL (1.6 mmol) of dimethylacetylene dicarboxylate and a catalytic amount of Rh₂(OAc)₄. The mixture was heated at reflux for 2 h. The reaction mixture was then concentrated under reduced pressure and subjected to flash silica gel chromatography to give 0.14 g (98%) of cycloadduct **23** as a colorless oil; IR (neat) 2991, 2954, 2832, 2279, 1732, 1600, and 1442 cm⁻¹; ¹H-NMR (400 MHz, CDCl₃) δ 1.38 (t, 3H, *J* = 7.1 Hz), 2.05 (s, 3H), 3.22 (s, 3H), 3.45 (s, 3H), 3.92 (s, 3H), 4.39 (q, 2H, *J* = 7.0 Hz), 7.00–7.03 (m, 1H), 7.23–7.25 (m, 2H), and 7.71–7.73 (m, 1H); ¹³C-NMR (100 MHz, CDCl₃) δ 14.3, 22.5, 51.4, 52.1, 53.1, 62.1, 94.7, 109.7, 116.2, 125.6, 126.8, 127.2, 127.6, 128.9, 130.0, 137.4, 140.0, 157.2, 161.3, 162.0, and 163.4; HRMS Calcd. for [C₂₁H₂₁NO₉ + H⁺]: 432.1294. Found 432.1292.

4.1.9 Rh(II)-Catalyzed Dipolar Cycloadducts 24a and 25a—To a solution of 0.1 g (0.32 mmol) of α-diazo hydroindolinone **20a** in 10 mL of benzene was added 0.1 mL (0.96 mmol) of methyl propiolate and a catalytic amount of Rh₂(OAc)₄. The mixture was heated at reflux for 1 h, concentrated under reduced pressure and subjected to flash silica gel

chromatography to give 0.1 g (80% as a 1:1 mixture of regioisomers) of cycloadducts **24a** and **25a** as a clear oil which could not be separated by silica gel column chromatography.

4.1.10 Rh(II)-Catalyzed Dipolar Cycloadduct 26—To a solution of 0.07 g (0.2 mmol) of α -diazo hydroindolinone **20a** in 15 mL of benzene was added 0.11 g (0.66 mmol) of *N*-phenylmaleimide and a catalytic amount of Rh₂(OAc)₄. The mixture was heated at reflux for 2 h, concentrated under reduced pressure and subjected to flash silica gel chromatography to give a mixture of diastereoisomeric cycloadducts **26a** (0.06 g, 57%) and **26b** (0.04 g, 38%); isomer **26a**: IR (neat) 2986, 2940, 2835, 1756, 1722, 1607, 1484, and 1378 cm⁻¹; ¹H-NMR (400 MHz, CDCl₃) δ 1.41 (t, 3H, *J* = 7.1 Hz), 2.11 (s, 3H), 3.20 (s, 3H), 4.18 (d, 1H, *J* = 8.6 Hz), 4.28 (d, 1H, *J* = 8.6 Hz), 4.47 (q, 2H, *J* = 7.3 Hz), 7.10 (d, 2H, *J* = 7.0 Hz), and 7.24–7.49 (m, 7H); ¹³C-NMR (100 MHz, CDCl₃) δ 14.3, 23.4, 48.7, 52.3, 54.0, 63.6, 79.6, 91.2, 105.8, 114.0, 125.7, 126.2, 126.7, 126.8, 129.4, 129.5, 131.3, 133.2, 136.3, 163.4, 163.5, 169.3, and 169.7.

Isomer **26b**: IR (neat) 2985, 2941, 1762, 1722, 1598, 1484, and 1214 cm⁻¹; ¹H-NMR (400 MHz, CDCl₃) δ 1.41 (t, 3H, *J* = 7.1 Hz), 2.06 (s, 3H), 3.45 (s, 3H), 3.90 (d, 1H, *J* = 6.7 Hz), 4.03 (d, 1H, *J* = 6.7 Hz), 4.45 (q, 2H, *J* = 7.0 Hz), and 7.25–7.52 (m, 9H); ¹³C-NMR (100 MHz, CDCl₃) δ 14.3, 27.4, 47.9, 51.9, 55.5, 63.2, 82.1, 88.0, 106.5, 114.6, 124.5, 125.8, 126.7, 129.5, 129.6, 130.2, 131.5, 134.2, 137.7, 161.4, 169.0, 170.8, and 171.0.

4.1.11 Rh(II)-Catalyzed Dipolar Cycloadduct 28—To a solution of 0.1 g (0.3 mmol) of α -diazo hydroindolinone **20a** in 15 mL of benzene was added 0.1 mL (0.96 mmol) of benzaldehyde and a catalytic amount of Rh₂(OAc)₄. The mixture was heated at reflux for 1 h, concentrated under reduced pressure and subjected to flash silica gel chromatography to give 0.12 g (98%) of cycloadduct **28** as a clear oil; IR (neat) 2985, 2677, 2565, 2142, 1770, 1726, 1689, and 1454 cm⁻¹; ¹H-NMR (400 MHz, CDCl₃) δ 1.32 (t, 3H, *J* = 7.0 Hz), 1.63 (s, 3H), 3.09 (s, 3H), 4.26–4.32 (m, 2H), 7.37 (d, 1H, *J* = 7.6 Hz), 7.47–7.51 (m, 3H), 7.61–7.67 (m, 2H), and 8.13 (d, 4H, *J* = 7.3 Hz); ¹³C-NMR (100 MHz, CDCl₃) δ 14.2, 24.7, 53.4, 62.0, 79.9, 114.3, 123.8, 125.3, 127.9, 128.4, 129.3, 129.9, 130.1, 133.7, 139.1, 160.0, 160.1, 172.1, and 175.7; HRMS Calcd. for [C₂₂H₂₁NO₆ + H⁺]: 396.1447. Found 396.1445.

4.1.12 3-(*tert*-Butyl-dimethyl-silanyloxy)-3-methyl-1,3-dihydro-indol-2-one—To a solution of 0.2 g (1.2 mmol) of 3-hydroxy-3-methyl-1,3-dihydro-indol-2-one³⁸ in 10 mL of CH₂Cl₂ at 0 °C was added 1.0 mL (7.0 mmol) of triethylamine and 0.7 mL (3.0 mmol) of *tert*-butyldimethylsilyl trifluoromethanesulfonate (TBSOTf). After stirring for 15 min at 0 °C, the reaction mixture was allowed to warm to rt and was stirred for an additional 2 h. The mixture was extracted with CH₂Cl₂ and the combined organic layers were washed with brine, dried over Na₂SO₄, and concentrated under reduced pressure. The crude residue was then taken up in 2 mL of THF, followed by 2 mL of acetic acid and 2 mL of water. After heating at reflux for 3 h, the mixture was warmed to rt and was extracted with EtOAc. The combined organic layers were washed with water, brine, and dried over Na₂SO₄. After removal of the solvent under reduced pressure, the residue was subjected to flash silica gel chromatography to give 0.25 g (74%) of the titled compound as an off-white solid; mp 144–146 °C; IR (neat) 3168, 2927, 2856, 1733, 1691, 1622, and 1472 cm⁻¹; ¹H-NMR (400 MHz, CDCl₃) δ -0.21 (s, 3H), -0.03 (s, 3H), 0.88 (s, 9H), 1.57 (s, 3H), 6.88 (d, 1H, *J* = 7.9 Hz), 7.06 (dt, 1H, *J* = 7.6 and 1.0 Hz), 7.25 (dt, 1H, *J* = 7.9 and 1.0 Hz), 7.32 (d, 1H, *J* = 7.6 Hz), and 8.32 (brs, 1H); ¹³C-NMR (100 MHz, CDCl₃) δ -3.8, -3.4, 18.3, 25.9, 26.8, 75.8, 110.3, 123.0, 124.2, 129.4, 133.4, 139.6, and 180.1.

4.1.13 3-[3-(*tert*-Butyl-dimethyl-silanyloxy)-3-methyl-2-oxo-2,3-dihydroindol-1-yl]-2-diazo-3-oxo-propionic Acid Ethyl Ester (20b)—To a solution of 0.24 g (0.87 mmol) of the above compound in 10 mL of THF at 0 °C was added to 0.07 g (1.7 mmol) of

NaH (60% dispersion in mineral oil). After stirring for 30 min, 0.3 g (1.7 mmol) of ethyl 2-diazomalonyl chloride²⁹ was added and the solution was stirred for 3 h. The mixture was then quenched with water and extracted with EtOAc. The combined organic layers were washed with water, brine, and dried over Na₂SO₄. After removal of the solvent under reduced pressure, the residue was subjected to flash silica gel chromatography to give 0.31 g (86%) of **20b** as a yellow oil; IR (neat) 2956, 2929, 2857, 2140, 1773, 1732, 1667, 1609, and 1467 cm⁻¹; ¹H-NMR (400 MHz, CDCl₃) δ -0.17 (s, 3H), -0.06 (s, 3H), 0.88 (s, 9H), 1.31 (t, 3H, *J* = 7.1 Hz), 1.63 (s, 3H), 4.26–4.31 (m, 2H), 7.20 (dt, 1H, *J* = 7.6 and 1.0 Hz), 7.32–7.39 (m, 2H), and 7.62 (d, 1H, *J* = 7.9 Hz); ¹³C-NMR (100 MHz, CDCl₃) δ -3.6, -3.2, 14.5, 18.3, 25.8, 27.7, 62.1, 76.0, 114.5, 123.9, 125.2, 129.7, 132.4, 138.1, 160.3, 160.4, and 177.3.

4.1.14 Rh(II)-Catalyzed Dipolar Cycloadducts 24b and 25b—To a solution containing 0.085 g (0.2 mmol) of **20b** in 10 mL of benzene was added 0.06 mL (0.6 mmol) of methyl propiolate and a catalytic amount of Rh₂(OAc)₄. The mixture was heated at reflux for 1 h, concentrated under reduced pressure and subjected to flash silica gel chromatography to give 0.07 g (70% as a 1:3 mixture of regioisomers) of cycloadducts **24b** and **25b**. The major isomer **25b** showed the following spectral properties: IR (neat) 2955, 2932, 2897, 2858, 2271, 1730, 1598, and 1472 cm⁻¹; ¹H-NMR (400 MHz, CDCl₃) δ -0.17 (s, 3H), -0.02 (s, 3H), 0.94 (s, 9H), 1.34 (t, 3H, *J* = 7.0 Hz), 1.97 (s, 3H), 3.87 (s, 3H), 4.30–4.41 (m, 2H), 6.58 (s, 1H), 7.04 (d, 1H, *J* = 6.7 Hz), 7.19–7.26 (m, 2H), and 7.83 (d, 1H, *J* = 7.0 Hz). The minor isomer **24b** showed the following spectral properties: ¹H-NMR (400 MHz, CDCl₃) δ -0.24 (s, 3H), 0.04 (s, 3H), 0.92 (s, 9H), 1.39 (t, 3H, *J* = 7.3 Hz), 2.02 (s, 3H), 3.38 (s, 3H), 4.30–4.41 (m, 2H), 6.97 (d, 1H, *J* = 6.7 Hz), 7.19–7.26 (m, 2H), 7.47 (s, 1H), and 7.91 (d, 1H, *J* = 7.0 Hz); ¹³C-NMR (100 MHz, CDCl₃) δ -4.2, -3.9, -3.1, -2.8, 14.0, 14.2, 18.3, 25.7, 25.8, 27.6, 28.2, 51.2, 52.3, 61.2, 61.5, 73.1, 74.1, 110.0, 115.9, 120.2, 124.0, 124.5, 125.3, 125.8, 126.9, 127.0, 127.4, 128.1, 128.8, 129.9, 138.8, 140.3, 141.5, 142.6, 157.7, 158.1, 160.5, 161.8, 162.7, and 164.6.

4.1.15 3-But-3-enyl-3-hydroxy-1,3-dihydro-indol-2-one (29)—Magnesium turnings (0.37 g) were added to a flame-dried, magnetic-stirred flask which was sealed and evacuated under a nitrogen atmosphere. To this mixture was added 20 mL of THF followed by the addition of 1.0 mL (10.2 mmol) of 4-bromo-1-butene dropwise. After heating at reflux for 2 h, the reaction mixture was cooled to -78 °C and then 0.5 g (3.4 mmol) of isatin dissolved in 10 mL of THF was added. After stirring for 30 min, the solution was warmed to rt and was stirred for an additional 2 h. The mixture was then quenched with a saturated NH₄Cl solution and extracted with EtOAc. The combined organic extracts were washed with water, brine, dried over Na₂SO₄, and concentrated under reduced pressure. The residue was subjected to flash silica gel chromatography to give 0.15 g (22%) of **29** as a yellow solid; mp 71–73 °C; IR (neat) 3263, 2923, 1720, 1623, 1472, and 1215 cm⁻¹; ¹H-NMR (400 MHz, CDCl₃) δ 1.92–2.06 (m, 4H), 3.16 (s, 1H), 4.88–4.96 (m, 2H), 5.65–5.75 (m, 1H), 6.88 (d, 1H, *J* = 7.6 Hz), 7.07 (dt, 1H, *J* = 7.6 and 0.9 Hz), 7.27 (dt, 1H, *J* = 7.8 and 1.3 Hz), 7.36 (d, 1H, *J* = 7.9 Hz), and 8.27 (brs, 1H); ¹³C-NMR (100 MHz, CDCl₃) δ 27.6, 37.6, 76.9, 110.6, 115.4, 123.4, 124.6, 130.0, 130.5, 137.3, 140.6, and 180.6.

4.1.16 3-But-3-enyl-3-(*tert*-butyl-dimethylsilyloxy)-1,3-dihydro-indol-2-one (30)—To a solution of 0.15 g (0.7 mmol) of the above alcohol **29** in 10 mL of CH₂Cl₂ at 0 °C was added 0.6 mL (4.2 mmol) of triethylamine and 0.4 mL (1.8 mmol) of *tert*-butyldimethylsilyl trifluoromethanesulfonate (TBSOTf). After stirring for 15 min at 0 °C, the reaction mixture was allowed to warm to rt and was stirred for an additional 2 h. The mixture was extracted with CH₂Cl₂ and the combined organic layers were washed with brine, dried over Na₂SO₄, and concentrated under reduced pressure. The crude residue was then taken up in 1 mL of THF, followed by 1 mL of acetic acid and 1 mL of water. After heating at reflux for 3 h, the mixture was warmed to rt and was extracted with EtOAc. The combined organic

layers were washed with water, brine, and dried over Na₂SO₄. After removal of the solvent under reduced pressure, the residue was subjected to flash silica gel chromatography to give 0.2 g (89%) of **30** as a pale yellow oil; IR (neat) 3250, 2955, 2929, 2857, 1725, 1621, 1472 and 1189 cm⁻¹; ¹H-NMR (400 MHz, CDCl₃) δ -0.24 (s, 3H), -0.06 (s, 3H), 0.89 (s, 9H), 1.97–2.08 (m, 4H), 4.89–4.98 (m, 2H), 5.72–5.79 (m, 1H), 6.86 (d, 1H, *J* = 7.9 Hz), 7.06 (dt, 1H, *J* = 7.6 and 1.0 Hz), 7.24–7.31 (m, 2H), and 8.00 (brs, 1H); ¹³C-NMR (100 MHz, CDCl₃) δ -3.9, -3.4, 18.4, 25.9, 27.3, 39.4, 78.0, 110.1, 114.9, 122.9, 124.9, 129.5, 131.9, 138.0, 140.0, and 179.9.

4.1.17 3-[3-But-3-enyl-3-(*tert*-butyl-dimethylsilanyloxy)-2-oxo-2,3-dihydro-indol-1-yl]-2-diazo-3-oxo-propionic Acid Ethyl Ester (31)—To a solution of 0.36 g (1.1 mmol) of the above oxindole **30** in 10 mL of THF at 0 °C was added to 0.09 g (2.2 mmol) of NaH (60% dispersion in mineral oil). After stirring for 30 min, 0.4 g (2.2 mmol) of ethyl 2-diazomalonyl chloride²⁹ was added and the solution was stirred 3 h. The mixture was then quenched with water and extracted with EtOAc. The combined organic layers were washed with water, brine, and dried over Na₂SO₄. After removal of the solvent under reduced pressure, the residue was subjected to flash silica gel chromatography to give 0.3 g (57%) of **31** as a pale yellow oil; IR (neat) 3079, 2930, 2857, 2140, 1771, 1666, 1466 and 1350 cm⁻¹; ¹H-NMR (400 MHz, CDCl₃) δ -0.22 (s, 3H), -0.12 (s, 3H), 0.89 (s, 9H), 1.32 (t, 3H, *J* = 7.2 Hz), 1.97–2.04 (m, 2H), 2.12–2.28 (m, 2H), 4.25–4.34 (m, 2H), 4.92–5.02 (m, 2H), 5.72–5.84 (m, 1H), 7.21 (t, 3H, *J* = 7.5 Hz), 7.33–7.38 (m, 2H), and 7.63 (d, 1H, *J* = 7.6 Hz); ¹³C-NMR (100 MHz, CDCl₃) δ -3.8, -3.2, 14.5, 18.4, 25.9, 26.9, 40.2, 62.1, 78.2, 114.7, 115.1, 124.5, 125.1, 129.8, 131.2, 138.0, 138.7, 160.2, and 177.1.

4.1.18 Rh(II)-Catalyzed Dipolar Cycloadduct 32—To a solution of 0.26 g (0.6 mmol) of diazo hydroindolinone **31** in 15 mL of benzene was added a catalytic amount of Rh₂(OAc)₄ and the mixture was heated at reflux for 2 h. The reaction mixture was concentrated under reduced pressure and subjected to flash silica gel chromatography to give 0.18 g (70%) of cycloadduct **32** as a clear oil; IR (neat) 2955, 2930, 2857, 1757, 1607, 1474, and 1406 cm⁻¹; ¹H-NMR (400 MHz, CDCl₃) δ -0.32 (s, 3H), -0.08 (s, 3H), 0.84 (s, 9H), 1.36 (t, 3H, *J* = 7.1 Hz), 1.79–1.99 (m, 2H), 2.22–2.31 (m, 2H), 2.39–2.54 (m, 2H), 2.63 (dd, 1H, *J* = 14.0 and 7.0 Hz), 4.37 (q, 2H, *J* = 7.1 Hz), 7.17 (dt, 3H, *J* = 7.3 and 1.3 Hz), 7.36–7.41 (m, 2H), and 7.45 (d, 1H, *J* = 7.6 Hz); ¹³C-NMR (100 MHz, CDCl₃) δ -3.6, -3.4, 14.3, 18.4, 26.0, 29.6, 34.9, 45.8, 49.9, 62.3, 78.6, 95.9, 109.6, 114.3, 125.2, 126.8, 130.5, 134.3, 138.8, 162.8, and 164.9; HRMS Calcd. for [C₂₃H₃₁NO₅Si + H⁺]: 430.2044. Found 430.2040.

4.1.19 3-Hydroxy-3-pent-4-enyl-1,3-dihydro-indol-2-one (33)—Magnesium turnings (0.37 g) were added to a flame-dried, magnetic-stirred flask which was sealed and evacuated under a nitrogen atmosphere. To this mixture was added dropwise 20 mL of THF, followed by the addition of 1.2 mL (10.2 mmol) of 5-bromo-1-pentene. After heating at reflux for 2 h, the reaction mixture was cooled to -78 °C and 0.5 g (3.4 mmol) of isatin dissolved in 10 mL of THF was added. After stirring for 30 min, the reaction mixture was warmed to rt and was stirred for an additional 2 h. The mixture was then quenched with a saturated NH₄Cl solution and extracted with EtOAc. The combined organic extracts were washed with water, brine, dried over Na₂SO₄, and concentrated under reduced pressure. The residue was subjected to flash silica gel chromatography to give 0.32 g (43%) of **33** as a yellow solid; mp 72–74 °C; IR (neat) 3254, 2939, 1716, 1623, 1472, and 1207 cm⁻¹; ¹H-NMR (400 MHz, CDCl₃) δ 1.15–1.26 (m, 1H), 1.26–1.39 (m, 1H), 1.93–2.02 (m, 4H), 3.65 (s, 1H), 4.88–4.96 (m, 2H), 5.62–5.72 (m, 1H), 6.89 (d, 1H, *J* = 7.6 Hz), 7.07 (dt, 1H, *J* = 7.6 and 0.9 Hz), 7.24 (dt, 1H, *J* = 7.6 and 1.3 Hz), 7.35 (d, 1H, *J* = 7.3 Hz), and 8.81 (brs, 1H); ¹³C-NMR (100 MHz, CDCl₃) δ 22.5, 33.8, 38.0, 77.3, 110.7, 115.3, 123.3, 124.4, 129.8, 130.8, 138.1, 140.7, and 181.3.

4.1.20 3-(*tert*-Butyl-dimethylsilyloxy)-3-pent-4-enyl-1,3-dihydro-indol-2-one (34)

—To a solution containing 0.6 g (2.8 mmol) of the above alcohol **33** in 10 mL of CH₂Cl₂ at 0 °C was added 2.3 mL (16.6 mmol) of triethylamine and 1.6 mL (6.9 mmol) of *tert*-butyldimethylsilyl trifluoromethanesulfonate (TBSOTf). After stirring for 15 min, the reaction mixture was warmed to rt and was stirred for an additional 2 h. The mixture was then extracted with CH₂Cl₂ and the combined organic layers were washed with brine, dried over Na₂SO₄, and concentrated under reduced pressure. The crude residue was then taken up in 2 mL of THF, followed by 2 mL of acetic acid and 2 mL of water. After heating at reflux for 3 h, the reaction mixture was cooled to rt and extracted with EtOAc. The combined organic layers were washed with water, brine, and dried over Na₂SO₄. After removal of the solvent under reduced pressure, the residue was subjected to flash silica gel chromatography to give 0.85 g (92%) of **34** as a yellow oil; IR (neat) 3249, 2953, 2929, 1725, 1621, 1471, 1250 and 1121 cm⁻¹; ¹H-NMR (400 MHz, CDCl₃) δ -0.25 (s, 3H), -0.07 (s, 3H), 0.88 (s, 9H), 1.25–1.48 (m, 2H), 1.87–1.93 (m, 2H), 1.99–2.04 (m, 2H), 4.90–4.97 (m, 2H), 5.67–5.78 (m, 1H), 6.86 (d, 1H, *J* = 7.9 Hz), 7.05 (dt, 1H, *J* = 7.6 and 0.6 Hz), 7.22–7.30 (m, 2H), and 8.31 (brs, 1H); ¹³C-NMR (100 MHz, CDCl₃) δ -3.9, -3.4, 18.4, 22.2, 25.9, 33.9, 39.8, 78.3, 110.2, 115.0, 122.8, 124.9, 129.4, 132.1, 138.6, 140.1, and 180.2.

4.1.21 3-[3-(*tert*-Butyl-dimethylsilyloxy)-2-oxo-3-pent-4-enyl-2,3-dihydro-indol-1-yl]-2-diazo-3-oxo-propionic Acid Ethyl Ester (35)

—To a solution of 0.22 g (0.66 mmol) of the above oxindole **34** in 10 mL of THF at 0 °C was added 0.03 g (0.8 mmol) of NaH (60% dispersion in mineral oil). After stirring for 30 min, 0.17 g (1 mmol) of ethyl 2-diazomalonyl chloride²⁹ was added and the solution was stirred 3 h. The mixture was then quenched with water and extracted with EtOAc. The combined organic layers were washed with water, brine, and dried over Na₂SO₄. After removal of the solvent under reduced pressure, the residue was subjected to flash silica gel chromatography to give 0.19 g (60%) of **35** as a pale yellow oil; IR (neat) 2955, 2930, 2139, 1771, 1732, 1664 and 1465 cm⁻¹; ¹H-NMR (400 MHz, CDCl₃) δ -0.30 (s, 3H), -0.20 (s, 3H), 0.81 (s, 9H), 1.24 (t, 3H, *J* = 7.2 Hz), 1.35–1.53 (m, 2H), 1.82–1.86 (m, 2H), 1.94–2.00 (m, 2H), 4.19–4.25 (m, 2H), 4.84–4.93 (m, 2H), 5.62–5.71 (m, 1H), 7.12 (t, 1H, *J* = 7.6 Hz), 7.25–7.29 (m, 2H), and 7.54 (d, 1H, *J* = 7.6 Hz); ¹³C-NMR (100 MHz, CDCl₃) δ -3.8, -3.2, 14.5, 18.4, 21.7, 21.7, 25.9, 29.9, 33.8, 40.5, 60.6, 62.1, 78.4, 114.7, 115.0, 124.5, 125.0, 129.7, 131.3, 138.5, 160.2, and 177.0.

4.1.22 Rh(II)-Catalyzed Dipolar Cycloadduct 36—To a solution containing 0.19 g (0.4 mmol) of diazo hydroindolinone **35** in 15 mL of benzene was added a catalytic amount of Rh₂(OAc)₄ and the mixture was heated at reflux. After heating for 2 h, the reaction mixture was concentrated under reduced pressure and subjected to flash silica gel chromatography to give 0.13 g (69%) of cycloadduct **36** as a pale yellow oil; IR (neat) 2935, 2857, 1758, 1733, 1607, 1480, and 1252 cm⁻¹; ¹H-NMR (400 MHz, CDCl₃) δ -0.35 (s, 3H), -0.24 (s, 3H), 0.80 (s, 9H), 1.22–1.28 (m, 2H), 1.34 (t, 3H, *J* = 7.2 Hz), 1.79–1.87 (m, 2H), 1.97–2.55 (m, 3H), 2.38 (dt, 1H, *J* = 14.0 and 9.5 Hz), 2.56 (dd, 1H, *J* = 14.0 and 9.5 Hz), 4.30–4.40 (m, 2H), 7.14 (t, 1H, *J* = 7.3 Hz), and 7.34–7.41 (m, 3H); ¹³C-NMR (100 MHz, CDCl₃) δ -3.6, -3.5, 14.2, 18.5, 21.8, 25.9, 28.0, 33.6, 36.0, 43.8, 62.3, 75.3, 91.2, 102.4, 114.5, 124.8, 131.0, 135.3, 136.4, 164.0, and 165.1; HRMS Calcd. for [C₂₄H₃₃NO₅Si + H⁺]: 444.2201. Found 444.2196.

4.1.23 3-Allyloxy-3-methyl-1,3-dihydro-indol-2-one—To a solution of 0.15 g (0.9 mmol) of 3-hydroxy-3-methyl-1,3-dihydro-indol-2-one³⁸ in 10 mL of toluene was added 0.3 mL (4.6 mmol) of allyl alcohol and 0.03 g (0.1 mmol) of *p*-toluenesulfonic acid. The reaction mixture was heated at reflux for 12 h. After cooling to rt, the mixture was extracted with EtOAc. The combined organic extracts were washed with a sodium bicarbonate solution, H₂O, brine, and dried over Na₂SO₄. Concentration under reduced pressure followed by flash silica gel chromatography gave 0.12 g (65%) of the titled compound as a clear oil; IR (neat) 3251, 2981,

2928, 2862, 1725, 1621, and 1472 cm^{-1} ; $^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 1.63 (s, 3H), 3.61–3.66 (m, 1H), 3.75–3.80 (m, 1H), 5.11 (dd, 1H, $J = 10.5$ and 1.6 Hz), 5.20 (dd, 1H, $J = 17.2$ and 1.6 Hz), 5.82–5.92 (m, 1H), 6.98 (d, 1H, $J = 7.6$ Hz), 7.10 (dt, 1H, $J = 7.6$ and 1.0 Hz), 7.29 (dt, 1H, $J = 7.6$ and 1.3 Hz), 7.34 (d, 1H, $J = 7.6$ Hz), and 9.45 (brs, 1H); $^{13}\text{C-NMR}$ (100 MHz, CDCl_3) δ 24.4, 67.2, 79.8, 110.9, 117.7, 123.3, 124.3, 129.5, 130.0, 134.2, 140.8, and 179.9.

4.1.24 3-(3-Allyloxy-3-methyl-2-oxo-2,3-dihydro-indol-1-yl)-2-diazo-3-oxo-propionic Acid Ethyl Ester (37)—To a solution of 0.12 g (0.6 mmol) of the above lactam in 10 mL of THF at 0 °C was added to 0.05 g (1.2 mmol) of NaH (60% dispersion in mineral oil). After stirring for 30 min, 0.2 g (1.2 mmol) of ethyl 2-diazomalonyl chloride²⁹ was added and the solution was stirred for 3 h. The mixture was then quenched with water and extracted with EtOAc. The combined organic layers were washed with water, brine, and dried over Na_2SO_4 . After removal of the solvent under reduced pressure, the residue was subjected to flash silica gel chromatography to give 0.15 g (75%) of **37** as a yellow oil; IR (neat) 2984, 2928, 2856, 2145, 1771, 1732, 1667, 1608, and 1466 cm^{-1} ; $^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 1.31 (t, 3H, $J = 7.3$ Hz), 1.65 (s, 3H), 3.62–3.67 (m, 1H), 3.72–3.77 (m, 1H), 4.24–4.32 (m, 2H), 5.11 (d, 1H, $J = 10.5$ Hz), 5.18 (d, 1H, $J = 17.2$ Hz), 5.78–5.87 (m, 1H), 7.24 (t, 1H, $J = 7.5$ Hz), 7.36–7.40 (m, 2H), and 7.65 (d, 1H, $J = 7.8$ Hz); $^{13}\text{C-NMR}$ (100 MHz, CDCl_3) δ 14.5, 25.3, 62.2, 67.5, 79.6, 114.6, 117.9, 124.0, 125.5, 128.6, 130.2, 134.0, 139.2, 160.3, 160.4, and 176.1.

4.1.25 Rh(II)-Catalyzed Dipolar Cycloadduct 40—To a solution containing 0.15 g (0.44 mmol) of **37** in 10 mL of benzene was added a catalytic amount of $\text{Rh}_2(\text{OAc})_4$ and the mixture was heated at reflux for 2 h. The reaction mixture was then concentrated under reduced pressure and subjected to flash silica gel chromatography to give 0.1 g (72%) of cycloadduct **40** as a clear oil; IR (neat) 2980, 2870, 1754, 1608 and 1471 cm^{-1} ; $^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 1.39 (t, 3H, $J = 7.3$ Hz), 1.65 (s, 3H), 2.29 (dd, 1H, $J = 12.7$ and 6.0 Hz), 2.53 (dd, 1H, $J = 12.7$ and 8.3 Hz), 2.66–2.74 (m, 1H), 3.66 (dd, 1H, $J = 11.4$ and 8.6 Hz), 4.12 (dd, 1H, $J = 8.3$ and 7.0 Hz), 4.39–4.45 (m, 2H), 7.19 (dt, 1H, $J = 7.3$ and 1.6 Hz), 7.35–7.41 (m, 2H), and 7.46 (d, 1H, $J = 7.6$ Hz); $^{13}\text{C-NMR}$ (100 MHz, CDCl_3) δ 14.1, 18.9, 32.0, 50.7, 62.6, 70.1, 80.2, 96.7, 111.3, 114.3, 125.6, 130.3, 133.6, 137.0, 161.9, and 164.3; HRMS Calcd. for $[\text{C}_{17}\text{H}_{17}\text{NO}_5 + \text{H}^+]$: 316.1185. Found 316.1183.

4.1.26 3-But-3-enyloxy-3-methyl-1,3-dihydro-indol-2-one—To a solution containing 0.15 g (0.9 mmol) of 3-hydroxy-3-methyl-1,3-dihydro-indol-2-one³⁸ in 10 mL of toluene was added 0.4 mL (4.5 mmol) of 3-buten-1-ol and 0.03 g (0.1 mmol) of *p*-toluenesulfonic acid. The reaction mixture was heated at reflux for 12 h. After cooling to rt, the mixture was extracted with EtOAc. The combined organic extracts were washed with a sodium bicarbonate solution, water, brine, and dried over Na_2SO_4 . Concentration under reduced pressure followed by flash silica gel chromatography gave 0.13 g (67%) of the titled compound as a red oil; IR (neat) 3250, 3081, 2980, 2928, 2870, 1725, 1621, and 1472 cm^{-1} ; $^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 1.60 (s, 3H), 2.28–2.34 (m, 2H), 3.10 (dt, 1H, $J = 8.3$ and 7.0 Hz), 3.28 (dt, 1H, $J = 8.3$ and 7.0 Hz), 4.98–5.06 (m, 2H), 5.68–5.79 (m, 1H), 6.97 (d, 1H, $J = 7.6$ Hz), 7.10 (t, 1H, $J = 7.3$ Hz), 7.28–7.32 (m, 2H), and 9.22 (brs, 1H); $^{13}\text{C-NMR}$ (100 MHz, CDCl_3) δ 24.4, 34.4, 65.1, 79.8, 110.8, 116.7, 123.3, 124.3, 129.7, 129.8, 134.9, 140.7, and 179.9.

4.1.27 3-(3-But-3-enyloxy-3-methyl-2-oxo-2,3-dihydro-indol-1-yl)-2-diazo-3-oxo-propionic Acid Ethyl Ester (38)—To a solution of 0.13 g (0.6 mmol) of the above lactam in 10 mL THF at 0 °C was added to 0.05 g (1.2 mmol) of NaH (60% dispersion in mineral oil). After stirring for 30 min, a 0.2 g (1.2 mmol) sample of ethyl 2-diazomalonyl chloride²⁹ was added and the solution was stirred for 3 h. The mixture was then quenched with water and

extracted with EtOAc. The combined organic layers were washed with water, brine, and dried over Na₂SO₄. After removal of the solvent under reduced pressure, the residue was subjected to flash silica gel chromatography to give 0.18 g (85%) of **38** as an orange oil; IR (neat) 2983, 2929, 2873, 2145, 1771, 1731, 1608, and 1466 cm⁻¹; ¹H-NMR (400 MHz, CDCl₃) δ 1.31 (t, 3H, *J* = 7.3 Hz), 1.63 (s, 3H), 2.26–2.31 (m, 2H), 3.11 (dt, 1H, *J* = 8.3 and 7.0 Hz), 3.24 (dt, 1H, *J* = 8.3 and 7.0 Hz), 4.25–4.31 (m, 2H), 4.98–5.06 (m, 2H), 5.68–5.79 (m, 1H), 7.24 (t, 1H, *J* = 7.5 Hz), 7.35–7.39 (m, 2H), and 7.65 (d, 1H, *J* = 7.9 Hz); ¹³C-NMR (100 MHz, CDCl₃) δ 14.5, 25.3, 34.4, 62.2, 65.5, 79.8, 114.6, 116.8, 124.0, 125.5, 128.9, 130.1, 134.8, 139.2, 160.3, 160.4, and 176.2.

4.1.28 Rh(II)-Catalyzed Dipolar Cycloadduct 41—To a solution containing 0.12 g (0.34 mmol) of **38** in 10 mL of benzene was added a catalytic amount of Rh₂(OAc)₄ and the mixture was heated at reflux for 2 h. The reaction mixture was then concentrated under reduced pressure and subjected to flash silica gel chromatography to give 0.09 g (82%) of cycloadduct **41** as a clear oil; IR (neat) 2983, 2935, 2876, 1756, 1734, 1607, and 1477 cm⁻¹; ¹H-NMR (400 MHz, CDCl₃) δ 1.38 (t, 3H, *J* = 7.1 Hz), 1.62 (s, 3H), 1.62–1.71 (m, 1H), 1.88–1.95 (m, 1H), 2.00 (dd, 1H, *J* = 12.7 and 4.8 Hz), 2.32–2.39 (m, 1H), 2.59 (dd, 1H, *J* = 12.7 and 8.3 Hz), 3.65 (dt, 1H, *J* = 11.1 and 4.4 Hz), 3.95–4.00 (m, 1H), 4.37–4.43 (m, 2H), 7.15 (dt, 1H, *J* = 7.3 and 1.3 Hz), and 7.30–7.39 (m, 3H); ¹³C-NMR (100 MHz, CDCl₃) δ 14.2, 22.1, 27.9, 34.7, 39.4, 61.9, 62.4, 78.4, 90.7, 100.5, 113.9, 124.3, 125.2, 130.1, 134.6, 136.8, 163.6, and 164.8; HRMS Calcd. for [C₁₈H₁₉NO₅ + H⁺]: 330.1341. Found 330.1340.

4.1.29 3-Methyl-3-pent-4-enyloxy-1,3-dihydro-indol-2-one—To a solution of 0.2 g (1.2 mmol) of 3-hydroxy-3-methyl-1,3-dihydro-indol-2-one³⁸ in 10 mL of toluene was added 1.3 mL (12.0 mmol) of 4-penten-1-ol and 0.03 g (0.1 mmol) of *p*-toluenesulfonic acid. The reaction mixture was heated at reflux for 12 h. After cooling to rt, the mixture was extracted with EtOAc. The combined organic extracts were washed with a sodium bicarbonate solution, water, brine, and dried over Na₂SO₄. Concentration under reduced pressure followed by flash silica gel chromatography gave 0.16 g (57%) of the titled compound as a pale yellow oil; IR (neat) 3250, 3079, 2979, 2929, 2871, 1725, 1621, 1472, 1205 and 1132 cm⁻¹; ¹H-NMR (400 MHz, CDCl₃) δ 1.59 (s, 3H), 1.61–1.68 (m, 2H), 2.02–2.15 (m, 2H), 3.03–3.15 (m, 1H), 3.21–3.26 (m, 1H), 4.89–5.08 (m, 2H), 5.69–5.79 (m, 1H), 6.92 (d, 1H, *J* = 7.6 Hz), 7.10 (t, 1H, *J* = 7.5 Hz), 7.26–7.32 (m, 2H), and 8.46 (brs, 1H); ¹³C-NMR (100 MHz, CDCl₃) δ 24.4, 29.3, 30.3, 65.2, 79.5, 110.5, 114.9, 123.3, 124.3, 129.8, 130.0, 138.3, 140.5, and 179.3.

4.1.30 2-Diazo-3-(3-methyl-2-oxo-3-pent-4-enyloxy-2,3-dihydro-indol-1-yl)-3-oxo-propionic Acid Ethyl Ester (39)—To a solution of 0.16 g (0.7 mmol) of the above compound in 10 mL of THF at 0 °C was added 0.06 g (1.4 mmol) of NaH (60% dispersion in mineral oil). After stirring for 30 min, 0.2 g (1.4 mmol) of ethyl 2-diazomalonyl chloride²⁹ was added and the solution was stirred at rt for 3 h. The mixture was then quenched with water and extracted with EtOAc. The combined organic layers were washed with water, brine, and dried over Na₂SO₄. After removal of the solvent under reduced pressure, the residue was subjected to flash silica gel chromatography to give 0.19 g (72%) of **39** as a pale yellow oil; IR (neat) 2928, 2143, 1772, 1730, 1665, 1466 and 1350 cm⁻¹; ¹H-NMR (400 MHz, CDCl₃) δ 1.32 (t, 3H, *J* = 7.2 Hz), 1.59–1.66 (m, 2H), 1.63 (s, 3H), 2.05–2.10 (m, 2H), 3.05–3.09 (m, 1H), 3.17–3.21 (m, 1H), 4.25–4.34 (m, 2H), 4.90–5.00 (m, 2H), 5.70–5.81 (m, 1H), 7.24 (t, 1H, *J* = 7.6 Hz), 7.35–7.40 (m, 2H), and 7.66 (d, 1H, *J* = 7.9 Hz); ¹³C-NMR (100 MHz, CDCl₃) δ 14.6, 25.4, 29.3, 30.3, 62.2, 65.6, 79.7, 114.6, 115.0, 123.9, 125.5, 129.0, 130.0, 138.3, 139.2, 160.1, 160.2, and 178.2.

4.1.31 Rh(II)-Catalyzed Dipolar Cycloadduct 42—To a solution of 0.18 g (0.5 mmol) of α-diazo hydroindolinone **39** in 15 mL of benzene was added a catalytic amount of

$\text{Rh}_2(\text{OAc})_4$ and the mixture was heated at reflux for 2 h. The reaction mixture was then concentrated under reduced pressure and subjected to flash silica gel chromatography to give 0.13 g (80%) of cycloadduct **42** as a clear oil; IR (neat) 3482, 2986, 2946, 1758, 1607, 1486, and 1374 cm^{-1} ; ^1H -NMR (400 MHz, CDCl_3) δ 1.36 (t, 3H, $J = 7.1$ Hz), 1.79 (s, 3H), 1.76–1.87 (m, 5H), 2.70–2.78 (m, 2H), 3.97–3.99 (m, 2H), 4.33–4.41 (m, 2H), 7.16 (t, 1H, $J = 7.5$ Hz), 7.31–7.35 (m, 2H), and 7.42 (d, 1H, $J = 7.6$ Hz); ^{13}C -NMR (100 MHz, CDCl_3) δ 14.4, 20.4, 31.0, 33.6, 36.7, 46.2, 62.5, 66.4, 83.5, 88.4, 109.3, 114.0, 124.3, 125.4, 130.3, 136.3, 137.8, 165.2, and 168.1; HRMS Calcd. for $[\text{C}_{19}\text{H}_{21}\text{NO}_5 + \text{H}^+]$: 344.1493. Found 344.1489.

4.1.32 3-But-3-enyloxy-3-pent-4-enyl-1,3-dihydro-indol-2-one—To a solution of 0.16 g (0.7 mmol) of alcohol **33** in 10 mL of toluene was added 0.3 mL (3.5 mmol) of 3-buten-1-ol and 0.03 g (0.1 mmol) of *p*-toluenesulfonic acid. The reaction mixture was heated at reflux for 12 h. After cooling to rt, the mixture was extracted with EtOAc. The combined organic extracts were washed with a sodium bicarbonate solution, water, brine, and dried over Na_2SO_4 . Concentration under reduced pressure followed by flash silica gel chromatography gave 0.11 g (55%) of the titled compound as a yellow solid; mp 78–79 °C; IR (neat) 3250, 2925, 2869, 1722, 1621, and 1471 cm^{-1} ; ^1H -NMR (400 MHz, CDCl_3) δ 1.21–1.40 (m, 2H), 1.95–2.03 (m, 4H), 2.27–2.32 (m, 2H), 3.10 (dt, 1H, $J = 8.6$ and 7.0 Hz), 3.27 (dt, 1H, $J = 8.6$ and 7.0 Hz), 4.89–5.06 (m, 4H), 5.66–5.77 (m, 2H), 6.92 (d, 1H, $J = 8.1$ Hz), 7.10 (t, 1H, $J = 6.8$ Hz), 7.26–7.30 (m, 2H), and 8.68 (brs, 1H); ^{13}C -NMR (100 MHz, CDCl_3) δ 22.2, 33.9, 34.4, 37.6, 64.9, 82.9, 110.5, 115.2, 116.7, 123.2, 124.8, 128.5, 129.8, 135.0, 138.3, 141.2, and 179.2.

4.1.33 3-(3-But-3-enyloxy-2-oxo-3-pent-4-enyl-2,3-dihydro-indol-1-yl)-2-diazo-3-oxo-propionic Acid Ethyl Ester (43)—To a solution of 0.11 g (0.4 mmol) of the above compound in 10 mL of THF at 0 °C was added 0.03 g (0.8 mmol) of NaH (60% dispersion in mineral oil). After stirring for 30 min, 0.14 g (0.8 mmol) of ethyl 2-diazomalonyl chloride²⁹ was added and the solution was stirred for 3 h. The mixture was then quenched with water and extracted with EtOAc. The combined organic layers were washed with water, brine, and dried over Na_2SO_4 . After removal of the solvent under reduced pressure, the residue was subjected to flash silica gel chromatography to give 0.1 g (60%) of **43** as a dark yellow oil; IR (neat) 3078, 2926, 2872, 2140, 1770, 1731, 1662, and 1465 cm^{-1} ; ^1H -NMR (400 MHz, CDCl_3) δ 1.32 (t, 3H, $J = 7.1$ Hz), 1.25–1.48 (m, 2H), 1.95–2.05 (m, 4H), 2.26–2.31 (m, 2H), 3.11 (dt, 1H, $J = 8.6$ and 7.0 Hz), 3.25 (dt, 1H, $J = 8.6$ and 7.0 Hz), 4.26–4.33 (m, 2H), 4.91–5.07 (m, 4H), 5.67–5.78 (m, 2H), 7.24 (dt, 1H, $J = 7.6$ and 1.0 Hz), 7.33–7.40 (m, 2H), 7.66 (d, 1H, $J = 7.9$ Hz); ^{13}C -NMR (100 MHz, CDCl_3) δ 14.6, 21.8, 33.8, 34.4, 38.4, 62.2, 65.4, 82.7, 114.8, 115.2, 116.8, 124.5, 125.4, 127.8, 130.0, 134.9, 138.3, 139.6, 160.3, and 175.9.

4.1.34 Rh(II)-Catalyzed Dipolar Cycloadducts 46 and 47—To a solution containing 0.1 g (0.24 mmol) of **43** in 10 mL of benzene was added a catalytic amount of $\text{Rh}_2(\text{OAc})_4$ and the mixture was heated at reflux for 2 h. The reaction mixture was then concentrated under reduced pressure and subjected to flash silica gel chromatography to give a mixture of cycloadducts. The major isomer **46** (55%) was obtained as a clear oil; IR (neat) 3076, 2951, 2862, 1732, 1640, 1607, and 1480 cm^{-1} ; ^1H -NMR (400 MHz, CDCl_3) δ 1.39 (t, 3H, $J = 7.3$ Hz), 1.47–1.67 (m, 3H), 1.83–1.92 (m, 2H), 1.96–2.04 (m, 4H), 2.22–2.29 (m, 1H), 2.55 (dd, 1H, $J = 12.4$ and 8.3 Hz), 3.55 (dt, 1H, $J = 11.4$ and 2.9 Hz), 3.93–3.97 (m, 1H), 4.37–4.43 (m, 2H), 4.92 (d, 1H, $J = 10.2$ Hz), 4.97 (dd, 1H, $J = 17.2$ and 1.9 Hz), 5.72–5.78 (m, 1H), 7.15 (dt, 1H, $J = 7.6$ and 1.3 Hz), and 7.31–7.40 (m, 3H); ^{13}C -NMR (100 MHz, CDCl_3) δ 14.4, 22.1, 28.8, 34.0, 35.1, 35.8, 40.9, 62.6, 62.8, 80.6, 91.3, 100.0, 114.2, 114.9, 125.0, 125.6, 130.2, 134.4, 135.5, 138.7, 163.0, and 165.1; HRMS Calcd. for $[\text{C}_{22}\text{H}_{25}\text{NO}_5 + \text{H}^+]$: 384.1806. Found 384.1803.

The minor isomer (11%) **47** was obtained as a clear oil; IR (neat) 3075, 2939, 2868, 1756, 1732, 1606, and 1480 cm^{-1} ; ^1H -NMR (400 MHz, CDCl_3) δ 1.25–1.28 (m, 2H), 1.37 (t, 3H, $J = 7.0$ Hz), 1.82–1.92 (m, 2H), 1.95–2.05 (m, 3H), 2.16–2.21 (m, 2H), 2.42–2.47 (m, 1H), 2.54–2.59 (m, 1H), 3.01 (dt, 1H, $J = 8.6$ and 7.0 Hz), 3.55 (dt, 1H, $J = 8.6$ and 7.0 Hz), 4.35–4.43 (m, 2H), 4.92–5.00 (m, 2H), 5.63–5.73 (m, 1H), 7.20 (dt, 1H, $J = 7.3$ and 1.0 Hz), and 7.37–7.48 (m, 3H); ^{13}C -NMR (100 MHz, CDCl_3) δ 14.4, 22.0, 28.1, 31.4, 34.7, 35.6, 44.8, 62.5, 63.7, 91.4, 101.8, 114.3, 116.3, 125.0, 125.3, 131.2, 133.3, 135.4, 135.8, and 165.0.

4.1.35 3-But-3-enyl-3-but-3-enyloxy-1,3-dihydro-indol-2-one—To a solution of 0.16 g (0.8 mmol) of alcohol **29** in 10 mL of toluene was added 0.3 mL (4.0 mmol) of 3-buten-1-ol and 0.03 g (0.1 mmol) of *p*-toluenesulfonic acid. The reaction mixture was heated at reflux for 12 h. After cooling to rt, the mixture was extracted with EtOAc. The combined organic extracts were washed with a sodium bicarbonate solution, water, brine, and dried over Na_2SO_4 . Concentration under reduced pressure followed by flash silica chromatography gave 0.12 g (57%) of the titled compound as a clear oil; IR (neat) 3256, 3079, 2923, 1721, 1621, and 1471 cm^{-1} ; ^1H -NMR (400 MHz, CDCl_3) δ 1.95–2.08 (m, 4H), 2.28–2.33 (m, 2H), 3.11 (dt, 1H, $J = 8.3$ and 7.0 Hz), 3.29 (dt, 1H, $J = 8.3$ and 7.0 Hz), 4.88 (d, 1H, $J = 10.2$ Hz), 4.94 (d, 1H, $J = 17.2$ Hz), 4.99 (d, 1H, $J = 10.2$ Hz), 5.03 (dd, 1H, $J = 17.2$ and 1.6 Hz), 5.69–5.78 (m, 2H), 6.95 (d, 1H, $J = 7.8$ Hz), 7.10 (t, 1H, $J = 7.5$ Hz), 7.28–7.31 (m, 2H), and 9.10 (brs, 1H); ^{13}C -NMR (100 MHz, CDCl_3) δ 27.2, 34.4, 37.1, 65.0, 82.7, 110.7, 115.0, 116.7, 123.2, 124.8, 128.3, 129.9, 135.0, 137.6, 141.3, and 179.3.

4.1.36 3-(3-But-3-enyl-3-but-3-enyloxy-2-oxo-2,3-dihydro-indol-1-yl)-2-diazo-3-oxo-propionic Acid Ethyl Ester (44)—To a solution of 0.09 g (0.35 mmol) of the above compound in 10 mL of THF at 0 $^\circ\text{C}$ was added 0.03 g (0.7 mmol) of NaH (60% dispersion in mineral oil). After stirring for 30 min, a 0.12 g (0.7 mmol) sample of ethyl 2-diazomalonate²⁹ was added and the solution was stirred for 3 h. The mixture was then quenched with water and extracted with EtOAc. The combined organic layers were washed with water, brine, and dried over Na_2SO_4 . After removal of the solvent under reduced pressure, the residue was subjected to flash silica gel chromatography to give 0.13 g (92%) of **44** as a yellow oil; IR (neat) 3078, 2925, 2855, 2141, 1770, 1731, 1664, and 1607 cm^{-1} ; ^1H -NMR (400 MHz, CDCl_3) δ 1.24–1.28 (m, 2H), 1.32 (t, 3H, $J = 7.1$ Hz), 2.00–2.14 (m, 2H), 2.26–2.31 (m, 2H), 3.12 (dt, 1H, $J = 8.6$ and 6.7 Hz), 3.25 (dt, 1H, $J = 8.6$ and 6.7 Hz), 4.26–4.33 (m, 2H), 4.90 (d, 1H, $J = 10.2$ Hz), 4.95 (d, 1H, $J = 17.2$ Hz), 5.01 (d, 1H, $J = 10.5$ Hz), 5.05 (d, 1H, $J = 17.2$ Hz), 5.70–5.79 (m, 2H), 7.24 (dt, 1H, $J = 7.6$ and 1.0), 7.34–7.41 (m, 2H), and 7.67 (d, 1H, $J = 8.3$ Hz); ^{13}C -NMR (100 MHz, CDCl_3) δ 14.5, 27.0, 34.4, 38.0, 62.2, 65.4, 82.4, 114.8, 115.2, 116.8, 124.5, 125.4, 127.6, 130.1, 134.9, 137.6, 139.6, 160.2, and 175.8.

4.1.37 Rh(II)-Catalyzed Dipolar Cycloadduct 48—To a solution containing 0.1 g (0.25 mmol) of **44** in 10 mL of benzene was added a catalytic amount of $\text{Rh}_2(\text{OAc})_4$ and the mixture was heated at reflux for 2 h. The reaction mixture was concentrated under reduced pressure and subjected to flash silica gel chromatography to give 0.07 g (79%) of cycloadduct **48** as a clear oil; IR (neat) 3076, 2978, 2949, 2859, 1756, 1734, 1607, and 1478 cm^{-1} ; ^1H -NMR (400 MHz, CDCl_3) δ 1.38 (t, 3H, $J = 7.0$ Hz), 1.59–1.66 (m, 1H), 1.82–2.29 (m, 7H), 2.55 (dd, 1H, $J = 12.4$ and 8.3 Hz), 3.55 (dt, 1H, $J = 11.4$ and 2.9 Hz), 3.93–3.98 (m, 1H), 4.36–4.43 (m, 2H), 4.90 (d, 1H, $J = 10.2$ Hz), 4.98 (dd, 1H, $J = 17.2$ and 1.6 Hz), 5.70–5.80 (m, 1H), 7.15 (t, 1H, $J = 6.7$ Hz), and 7.32–7.40 (m, 3H); ^{13}C -NMR (100 MHz, CDCl_3) δ 14.1, 27.0, 28.6, 34.8, 35.3, 40.7, 62.3, 62.6, 80.2, 91.1, 99.7, 114.0, 114.7, 124.8, 125.4, 130.1, 134.2, 135.1, 138.0, 162.7, and 164.8.

4.1.38 3-Allyloxy-3-pent-4-enyl-1,3-dihydro-indol-2-one—To a solution of 0.32 g (1.5 mmol) of alcohol **33** in 10 mL of toluene was added 0.5 mL (7.5 mmol) of allyl alcohol and

0.03 g (0.1 mmol) of *p*-toluenesulfonic acid. The reaction mixture was heated at reflux for 12 h. After cooling to rt, the mixture was extracted with EtOAc. The combined organic extracts were washed with a sodium bicarbonate solution, water, brine, and dried over Na₂SO₄. Concentration under reduced pressure followed by flash silica gel chromatography gave 0.11 g (30%) of the titled compound as a yellow oil; IR (neat) 3253, 3079, 2926, 2863, 1723, 1621, and 1471 cm⁻¹; ¹H-NMR (400 MHz, CDCl₃) δ 1.22–1.29 (m, 1H), 1.34–1.44 (m, 1H), 1.96–2.06 (m, 4H), 3.61–3.66 (m, 1H), 3.74–3.80 (m, 1H), 4.88–4.96 (m, 2H), 5.09 (dd, 1H, *J* = 10.5 and 1.3 Hz), 5.19 (dd, 1H, *J* = 17.2 and 1.3 Hz), 5.65–5.72 (m, 1H), 5.81–5.90 (m, 1H), 6.97 (d, 1H, *J* = 7.6 Hz), 7.10 (t, 1H, *J* = 7.9 Hz), 7.27–7.32 (m, 2H), and 9.37 (brs, 1H); ¹³C-NMR (100 MHz, CDCl₃) δ 22.2, 33.9, 37.6, 66.9, 83.0, 110.8, 115.2, 117.4, 123.2, 124.8, 128.2, 130.0, 134.3, 138.2, 141.4, and 179.6.

4.1.39 3-(3-Allyloxy-2-oxo-3-pent-4-enyl-2,3-dihydro-indol-1-yl)-2-diazo-3-oxo-propionic Acid Ethyl Ester (45)—To a solution of 0.07 g (0.3 mmol) of the above compound in 10 mL of THF at 0 °C was added to 0.02 g (0.6 mmol) of NaH (60% dispersion in mineral oil). After stirring for 30 min, a 0.1 g (0.6 mmol) sample of ethyl 2-diazomalonyl chloride²⁹ was added and the solution was stirred for 3 h. The mixture was then quenched with water and extracted with EtOAc. The combined organic layers were washed with water, brine, and dried over Na₂SO₄. After removal of the solvent under reduced pressure, the residue was subjected to flash silica gel chromatography to give 0.09 g (78%) of **45** as a yellow oil; IR (neat) 3079, 2982, 2930, 2871, 2148, 1774, 1730, 1705, 1662, and 1465 cm⁻¹; ¹H-NMR (400 MHz, CDCl₃) δ 1.32 (t, 3H, *J* = 7.1 Hz), 1.30–1.38 (m, 1H), 1.40–1.50 (m, 1H), 1.97–2.05 (m, 4H), 3.63–3.67 (m, 1H), 3.73–3.77 (m, 1H), 4.26–4.36 (m, 2H), 4.90–4.98 (m, 2H), 5.11 (dd, 1H, *J* = 10.5 and 1.6 Hz), 5.19 (dd, 1H, *J* = 17.5 and 1.6 Hz), 5.66–5.88 (m, 2H), 7.24 (dt, 1H, *J* = 7.6 and 1.0 Hz), 7.36–7.41 (m, 2H), and 7.66 (d, 1H, *J* = 7.9 Hz); ¹³C-NMR (100 MHz, CDCl₃) δ 14.5, 21.9, 33.8, 38.4, 63.0, 67.3, 82.6, 114.8, 115.3, 117.6, 124.5, 125.4, 127.5, 130.2, 134.1, 138.2, 139.6, 158.8, 160.2, and 175.9.

4.1.40 Rh(II)-Catalyzed Dipolar Cycloadduct 49—To a solution containing 0.09 g (0.22 mmol) of **45** in 10 mL of benzene was added a catalytic amount of Rh₂(OAc)₄ and the mixture was heated at reflux for 2 h. The reaction mixture was concentrated under reduced pressure and subjected to flash silica gel chromatography to give 0.06 g (75%) of **49** as a clear oil; IR (neat) 3075, 2980, 2944, 2869, 1757, 1608, and 1471 cm⁻¹; ¹H-NMR (400 MHz, CDCl₃) δ 1.29–1.40 (m, 1H), 1.38 (t, 3H, *J* = 7.1 Hz), 1.61–1.69 (m, 1H), 1.95–2.06 (m, 4H), 2.25 (dd, 1H, *J* = 12.7 and 6.0 Hz), 2.48 (dd, 1H, *J* = 12.7 and 8.3 Hz), 2.63–2.71 (m, 1H), 3.65 (dd, 1H, *J* = 11.8 and 8.6 Hz), 4.12 (dd, 1H, *J* = 8.3 and 7.0 Hz), 4.38–4.43 (m, 2H), 4.93 (d, 1H, *J* = 10.2 Hz), 4.98 (dd, 1H, *J* = 17.2 and 1.9 Hz), 5.70–5.78 (m, 1H), 7.17 (dt, 1H, *J* = 7.6 and 1.6 Hz), and 7.33–7.42 (m, 3H); ¹³C-NMR (100 MHz, CDCl₃) δ 14.4, 22.8, 32.0, 33.3, 34.0, 51.0, 62.8, 70.9, 82.7, 97.0, 111.5, 114.5, 115.2, 125.7, 126.4, 130.5, 134.2, 136.4, 138.3, 162.3, and 164.5; HRMS Calcd. for [C₂₁H₂₃NO₅ + H⁺]: 370.1649. Found 370.1644.

Acknowledgements

We greatly appreciate the financial support provided by the National Institutes of Health (GM 059384) and the National Science Foundation (Grant CHE-0450779).

References

- (a) Kuehne ME, Matsko TH, Bohnert JC, Motyka L, Oliver-Smith D. J Org Chem 2002;46:1981. (b) Kuehne ME, Earley WG. Tetrahedron 1983;39:3707. (c) Kuehne ME, Brook CS, Xu F, Parsons R. Pure Appl Chem 1994;66:2095. (d) Nkiliza J, Vercauteren J. Tetrahedron Lett 1991;32:1787. (e) Rawal VH, Michoud C, Monestel RF. J Am Chem Soc 1993;46:3030.

2. (a) Overman LE, Sworin M, Burk RM. *J Org Chem* 1983;48:2685. (b) Overman LE, Sugai S. *Helv Chim Acta* 1985;68:745. (c) Overman LE, Robertson G, Robichaud AJ. *J Org Chem* 1989;54:1236.
3. (a) Gallagher T, Magnus P, Huffman JC. *J Am Chem Soc* 1983;105:4750. (b) Magnus P, Gallagher T, Brown P, Pappalardo P. *Acc Chem Res* 1984;17:35.
4. (a) Rigby JH, Qabar MH. *J Org Chem* 1993;58:4473. (b) Rigby JH, Qabar M, Ahmed G, Hughes RC. *Tetrahedron* 1993;49:10219. (c) Rigby JH, Hughes RC, Heeg MJ. *J Am Chem Soc* 1995;117:7834. (d) Rigby JH, Mateo ME. *Tetrahedron* 1996;52:10569. (e) Rigby JH, Laurent S, Cavezza A, Heeg MJ. *J Org Chem* 1998;63:5587.
5. (a) Kraus GA, Thomas PJ, Bougie D, Chen L. *J Org Chem* 1990;55:1624. (b) Sole D, Bonjoch J. *Tetrahedron Lett* 1991;32:5183. (c) Bonjoch J, Sole D, Bosch J. *J Am Chem Soc* 1993;115:2064. (d) Schultz AG, Holoboski MA, Smyth MS. *J Am Chem Soc* 1993;115:7904. (e) Schultz AG, Guzzo PR, Nowak DM. *J Org Chem* 1995;60:8044. (f) Schultz AG, Holoboski MA, Smyth MS. *J Am Chem Soc* 1996;118:6210.
6. Pearson WH, Postich MJ. *J Org Chem* 1994;59:5662.
7. (a) Takano S, Inomata K, Ogasawara K. *Chem Lett* 1992:443. (b) Uesaka N, Saitoh F, Mori M, Shibasaki M, Okamura K, Date T. *J Org Chem* 1994;59:5633. (c) Mori M, Kuroda S, Zhang C, Sato Y. *J Org Chem* 1997;62:3263. [PubMed: 11671713]
8. Tietze LF. *Chem Rev* 1996;96:115. [PubMed: 11848746]
9. Tietze LF, Beifuss U. *Angew Chem, Int Ed Engl* 1993;32:131.
10. Ho, TL. *Tandem Organic Reactions*. Wiley; New York: 1992. (b) Bunce RA. *Tetrahedron* 1995;51:13103. (c) Parsons PJ, Penkett CS, Shell AJ. *Chem Rev* 1995;95:195.
11. Ziegler, FE. Chapter 7.3. In: Paquette, LA., editor. *Comprehensive Organic Synthesis, Combining C-C π -Bonds*. 5. Pergamon Press; Oxford: 1991.
12. Waldmann, H. *Domino Reaction in Organic Synthesis Highlight II*. Waldmann, H., editor. VCH; Weinheim: 1995. p. 193-202.
13. Curran, DP. *Comprehensive Organic Synthesis*. Trost, BM.; Fleming, I., editors. 4. Pergamon; Oxford: 1991. p. 779
14. Wender PA. *Frontiers in Organic Synthesis*. *Chem Rev* 1996;96:1–600. [PubMed: 11848741]
15. (a) Padwa A, Hornbuckle SF. *Chem Rev* 1991;91:263. (b) Padwa A, Weingarten MD. *Chem Rev* 1996;96:223. [PubMed: 11848752] (c) Padwa A. *Top Curr Chem* 1997;189:121.
16. Doyle, MP.; McKervey, MA.; Ye, T. *Modern Catalytic Methods for Organic Synthesis with Diazo Compounds*. John Wiley and Sons; New York: 1998.
17. Padwa A. *Helv Chim Acta* 2005;88:1357.
18. (a) Padwa A. *Top Curr Chem* 1997;189:121. (b) Mehta G, Muthusamy S. *Tetrahedron* 2002;58:9477. (c) McMills, MC.; Wright, D. Chapter 4. In: Padwa, A.; Pearson, WH., editors. *Synthetic Application of 1,3-Dipolar Cycloaddition Chemistry Toward Heterocycles and Natural Products*. Wiley; New York: 2002. (d) Padwa A. *J Organomet Chem* 2005;690:5533.
19. (a) Padwa A, Curtis EA, Sandanayaka VP. *J Org Chem* 1997;62:1317. (b) McMorris TC, Hu Y, Yu J, Kelner MJ. *J Chem Soc Chem Commun* 1997:315. (c) Kinder FR Jr, Bair KW. *J Org Chem* 1994;59:6965.
20. (a) McMills MC, Zhuang L, Wright DL, Watt W. *Tetrahedron Lett* 1994;34:8311. (b) Dauben WG, Dinges J, Smith TC. *J Org Chem* 1993;58:7635.
21. (a) Hodgson DM, Bailey JM, Harrison T. *Tetrahedron Lett* 1996;37:4623. (b) Hodgson DM, Villalonga-Barber C. *Tetrahedron Lett* 2000;41:5597. (c) Chin P, Chen B, Cheng KF. *Org Lett* 2001;3:1721. [PubMed: 11405695] (d) Graening T, Bette V, Neudörf J, Lex J, Schmalz GG. *Org Lett* 2005;7:4317. [PubMed: 16178522] (e) Hodgson DM, Avery TD, Donohue AC. *Org Lett* 2002;4:1809. [PubMed: 12000305] (f) Hodgson DM, Le Strat F, Avery TD, Donohue AC, Brühl J. *J Org Chem* 2004;69:8796. [PubMed: 15575760] (g) Hodgson DM, Le Strat F. *Chem Commun* 2004:822.
22. For examples, see: (a) Kitagaki S, Yasugahira M, Anada M, Nakajima M, Hashimoto S. *Tetrahedron Lett* 2000;41:5931. (b) Suga H, Kakehi A, Ito S, Inoue K, Ishida H, Ibata T. *Org Lett* 2000;2:3145. [PubMed: 11009367] (c) Hodgson DM, Glen R, Redgrave AJ. *Tetrahedron Lett* 2002;43:3927. (d) Hodgson DM, Stuppel PA, Pierard FYTM, Labaunde AH, Johnston C. *Chem Eur J* 2001;7:4465. (e) Hodgson DM, Glen R, Grant GH, Redgrave AJ. *J Org Chem* 2003;68:581. [PubMed: 12530886] (f)

- Inoue K, Suga H, Inoue S, Sato H, Kakehi A. *Synthesis* 2003;1413. (g) Suga H, Inoue K, Inoue S, Kahehi A, Shiro M. *J Org Chem* 2005;70:47. [PubMed: 15624905] (h) Hodgson DM, Brückl J, Glenn R, Labande AH, Selden DA, Dosseter AG, Redgrave AJ. *Proc Natl Acad Sci USA* 2004;101:5450. [PubMed: 15037752]
23. (a) Padwa A, Marino JP Jr, Osterhout MH. *J Org Chem* 1995;60:2704. (b) Padwa A, Marino JP Jr, Osterhout MH, Price AT, Semones MA. *J Org Chem* 1994;59:5518. (c) Hertzog DL, Nadler WR, Osterhout MH, Price AT, Padwa A. *J Org Chem* 1994;59:1418. (d) Hertzog DL, Austin DJ, Nadler WR, Padwa A. *Tetrahedron Lett* 1992;33:4731. (e) Osterhout MH, Nadler WR, Padwa A. *Synthesis* 1994:123.
24. Weingarten MD, Prein M, Price AT, Snyder JP, Padwa A. *J Org Chem* 1997;62:2001. [PubMed: 11671503]
25. (a) Padwa A, Price AT. *J Org Chem* 1995;60:6258. (b) Padwa A, Price AT. *J Org Chem* 1998;63:556. [PubMed: 11672045] (c) Mejía-Oneto JM, Padwa A. *Org Lett* 2004;6:3241. [PubMed: 15355022] (d) Padwa A, Lynch SM, Mejía-Oneto JM, Zhang H. *J Org Chem* 2005;70:2206. [PubMed: 15760207]
26. Mejía-Oneto JM, Padwa A. *Org Lett* 2006;8:3275. [PubMed: 16836384]
27. (a) Atta-ur-Rahman Sultana M. *Heterocycles* 1984;22:841. (b) Saxton JE. *Nat Prod Rep* 1996;13:327. (c) Node M, Nagasawa H, Fuji K. *J Org Chem* 1990;55:517.
28. England DB, Padwa A. *Org Lett* 2007;9:3249. [PubMed: 17658832]
29. Marino JP Jr, Osterhout MH, Price A, Sheehan SM, Padwa A. *Tetrahedron Lett* 1994;35:849.
30. (a) Padwa A, Fryxell GE, Zhi L. *J Am Chem Soc* 1990;112:3100. (b) Padwa A, Chinn RL, Hornbuckle SF, Zhang ZJ. *J Org Chem* 1991;56:3271. (c) Padwa A, Chinn RL, Hornbuckle SF, Zhang ZJ. *Tetrahedron Lett* 1989;30:301.
31. (a) Muthusamy S, Babu SA, Nethaji M. *Tetrahedron* 2003;59:8117. (b) Muthusamy S, Babu SA, Gunanathan C, Ganguly B, Suresh E, Dastidar P. *J Org Chem* 2002;67:8019. [PubMed: 12423129] (c) Nair V, Sheela KC, Sethumadhavan D, Dhanya R, Rath NP. *Tetrahedron* 2002;58:4171. (d) Muthusamy S, Babu SA, Gunathan C. *Tetrahedron Lett* 2002;43:3931. (e) Muthusamy S, Babu SA, Gunathan C, Suresh E, Dastidar P, Jasra RV. *Tetrahedron* 2001;57:7009. (f) Nair V, Sheela KC, Sethumadhavan D, Bindu S, Rath NP, Eigendorf GK. *Synlett* 2001:272. (g) Muthusamy S, Babu SA, Gunanathan C. *Tetrahedron Lett* 2000;41:8839. (h) Pirrung MC, Kaliappan KP. *Org Lett* 2000;3:353. [PubMed: 10814321]
32. Padwa A, Hornbuckle SF, Fryxell GE, Zhang ZJ. *J Org Chem* 1992;57:5747.
33. Padwa A, Chinn RL, Hornbuckle SF, Zhang ZJ. *J Org Chem* 1991;56:3271.
34. Huisgen, R. 1,3-Dipolar Cycloaddition Chemistry. Padwa, A., editor. I. Wiley-Interscience; New York: 1984.
35. (a) Sustmann R. *Tetrahedron Lett* 1971:277. (b) Sustmann R, Trill H. *Angew Chem, Int Ed Engl* 1972;11:838.
36. Bracher F, Hildebrand D. *Liebigs Ann Chem* 1992:1315.
37. Hinman RL, Bauman CP. *J Org Chem* 1964;24:31
38. Ogata M, Matsumoto H, Tawara K. *Eur J Med Chem Chim Ther* 1981:373.

23; R₁ = OMe; R₂ = R₃ = CO₂Me
24a; R₁ = OMe; R₂ = CO₂Me, R₃ = H
25a; R₁ = OMe; R₂ = H, R₃ = CO₂Me
24b; R₁ = OTBS; R₂ = CO₂Me, R₃ = H
25b; R₁ = OTBS; R₂ = H, R₃ = CO₂Me

26a; exo
26b; endo

27 (4 isomers)

28

Figure 2.

Scheme 1.

Scheme 2.

Scheme 3.

Scheme 4.

Scheme 5.

Scheme 6.

Scheme 7.

Scheme 8.

Table 1Bimolecular Cycloadditions of α -Diazo Amide **13** Using Various Dipolarophiles

entry	dipolarophile	product (yield)
1		 16 (75%)
2		 17 (77%)
3		 18 (78%)
4		 19 (85%)