CASE I.—Skiagraph, posterior view, showing shadow of calculus, c, in left ureter. Inset, photograph of calculus, after removal; natural size.

CASE II.—Skiagraph, posterior view, showing shadows of three calculi—c, c, c—in right ureter, and of a pseudo-calculus (calcified gland), P, on the left side. Inset, photograph of calculi after removal; natural size.

THE

British Medical Journal.

THE JOURNAL OF THE BRITISH MEDICAL ASSOCIATION.

LONDON: SATURDAY, JANUARY 1st, 1910.

ON
URETERAL CALCULI:

A REVIEW OF THE METHODS EMPLOYED FOR THEIR REMOVAL,
WITH NOTES OF TWO SUCCESSFUL CASES OF
TRANSFIBROPERITONEAL URETERO-LITHOTOMY.

BY **SINCLAIR WHITE, M.Ch., F.R.C.S.,**
SENIOR HONORARY SURGEON TO THE ROYAL INFIRMARY, AND LECTURER
ON SURGERY IN THE UNIVERSITY OF SHEFFIELD.

[WITH SPECIAL PLATE.]

THE employment of the Roentgen rays in the diagnosis of ureteral calculi has added greatly to our knowledge of, and the interest taken in, this field of surgery.

Formerly, unless the calculus could be felt through the rectum or vagina, or was arrested at the extreme end of the ureter and actually protruded into the bladder, it was difficult or impossible to distinguish between renal and ureteral stones previous to operation. The routine procedure was to explore the kidney, and then to search the ureter by means of a long flexible probe passed downwards from the renal pelvis. The angle formed by the ureter in its course over the pelvic brim was, however, a serious stumbling-block, and calculi in the pelvic portion of the ureter were not uncommonly overlooked.

The Roentgen rays have altered all this, and we are now able to predict with more or less certainty the presence or absence of a calculus in all suspected cases. Moreover—and this is a point of much importance—when a calculus is present, we can locate the exact point in the ureter where it will be found.

In the hands of an experienced radiographer it may be taken as generally correct that a negative finding excludes a calculus. Where a positive finding has been obtained the case is a little different, because phleboliths in the uterine or other pelvic veins, calcareous patches in atheromatous arteries, and especially tuberculous glands that have undergone calcification, all cast shadows which resemble in varying degrees that of an ureteral calculus. A careful scrutiny of the picture may show that the shadow of the pseudo-calculus does not lie on the anatomical line of the ureter, and that it lacks the clear-cut outline of a genuine calculus; but, in the light of cases recorded in which the mistake has been made, it is impossible to ignore altogether this source of error.

The *x* rays have demonstrated that, contrary to the belief formerly held, calculi are found in the ureters more frequently than in the kidneys. Leonard,¹ from a careful analysis of a series of cases, estimated the ratio as being two ureteral to one renal stone.

Morris,² Bovee,³ and Schenck,⁴ have shown that the vast majority of ureteral calculi are arrested at one or another of certain definite points in the ureter, and a study of its anatomy reveals why such should be the case. The lumen of a normal ureter is greatest in its central portion, where its diameter amounts to as much as one quarter of an inch. Towards the renal end it narrows to about one-seventh of an inch, and at its bladder end to one-tenth of an inch. There is a third slight constriction where it passes over the brim of the pelvis, and at this point it makes an angle of 135°.

Morris has tabulated 44 cases in which the site of the calculus was noted. Of these, 19 were found within an inch or two of the upper end, 10 at the brim of the pelvis, and 15 in the pelvic portion of the ureter.

The combined cases collected by Bovee and Schenck, omitting a few in which the stone was found in the central portion of the ureter, amount to 125. Of these, 41 were at the upper end, 25 at the pelvic brim, and 59 in the pelvic ureter.

Primary calculi of the ureters are very rare. They form around foreign bodies, or on ulcerated surfaces, and are usually phosphatic in character. The vast majority of ureteral calculi descend from the pelvis of the kidney, and most of these belong to the uric acid group, are oval in shape, and quite smooth. A small minority are oxalates with prickly surfaces which cause much irritation. When a calculus is impacted in a ureter the latter hypertrophies and becomes dilated above the stone, which frequently acts as a ball valve, and produces intermittent hydronephrosis. The calculus increases in size slowly but steadily, and if it does not escape into the bladder soon after entering the ureter there is no hope of it doing so later.

The treatment of abnormally placed ureteral calculi calls for little comment. The symptoms will be those of renal colic, and in nearly every case operative measures will include an exploration of the kidney. By prolonging the usual oblique nephrotomy incision downwards and forwards the whole of the abdominal portion of the ureter may be easily exposed, lying immediately behind the peritoneum, to which it is loosely attached. The calculus may be manipulated upwards to the renal pelvis, or, if it is not possible to do so, the ureter can be incised retroperitoneally over or, better still, just above the stone. After extracting the calculus and searching the ureter upwards and downwards the wound in the tube is closed by two or three fine catgut sutures, which should not engage the mucous coat.

In the pelvic portion the problem is much more difficult, and not a little ingenuity has been displayed in devising methods for approaching this part of the ureter. The dominant idea in planning almost all of these methods has been to reach and incise the ureter extraperitoneally.

Among the routes suggested and actually practised the following may be mentioned:

1. The intravesical.
2. The perineal.
3. The parasacral.
4. The trans-sacral.
5. The vaginal.
6. The rectal.
7. The iliac extraperitoneal.
8. The combined intraperitoneal and extraperitoneal and
9. The transperitoneal.

The vesical route is applicable to cases in which the calculus is arrested at the outlet of the ureter. In most instances the lower end of the stone protrudes into the bladder and can be felt with a searcher or actually seen through a cystoscope. Suprapubic cystotomy in the male and dilatation of the urethra in the female afford the necessary approaches.

The peritoneum covering the pelvic floor in the male is reflected on to the posterior wall of the bladder two-thirds of an inch above the lower end of the ureter, and short of

this extent the male ureter may be laid open from the bladder without injuring the peritoneum. In the female vesico-utero-lithotomy is less safe, and Thornton has recorded a fatality from peritonitis after this operation in a woman.

The perineal route was devised and successfully practised by Fenwick.⁵ Through it the lower end of the ureter is reached after a deep dissection between the urethra and prostate and the rectum. The procedure is admittedly a difficult one, and in at least one case had to be abandoned by the operator.

The para-sacral route was designed by Sir Henry Morris, and is fully described in his classical work on *Surgical Diseases of the Kidney and Ureter*. He employed it successfully in two female cases. Rigby⁶ had recourse to it in four male patients, all of whom recovered, but in one case he failed to find a calculus. The operation commences with an incision 5 in. long, parallel with, and a finger's breadth external to, the lower part of the sacrum and the coccyx. The posterior fibres of the gluteus maximus muscle, and, deeper, the sacro-sciatic ligaments are divided. The steps of the operation have been carefully thought out, and, although it involves a tedious dissection, it is undoubtedly superior to any of the other procedures conducted through the floor of the pelvis.

Cabot⁷ is responsible for suggesting the trans-sacral route, the preliminary steps of which are identical with those of Kraske's operation for cancer of the rectum. Sick relates a remarkable case in which he attempted to deal with a large pelvic calculus in this way. A portion only of the stone was removed, while the remainder worked its way up the ureter to the pelvis of the kidney, where it was discovered after a subsequent nephrectomy.

The vaginal route has been practised by Emmet, Cabot, Freyer, and others with success. At first sight it would appear to be a comparatively easy operation, but inability to fix the stone while the ureter is being opened is a drawback. In one of Freyer's cases the calculus became displaced and could not be found. Fortunately it subsequently escaped through the vaginal wound and was caught on the dressing.

Emmet used scissors for making the incision, and closed the opening in the ureter with interrupted sutures. There was no subsequent leakage. Cabot⁸ records the case of a woman aged 39, who had suffered from attacks of renal colic for sixteen years. There was a left pyonephrosis, and a small, hard mass could be felt in the left broad ligament close to the cervix uteri. After incising the ureter and removing the stone 10 oz. of pus escaped and the renal swelling disappeared. The patient ultimately recovered, but a uretero-vaginal fistula persisted for four months.

The rectal route need only be mentioned as an example of surgical temerity. Cecci has placed on record a successful case, but in the hands of other operators it has been attended with disaster.

In recent years there has been a growing tendency to attack pelvically placed calculi from above, and here, again, there is a choice of methods. The operation may be performed retroperitoneally; the peritoneum may be opened and the stone pushed up to a point above the pelvis, from whence it is removed retroperitoneally; or the stone may be dealt with transperitoneally.

The first of these methods is known as the iliac extra-peritoneal operation. It has been practised many times, and most surgeons accept it as "the operation of choice." It is carried out as follows: The abdominal parietes down to the peritoneum are divided from a point slightly above and to the inner side of the anterior superior spine of the ilium downwards and inwards to 1 in. above the centre of Poupart's ligament, and then horizontally inwards above the internal inguinal ring to the outer margin of the rectus muscle. The edges of the wound are retracted widely and the peritoneum lining the side of the pelvis is displaced inwards until the ureter is reached. The stone is now located and, if possible, pushed up above the pelvic brim, from whence it is removed through a vertical incision in the posterior wall of the ureter.

After searching for other calculi, both upwards and downwards, the ureteral wound is closed by inserting two or more Lambert sutures. If the calculus cannot be displaced upwards, the ureter is incised over it, but in these cases, owing to the great depth at which the operator is

working, it is usually quite hopeless to attempt to close the wound in the ureter by sutures, and ample provision must be made for urinary leakage which for a time will follow.

The second method—the combined intraperitoneal and extraperitoneal operation—has been resorted to by R. B. Hall⁹ and Arbuthnot Lane,¹⁰ each of whom records a successful case. Its technique consists in opening the abdomen and introducing a part, or the whole, of one hand, with which the stone is pushed up to a point above the iliac crest, from whence it is removed extraperitoneally through a small opening in the flank.

A modification of this method is to open the abdomen through one or other semilunar line; locate the stone by intraperitoneal palpation; displace it upwards as far as possible; strip the parietal peritoneum on the outer side of the wound downwards until the stone is reached, and open the ureter extraperitoneally. Gibbon¹¹ records particulars of two cases successfully operated on in this way. No attempt was made to suture the wound in the ureter, and in both cases a temporary urinary fistula resulted. A third case is reported by Erdman.¹² The patient was 7 years old, and had had symptoms for three years. The wound healed in two weeks.

There remains for consideration the transperitoneal method, which I used in the two cases recorded in this paper. It has received severe and, I think, altogether unmerited condemnation from Morris, Freyer, and others.

Morris¹³ says, "Nothing should induce the surgeon ever to incise a ureter through the peritoneum unless he is absolutely sure of the aseptic state of the urine."

Freyer¹⁴ writes: "Considering the grave danger of urinary leakage I hold that, even when the abdomen is opened for the purpose of effecting a diagnosis, if a stone is located in the ureter the abdomen should be closed and the calculus removed at a subsequent date extraperitoneally."

These opinions, coming as they do from workers who have done so much to advance our knowledge of ureteral surgery, necessarily carry much weight; but when one attempts to ascertain the facts on which they are founded there is a singular absence of evidence. I have gone carefully through the literature on the subject to which I have had access and with the exception of a case reported by Cullingworth I have failed to find a single death following the transperitoneal operation.

As much prominence has been accorded to Cullingworth's¹⁵ fatal case it may be well to reproduce his account of it.

A female, aged 30 years, admitted to St. Mary's Hospital, Manchester, 1884. Right pyonephrosis. A vaginal examination revealed a mass of stony hardness, of the size of a walnut, to the right of the uterus, and a smaller equally hard mass to the left. The urine contained pus to the amount of one-fourth of its volume. Diagnosis—right pyonephrosis, with independent disease of both ovaries.

Exploratory laparotomy through right semilunar line; right ureter enormously dilated and very tense. On following it down to the pelvis a large stone was found. Ureter incised in its long axis above stone; escape of a large quantity of highly offensive pus. Stone removed; weighed 270 grains; wound in ureter closed with difficulty by five interrupted sutures of fine carbolized silk; opposite ureter not examined because of patient's critical condition; glass tube inserted. Operation lasted an hour and a half. Death from uraemia eighty hours afterwards.

Autopsy.—Right kidney—numerous small abscesses; two calculi. Left kidney—pelvis and calyces formed a large irregular abscess cavity; left ureter greatly dilated; half an inch from bladder large stone.

It is hardly necessary to say that almost any operation performed on a patient in so desperate a condition would end fatally, and that it is illogical to quote this case as evidence of the danger attending transperitoneal uretero-lithotomy.

Let us, in the absence of facts, examine for a little while the reasons for assuming that the transperitoneal operation is a dangerous one. They resolve themselves into (a) fear of infecting the peritoneum during the operation, and (b) danger of urinary leakage subsequently.

The urine in most cases of impacted ureteral calculi is sterile, and when it is not so the virulence of the organisms it contains is no greater than that of those found in a suppurating appendix, or in a pyosalpinx, lesions which when dealt with by modern methods have scarcely a mortality. Again, in Wertheim's operation for advanced uterine cancer, the ureters are divided and implanted into the bladder, when necessary, with the happiest results.

As I hope to show presently, with proper technique, a wound in the ureter—even in the pelvic ureter—is almost as easily closed by suturing as a wound in the large intestine, while it is infinitely less likely to reopen from subsequent tension on the sutures. Surgical traditions have at all times stood in the way of progress, and nowhere could a better instance of this be found than here, where the *noli me tangere* attitude of a past generation of surgeons towards interfering with the peritoneum still holds sway, and fortified by the dicta of high authorities constitutes the last citadel of a discredited theory.

That it should be so may be explained, partially at least, by the experience of operators who have confined their work to extraperitoneal methods, whether conducted from above or below. In all of these operations the surgeon works at the bottom of a deep and narrow wound, where space and light are at a minimum and the accurate placing of sutures wellnigh an impossibility. Indeed, many operators, impressed by the hopelessness of the task, make no attempt to suture the ureter.

The conditions are, however, vastly different when the ureter is explored intraperitoneally. By placing the patient in the exaggerated Trendelenburg posture and employing the median suprapubic route the intestines fall away to the upper abdomen, where they are securely walled off by one or more towels. If retractors are now used to separate widely the edges of the abdominal wound, a clear and uninterrupted view is obtained of the line of the right ureter throughout its lower two-thirds, while the line of the left ureter can be almost equally well observed if the meso-sigmoid is displaced inwards below and outwards above the pelvic brim. In thin subjects the ureters can be distinctly seen where they cross the brim of the pelvis, and by running a finger along their course, so as to displace the subperitoneal fat, it is possible to follow them with the eye for some distance both upwards and downwards.

But whether they are visible or not, they can be easily palpated with the fingers, and there should be no question of missing a calculus in the pelvic ureter. The stone can usually be displaced up to the pelvic brim, which is the most convenient place from which to extract it. It is surprising to find how accessible the ureter is at this point. In one of my patients, a thin female, it was possible to lift the ureter covered with peritoneum quite to the level of the skin wound. The portion of ureter about to be opened is surrounded with gauze, and the peritoneum divided over it. The stone is next grasped between the fingers and thumb of the left hand, and, while so held, the ureter is clearly divided over it in a vertical direction. Should there be any leakage, this is at once mopped up. The wound in the ureter is closed by two or more fine catgut sutures passed after the manner of Lembert. These, while just missing the mucous membrane, secure firm and accurate apposition of the cut edges, which are slightly inverted. The small opening in the peritoneum is sutured in the usual way, and, after removal of the gauze and towels, the abdomen is closed without drainage.

Owing to the inverted posture of the patient, and to the fact that intraperitoneal manipulations temporarily inhibit the renal function, there is usually no escape of urine when the ureter is incised; but if there should be any reason for dreading this, the ureter above and below the site of operation may be temporarily occluded by ligatures. This precaution was actually adopted by Knaggs and Sick in cases which will be alluded to presently.

In cases of pyonephrosis and hydronephrosis, where the ureter is distended and tense, it should be aspirated and temporary ligatures applied before opening it. Finally, if drainage of the operation area is thought to be desirable it can be readily and efficiently provided through a small opening in the flank, the drainage tube ending just short of the place where the ureter has been incised, and of course extraperitoneally. Where drainage is contemplated the stone should be pushed up as high as the level of the iliac crest before it is removed. In order to render the line of suture in the ureter still more secure it has been proposed to surround the tube completely with peritoneum, either by infolding the adjacent portions or by detaching a strip and wrapping it around the ureter. This addition to the operation is, I think, wholly unnecessary, and by unduly interfering with the blood supply of the ureter may be actually harmful.

The advantages claimed for the transperitoneal operation are:

1. That it, and it alone, enables us to clear up doubtful diagnoses. Thus, in the second of my cases the patient complained chiefly of pain on the left side, and although the x-rays left no doubt as to the presence of calculi in the right ureter they also revealed a fairly well developed shadow on the left side, and as a matter of fact the diagnosis arrived at previous to exploration was bilateral ureteral calculi. Intraperitoneal palpation showed the left ureter to be free from a calculus, while the origin of the shadow of the pseudo-calculus was traced to a calcified mesenteric gland. Had the extraperitoneal route been adopted there would have been a second and wholly unnecessary operation performed in this case.

2. It is a less difficult operation than any of the extraperitoneal procedures. It ensures plenty of room and an abundance of light, and the operator is able to carry out the steps it entails with little or no disturbance of the surrounding parts, and with an amount of precision quite unattainable by any other method.

Probably in consequence of its denouncement by Morris and others, very few surgeons appear to have practised it.

F. J. Steward¹⁶ relates a successful case. The patient suffered from haematuria and painful micturition, which were thought to be due to a vesical calculus. A suprapubic cystotomy was performed. Nothing was found in the bladder, but through its walls a stone could be felt in the lower part of the right ureter. Eight days later the abdomen was opened through the lower part of the right linea semilunaris, and the stone manipulated up the ureter to a point above the iliac vessels, from whence it was removed. The wound in the ureter was closed with fine silk sutures, passed after the manner of Lembert, and a second row united the edges of the peritoneum over it. A gauze drain was inserted. There was no leakage, and the patient made a satisfactory recovery.

Lane is credited by Freyer and several other writers with a successful case, but a study of the original report shows it to be an example of the combined intraperitoneal and extraperitoneal operation.

Lawford Knaggs¹⁷ reports a case operated on after eleven years of suffering. Median laparotomy revealed a stone in the pelvic portion of the left ureter. The ureter was dilated and the left kidney hydronephrotic. The stone was pushed up to the loin, and the peritoneum over it divided. Strips of gauze were passed under the ureter above and below the stone. These served the double purpose of steadying the stone while the ureter was being incised and of preventing urinary leakage. The wound in the ureter was closed with fine silk sutures. A drainage tube was inserted through an opening in the left loin. Recovery took place without incident.

Another case is reported by Sick.¹⁸ The stone was arrested 2 in. from the bladder. A temporary ligature was applied to the ureter before opening it. The wound in the ureter was stitched up and covered with peritoneum. In addition to a peritoneal drain, a catheter was passed through the bladder into the lower end of the ureter and left in position. The patient recovered, and there was immediate resumption of the function of the corresponding kidney.

My own cases are:

CASE I.

A female, aged 18 years, who was an inmate of the Sheffield Royal Infirmary in January, 1909. She gave a history of numerous attacks of severe pain in the left loin extending downwards to the bladder. There had been haematuria from time to time, but when it was absent the urine was normal. A large fluctuating swelling could be felt in the left renal region, and exploration showed this to be a hydronephrosis. The left ureter was dilated, but the ureteral sound could not be passed beyond the brim of the pelvis. A skiagram of the pelvis taken a few days afterwards showed a large oval stone in the left ureter. The patient was operated on in the manner already described. There was no leakage of urine during the operation, which was completed in a quarter of an hour. I did not employ any form of drainage. The wound healed aseptically, and she has had no more pain.

CASE II.

A male, aged 27 years, admitted to the Sheffield Royal Infirmary, September, 1909.

History.—Since the age of 13 years he has had frequent attacks of pain in the left side, accompanied by irritability of the bladder, and occasionally by haematuria. During the last few months the attacks had recurred almost daily. His urine was acid, and contained an appreciable amount of blood. Skiagrams of the abdomen and pelvis revealed the shadows of three calculi, close together, in the pelvic portion of the right ureter, and a less clearly defined shadow of a calculus in the left ureter. On being closely questioned he now admitted that occasionally he had felt slight discomfort in the right side, but was emphatic that the severe pain had always been in the left side. A diagnosis of bilateral ureteral calculi was arrived at, and the abdomen was explored. The three calculi in the right ureter were without difficulty pushed up to a point above the pelvic brim and removed. After closing the wound in the ureter and suturing the peritoneum over it the left ureter was examined, but no stone could be found. The cause of the shadow on the left side was eventually traced to a calcified mesenteric gland situated in the mesentery. The abdomen was closed without drainage, and aseptic healing resulted. He has had no return of his pain, and has gone back to his work.

Morris has called attention to the remarkable fact that occasionally the pain induced by a renal calculus may be referred to the opposite side of the body, and there can be no doubt that this case is an instance of the phenomenon as applied to the ureters.

I am much indebted to Dr. Rupert Hallam, Medical Officer in charge of the Electrical Department at the Royal Infirmary, for the excellent skiagrams accompanying this paper, and to Dr. A. E. Barnes, Medical Registrar to the Infirmary, for his kindness in assisting me to compile the table of references.

REFERENCES.

- ¹Leonard: *Lancet*, 1905, vol. i, p. 1632. ²Morris: *Ibid.*, 1899, December 16th. ³Bovee: *Washington Medical Annals*, 1905, iv, p. 233. ⁴Schenck: *Journal of American Medical Association*, May 11th, 1901. ⁵Fenwick: *Edinburgh Medical Journal*, 1898, p. 281. ⁶Bigby: *Annals of Surgery*, 1907, vol. ii, p. 793. ⁷Cabot: *American Journal of Medical Sciences*, 1892, p. 281. ⁸Cabot: *American Textbook of Genito-Urinary Diseases*, vol. i, p. 497. ⁹Hall: *Medical Record*, 1890, October 18th, p. 430. ¹⁰Lane: *Lancet*, 1890, vol. ii, p. 967. ¹¹Gibbon: *Annals of Surgery*, 1906, vol. i, p. 743. ¹²Erdman: *Ibid.*, 1909, vol. ii, p. 797. ¹³Morris: *Surgical Diseases of the Kidney and Ureter*, 1901, vol. ii. ¹⁴Freyer: *Surgical Diseases of the Urinary Organs*. ¹⁵Cullingworth: *Pathological Transactions*, 1885, vol. xxxvi, p. 278. ¹⁶Steward: *Lancet*, 1901, vol. i, p. 1142. ¹⁷Knaggs: *Ibid.*, 1908, vol. i, p. 105. ¹⁸Sick: *Deut. med. Woch.*, 1906, p. 126.

An Address

ON THE

RATIONAL PUERPERIUM.

DELIVERED BEFORE THE EDINBURGH OBSTETRICAL SOCIETY.

BY J. W. BALLANTYNE, M.D., F.R.C.P.E.,

LECTURER ON MIDWIFERY AND GYNAECOLOGY, SURGEONS' HALL, EDINBURGH, ETC.

DR. HAULTAIN'S "plea for a rational puerperium," read before the Edinburgh Obstetrical Society last July,¹ was concerned wholly with one event in that period—namely, the resumption of the erect posture by the lying-in patient. The claim was then made that recumbency after the third day and in an ordinary case was unnecessary, and that it was rational management to allow the puerperal woman to sit up in bed on the second day and to rise on the third day. One hundred hospital patients were treated in this fashion, and it was stated that the morbidity was 3 per cent. Seven Fellows took part in the discussion which followed the reading of the paper, but, with one exception, the opinions expressed were critical. It seemed worth while, therefore, to raise the question afresh at the present time, both in view of the importance of the issues involved and because it might be anticipated that further evidence for or against early rising in the puerperium would be forthcoming.

Let me, in the first place, clearly state the question under discussion. The term "rational" has various meanings, and it may be well, in the first place, to decide upon its significance as applied adjectivally to the puerperium. The dictionary states that it may mean "endowed

with reason," "exercising or able to exercise one's reason in a proper manner," or "having sound judgement"; but obviously these definitions refer to a person rather than to a thing or a state such as the puerperium. Another meaning is "based on or derived from reason or reasoning"; but this, although nearer the sense, does not quite reach it. Perhaps the best definition is "agreeable to reason, reasonable, sensible," or, looking at it from the negative side, "not foolish, absurd, or extravagant." I am not forgetful of the fact that "rational" has had a special signification in the history of medicine, and that it has been used, as opposed to "empirical," for those physicians who deduce their treatment of cases from general principles; but I hardly think that this distinctive shade of meaning is contained in the phrase "rational puerperium," although it may not be altogether inappropriate to it. "Sensible, reasonable, not foolish, absurd, or extravagant," are doubtless the senses which underlie the word in its present application.

I take it, then, that Dr. Haultain was thus employing "rational" in his communication of July last, and that he was making a plea for early rising in the puerperium as being a reasonable and sensible procedure, and one which was neither foolish nor extravagant, although clearly opposed to custom and fashion, and, he added, to tradition and religion.

Of course, the *lying-in* period—if we may still use that term—presents other problems than the time of rising from bed. Some of these, such as the employment of vaginal douches, the management of the breasts, and the regulation of the bowels, may be regarded as solved; others, such as the nature of the diet, the giving of ergot, and the wearing of a binder, can scarcely be considered as quite settled. But, in the meantime, these matters are to be left out of account, and our attention is to be focussed on the date of the resumption of the erect posture by the puerperal patient. It may, however, be doubted whether it is possible to neglect the consideration of these other matters; whether, indeed, these details are not all bound up together with the one under consideration, and constitute, as a whole, the well-being or the undoing of the puerperal period. Still, so far as it can be done, let us deal solely with the question of the date of rising.

For some ten years or more the question of early rising has been exercising the obstetric mind on the Continent; but, rightly or wrongly, British practitioners and teachers have scarcely considered the matter, and might not now be debating the subject had not Dr. Haultain, by his experiment in the Edinburgh Royal Maternity Hospital last spring, forced it upon their notice. But the question, as a matter of fact, presented itself not ten but thirty-five years ago, when Dr. William Goodell of Philadelphia² described the method of puerperium followed at the Preston Retreat. I quote from his paper: "On the morning following the day of her labour the woman slips into a chair whilst her bed is making. This is repeated once or twice a day until the fourth or fifth day, when she, if so disposed, gets up and dresses herself. No patient quits her bed against her will, yet the force of example is so great that very few care to stay in bed when they see their companions up and about." This is a very remarkable statement to have been made thirty-five years ago, for it is almost word for word the description which we have been getting from Germany during the past five or six years. But, further, Dr. Goodell adduced the following reasons for early rising: first, labour being a physiological process, it should not be made to wear the livery of disease; second, the upright position excites the uterus to contract, and lessens the amount and duration of the lochia; third, uterine diseases are hardly known among the nations whose women early leave their beds; and, fourth, experience has shown him (Dr. Goodell) that convalescence is rendered far more prompt and sure. These, let it be noted, are, with perhaps the exception of the third one, the very reasons now being brought forward in support of the early resumption of the erect posture in the puerperium.

From Dr. Garrigues's paper,³ in which he criticizes the practice of Dr. Goodell, it may be learnt that the latter had some followers in America; but Dr. Garrigues was unable to find those who supported his views in Europe. "In the three chief countries of Europe," wrote Dr. Garrigues, "all the latest publications that have come