rib to be removed further experience may guide us; but it would seem reasonable to make the thoracic window of good size, and at least to remove any bone upon which the heart can have a direct impact.

Much stress is laid by several German writers upon the importance of removing all periosteum and perichondrium to prevent bone regeneration, while others emphasize the danger of so doing, and even claim that the posterior periosteum cannot be removed without injury to the pleura, which might, under the conditions, prove a most grave complication (König). Of actual facts to support either contention there are few, if any; but the dread of bone regeneration appears to be at least greatly exaggerated and based on the results of thoracoplasty for empyema, in which the conditions are very different. It would seem to the writer probable that, even if regeneration did tend to occur over the bulging heart, the latter would readily provide space for its action in the thin layers of bone which would first be formed, and that, as the cartilages would in no case regenerate, there would always be an ample gap in the chest wall; while the ends of the ribs, having no attachments, would accommodate themselves to the requirements of the heart. In the case of Umber and König, in which periosteum was not removed, the patient died two and a half years later, and at the *post-mortem* examination it was found that there had been no regeneration except for a few bony spicules. On these grounds I should in future be inclined to adopt König's suggestion—namely, to excise the superficial periosteum, leaving that on the deeper side of the ribs, unless it could be taken away with ease and rapidity; in a case such as Meyer-Westfeld's, in which thick and dense fibrous tissue is found below the chest wall, it is, however, both easy and advisable to remove the posterior periosteum (Umber).

Lastly, it is of essential importance to arrest all hæmorrhage before closing the wound, and it is probably wise to provide a drain for a few days, as the cut surfaces are extensive, and the movements of the heart cause a good deal of mechanical irritation in the flap, and will tend to aggravate the flow of serum.

REFERENCE.

¹ *Lancet*, July 14th, 1908, p. 7.

THE MECHANISM OF CARDIAC PAIN.

By OTTO MAY, M.A., M.B.CANTAB., M.R.C.P.LOND.,

MEDICAL REGISTRAR TO THE MIDDLESEX HOSPITAL, AND
PHYSICIAN TO OUT-PATIENTS TO THE EVELINA
HOSPITAL FOR CHILDREN.

AN investigation of the pain and other sensory disturbances occurring in a series of 50 cases of heart disease has elicited certain facts which seem to have a bearing on the mechanism of production of cardiac pain. The details of these cases are published elsewhere;¹ the purpose of the present article is to emphasize these facts, and to touch upon a few theoretical considerations arising therefrom.

The whole heart, and the aorta as far as the origin of the innominate artery, are developmentally "mesial" organs. Further, the auricles develop from the more caudal part of the primitive heart tube, the ventricles and aorta from the more cephalic part.

From these facts it could be confidently predicted (a) that the cardiac nerves are derived from both sides of the nervous axis; and (b) that the auricles are innervated from segments of the central nervous system caudal (or posterior) to those innervating the ventricles and aorta. The former statement is easily verified by the known facts of anatomy, the cardiac nerves being derived from both vago-spinal accessories, and both right and left sympathetics.

Little seems to be really known, however, as to the exact details of distribution—for example, whether the nerves from the right side supply the right chambers, and vice versa. G. A. Gibson² states that "the left ventricle and its attachments are connected mainly with the left coronary and left deep cardiac plexuses, which, along with the entire superficial cardiac plexus, are in relation with the left nerve centres." He appears to believe, also, that the right chambers are supplied from the right nerves, and, in confirmation, instances a case of failure of the

right heart in which "anginous" symptoms began on that side. On *a priori* grounds, such a scheme of innervation seems highly improbable, as the septation of the heart is a secondary process, having no relation whatever to the bilaterality of the organism.

Statement (b) receives neither verification nor denial from the anatomical textbooks, which contain no reference to the detailed innervation of the separate chambers in terms of the segments of the nervous axis. According to Head,³ the afferent paths from the auricles are connected with the fifth to eighth thoracic segments, those from the ventricles with the first to fifth thoracic, and those from the aortic arch with the third and fourth cervical, and first to fourth thoracic. These statements accord with (b), but they rest on theoretical grounds rather than on anatomical observation. Neither do they represent the whole afferent innervation, as they do not include those of the "bulbar autonomic system" (vago-spinal accessories). There is abundant physiological evidence that these nerves carry afferent cardiac fibres—for example, the "depressor" nerve of the rabbit, and the reflex effects of stimulating the central cut end of the cardiac branches of the vagus in mammals generally. Recently Einthoven⁴ has shown that the peripheral part of the cut vagus exhibits "negative variations" corresponding in frequency with the rate of the heart beat—strong presumptive evidence that the vagus transmits afferent cardiac impulses even during the normal heart rhythm. It is probable that the referred tenderness in the sterno-mastoid, to which Mackenzie⁵ has drawn special attention as a sign often elicitable in patients with failing compensation, is a clinical manifestation of these afferent cardiac impulses in the vago spinal accessories. This tenderness was present in 13 out of the 50 cases examined.

Pain was, or had been, present in 35 cases—that is, 70 per cent. In some it was slight, in others it formed the most striking feature in the clinical picture, being of a severity sufficient to dominate the patient's mode of life. Its distribution varied in different cases, the important point in this connexion being the absence of any definite relation between the site of the pain and the clinical diagnosis of the valvular lesion present. This statement is contrary to the usually expressed views. Thus Head³ describes cardiac pain as of two kinds, (1) local, (2) referred, according to the laws formulated by him and others for the reference of pain in visceral disease. He regards the pain of aortic disease as due to the left ventricle, and as situated over the upper part of the chest, the scapula, and the inner side of the left arm—that is, the territory of the upper thoracic nerves. Similarly, he explains the pain of mitral disease as auricular in origin, and referred to the territory of the middle four thoracic nerves. As regards the mechanism of production, he looks upon the causal factor as being the undue distension of the chamber in question during its diastolic pause. It is on these assumptions, in fact, that he has worked out the innervation of the different chambers as described above.

The discrepancy between Head's account and the findings in the present series of cases can best be exemplified by brief accounts of three cases of undoubted mitral disease in which the distribution of pain was of the type described by him as typically aortic. It is interesting to note that Mackenzie⁶ has described this distribution of pain as occurring in certain cases of mitral stenosis, especially in female patients. He regards it as a sign of exhausted power of contractility of the cardiac muscle.

CASE I.

M. R., female, aged 35. Rheumatic fever one year ago. Since then she has never felt strong, and, at fairly frequent intervals, has had very severe attacks of pain in the front of the chest and radiating down the inner side of the left arm. The pain has been so bad as to cause the patient to pace the room, nursing the arm with the other hand. She noticed, moreover, that these attacks left the arm with a feeling of soreness for some hours afterwards. She was sent up to hospital by her doctor with the tentative diagnosis of angina pectoris. (She never exhibited the immobility or "the fear of impending death" characteristic of the severe form of angina.)

Physical Signs.—Apex beat in fifth left space, just outside the nipple line; impulse fairly well localized, but not very strong. At the apex there is a double murmur, presystolic and systolic, and the second sound can be heard. At the base the second sound is rather accentuated to the left of the sternum; both sounds are weak to the right. The pulse is rather irregular and the liver cannot be felt.

CASE II.

H. D., female, aged 45. Two previous attacks of rheumatic fever. Admitted with all the signs of failing compensation—general oedema, ascites, orthopnoea, and cyanosis. She complained of sharp pain in the chest and left upper arm as far as the elbow.

Physical Signs in Heart.—H.A.B. fifth space, 5½ in. from the middle line. Impulse irregular and not very forcible. At apex both sounds heard, and, in addition, there is a loud, rough, systolic murmur conducted outwards to the axilla. At the base both sounds are heard and the second sound is accentuated at the pulmonary cartilage. There is no murmur at the base.

CASE III.

M. G., female. Aged 31. Four attacks of rheumatic fever. Complains of shortness of breath, palpitation, and pain. The latter is in various parts of the chest, but especially in the precordium, and often radiates down the left arm.

Physical Signs.—H.A.B. in fifth space, 3½ in. from the middle line. The impulse is slapping in character, and preceded by a well-marked presystolic thrill. At the apex a rough presystolic murmur is heard, followed by a short first sound. The second sound at the pulmonary cartilage is accentuated. The pulse is irregular, and the liver cannot be felt.

The occurrence of such cases is sufficient to cast doubt on the theory of the "mechanical" origin of cardiac pain. And a more detailed consideration of the conditions confirms the necessity of abandoning this theory. Thus, two possible methods of mechanical stimulation of the afferent nerve endings in the heart wall suggest themselves (a) by increased strain as a result of undue distension of the chamber; (b) by increased pressure as a result of excessive strength of contraction of the cardiac wall. As already mentioned, the former of these alternatives is apparently the one adopted by Head.³

If it were true, then the ideal condition for the production of heart pain would be the overfilling of a cardiac chamber during diastole. In other words, pain should be a prominent symptom in all cases of heart disease, with marked dilatation and insufficient emptying of the chambers. Now this condition is met with to an extreme degree in the common case of "double mitral disease" with failure of compensation; yet a patient afflicted in this manner not uncommonly exhibits all the physical signs of complete failure of compensation—cyanosis, orthopnoea, general anasarca—without at any time suffering any pain referable to the heart. In the series of 50 cases two well-marked examples of this condition occurred, both in young subjects, and both going to a fatal issue; yet the patients never manifested either pain or tenderness. Such cases would be quite inexplicable were the mechanical effects of cardiac dilatation the chief factor in the production of pain. The converse argument tells in the same direction—many of the severest cases of heart pain occur in patients in whom no dilatation can be observed either clinically or at autopsy.

The second alternative—increased pressure on the nerve endings as a result of excessive force of contraction—hardly calls for consideration. The condition of great muscular hypertrophy without the occurrence of any pain is so commonly met with, in athletes, in renal disease, etc., that this possibility could never be seriously urged as a fitting explanation.

It seems, then, that no mechanical theory is capable of offering a satisfactory explanation of the origin of the "painful" stimulus. But there is one factor which may be presumed to be present to some degree in all cases of heart disease, namely, some *chemical* change in the myocardium. The suggestion presents itself that the stimulus to the nerve endings may be a chemical one, some product of the disordered metabolism of the heart muscle. On consideration a number of facts can be adduced which fit in well with this suggestion. Thus, there is a general consensus of opinion that those heart affections are most characterized by pain in which, at autopsy, most evidence of myocardial degeneration is present. Again, the paroxysmal nature of the pain in many cases, without any corresponding change in the physical signs in the heart, is easier to explain on the assumption of a chemical stimulus than of a mechanical one.

No striking originality is claimed for this view; in fact, it is in some degree foreshadowed by Mackenzie's opinion, quoted above, that the pain often results from exhausted contractility, for there can be practically no doubt that the phenomena of muscular exhaustion must rest on a chemical basis. In the present state of biochemical knowledge it is impossible to suggest what actual com-

pound or compounds could be invoked as an adequate stimulus. But it is quite within the bounds of probability that competent chemical investigation of the myocardium of patients whose clinical history was known might throw considerable light on this question.

Further, this conception of the mechanism of cardiac pain tends to bring it into line with other painful conditions—notably, "cramp" of voluntary muscles and the "lightning pains" of tabes dorsalis. Mott and others believe that the latter are caused by gradual liberation of chemical irritants from the degenerating posterior root neurons. This view has displaced the earlier one—that the pain was due to pressure from sclerosis of the roots. It affords a ready explanation of the clinical observation that the pains often cease when ataxia develops, and that in those patients who never become ataxic pain continues to be a prominent feature throughout the course of the disease.

REFERENCES.

- ¹ O. May, *Archives of Middlesex Hospital*, October, 1909. ² G. A. Gibson, *Nervous Affections of the Heart*, 1905, p. 38. ³ H. Head, *Brain*, vol. xix (1896), p. 153 et seq. ⁴ W. Einthoven, *Quarterly Journal of Experimental Physiology*, vol. i, p. 244. ⁵ J. Mackenzie, *Lancet*, 1895, p. 16 et seq. ⁶ J. Mackenzie, *Diseases of the Heart*, 1908, p. 187.

INFECTIVE ENDOCARDITIS TREATED WITH ANTIDIPHThERIAL SERUM.

By W. ROSSELL JUDD, M.R.C.S.ENG., L.R.C.P.LOND.,
ASHTON-UNDER-LYNE.

IN the absence of bacteriological proof, some authorities state¹ that the cure of infective endocarditis cannot be accepted as such.

Time and the many exigencies of general practice make it almost impossible for the practitioner to make blood cultures, take opsonic controls, and indulge in the ultra-refinements of the bacteriological laboratory. The following notes are based purely on clinical observation, and their publication would not have been sought had not the case in question been seen by one whose authority goes without saying.

On August 14th, 1909, I was called to see a man, J. T., aged 30, and was told that he had just had "a severe shivering attack."

Previous History.

The patient, as far as he could remember, had always enjoyed good health, the only trouble being a "running ear," which sometimes discharged a little, but for long periods was quiescent; the condition of the ear he dated back to childhood. His habits had always been regular, and he had always been most abstemious. No history of rheumatism or injury could be elicited.

State on Examination.

I found the patient sweating profusely, the pulse was forcible, and the respirations hurried; he was pale, complained of headache, and had just had an attack of vomiting; the tongue was furred, dry, and tremulous; the temperature was 104° F. For some few days he had not felt quite as well as usual, but did not attach much importance to this condition. The provisional diagnosis made was some atypical form of typhoid.

Progress of the Case.

The pyrexial curve and absence of other typhoid symptoms dispelled the idea of typhoid fever, and I then regarded the case as one of septic infection, and by a process of exclusion infective endocarditis was surmised.

The patient, in spite of ordinary treatment, continued to have severe rigors, averaging three attacks in the twenty-four hours, with an intermittent pyrexia ranging between 97° and 104° to 105°, accompanied by delirium and followed by profuse sweats.

Examination of the heart gave at first indefinite results, but a systolic murmur could be heard over the aortic area on the third day; and on the fourteenth day a well-marked mitral systolic murmur could be heard carried up into the axilla.

The spleen was enlarged and tender. The urine contained albumen, but no casts. Examination of the left ear revealed a perforation of the tympanic membrane, with a bead of pus in the middle ear.

On the eighth day of the disease the condition of the patient was so grave that I deemed it necessary to call in further advice, and Professor Dixon Mann saw the patient with me. The diagnosis was confirmed, but the prognosis was hopeless.

Treatment.

At first the patient was put upon large doses of quinine and arsenic, but they had practically no influence upon the pyrexia; and, having to deal with a case of a pyaemic nature, whose blood stream was no doubt charged with infective bacteria, I decided to treat the patient as much as possible on germicidal lines.