nastin B2 had been followed in him on one occasion by a reaction of such violence that the form was deemed to be unsafe and was definitively thrown out of use. I had no reason at all for doubting the soundness of my supply; but here, it seemed, was an opportunity of testing such a suspicion as Dr. Lenz has more recently mentioned, and at my request the Chief Health Officer of New Zealand was good enough to turn over to me the remainder of his stock of nastin B2. I applied it in the following way:

H. F., male, aged 40, lepra tuberosa, duration unascertained (but probably several years), received 1 c.cm. of nastin B2 (N.Z.) on three days with two intervals of two days each. His temperature remained unaffected and no other result ensued.

Cases I and II (above) each had 1 c.cm. of nastin B2 (N.Z.) on each of two days; Case III had one such dose; in Case V the nastin B2 continuously administered on and after May 28th to the end of the year was part of this New Zealand supply.

In none of the patients did it cause any reaction, either local or general. These discrepancies are puzzling. It is possible that they have their psychological aspect.

As regards the bacilli in the cases described above, they had their usual appearances, and no changes could be detected. Professor Deycke's description⁷ of the degenerative alterations observed after treatment do not (I think) indicate anything unusual. In cases of leprosy, in which bacilli can be easily obtained from the surface of the body, it is generally found that wholly-stained specimens lie among a larger number of others which are either vacuolated, or dotted, or reduced to mere acid-fast granules. In many situations none wholly stained are found; in some, which perhaps are generally more recent deposits, all may be wholly stained. The imperfect bacilli can take a differential stain.

REFERENCES.

- ¹ 1909-10. J. Ashburton Thompson: Eighteenth and Nineteenth Annual Reports of the Board of Health on Leprosy in New South Wales.
- ² 1904-7. Deycke: V Internationale Dermatologen-Kongress, Berlin; *Deut. med. Woch.*, 1905, SS. 480, 545; *Lepra*, vii, p. 174.
- ³ 1909. Lenz: *Lepra*, ix, p. 19.
- ⁴ 1909. Ziemann: *Ibid.*, p. 23.
- ⁵ 1909. Rodriguez: F. A. Smith and Bissett, Beauchamp Williams: *Trans. of Bombay Medical Congress*. F. Raschid: *BRITISH MEDICAL JOURNAL*, ii, p. 1343.
- ⁶ 1909. Heiser: *Amer. Journ. of Med. Sci.*, cxxxviii, September.
- ⁷ 1908. Deycke: *BRITISH MEDICAL JOURNAL*, i, April 4th.

THE SCIENCE COMMITTEE

OF THE

British Medical Association.

REPORT CXX.

PERITONEAL BLOOD TRANSFUSION.

By A. E. BOYCOTT, B.Sc., M.D. OXON.,
LECTURER ON PATHOLOGY, GUY'S HOSPITAL.

BLOOD injected into the peritoneal cavity is quickly absorbed into the circulation, with the exception of a small quantity which remains entangled in the omentum. The volume of the circulating blood is within a few days adjusted to its proper level, so that the concentration of red cells is increased and a condition of polycythaemia produced. The general phenomena are fully described by Hunter,¹ where an abstract of the older literature will be found.

In previous investigations Boycott and Douglas² determined the rate at which the red cells contained in the blood of normal rabbits transfused by direct intravenous injection into other normal rabbits are destroyed. They found that the mean daily amount of haemoglobin destroyed, expressed in terms of oxygen capacity, varied from -0.19 c.cm. to 0.44 c.cm., on an average 0.24 c.cm. It was the immediate object of the present research to see whether the rate of destruction of red cells passed into the circulation via the peritoneum was greater or less than this. In a series of twenty-two experiments a mean daily rate of 0.55 c.cm. was found. From a consideration of the nature of the stimulus which excites a rabbit to destroy the red cells of another rabbit which it may find in its

circulating blood, it appeared that the relatively foreign nature of the cells was an important factor. The degree of foreignness of the cells is presumably increased by passing them through an extravascular position (peritoneum) on their way to the circulation, and it is suggested that this is the reason why cells so transfused are destroyed more rapidly than when they are put straight into the blood of the recipient. With intravenous transfusions the rate of daily destruction increases as the process advances, while with intraperitoneal transfusions the rate is much quicker during the first than during the second week.

The red cells are not destroyed by a process of intravascular haemolysis, nor has it been found possible to induce the production of antolytic or isolytic substances in the blood serum of rabbits by repeated intraperitoneal injections of whole or laked blood, nor by destroying a rabbit's red cells inside the body by such a poison as phenylhydrazine. The initial phase of destruction seems to find its morphological expression in abundant phagocytosis of red cells by the endothelial (stroma) cells of the lymphatic glands and spleen pulp. Within these cells the red corpuscles are broken up, and pass gradually to a condition of amorphous pigmented debris, some of which gives the reaction for inorganic iron. This phagocytosis is very much more abundant with intraperitoneal than with intravenous transfusions—an observation in accordance with the view that the cells are made more foreign by passage through the peritoneum.

Apart from the various other reasons which point in the same direction,³ these observations suggest that intravenous transfusion is to be preferred to intraperitoneal transfusion, as giving a prospect of more lasting benefit to the recipient. This conclusion does not, however, necessarily follow, since experiments show that considerations of whether the red cells are needed in the body can outweigh those concerning the precise degree by which they are foreign to the economy.

REFERENCES.

- ¹ *Journal of Anatomy and Physiology*, vol. xxi, 1887. ² *Journal of Pathology and Bacteriology*, vols. xiii and xiv; *BRITISH MEDICAL JOURNAL*, May 22nd, 1909. ³ Crile, *Haemorrhage and Transfusion*, 1909.

Memoranda:

MEDICAL, SURGICAL, OBSTETRICAL.

LYSSOPHOBIA OR HYDROPHOBIA?

In June, 1903, my orderly—a Mohammedan Sepoy of the Kapurthala Infantry—was bitten by a dog on the outer side of the left leg below the knee. The wounds (two) healed in eight days, and he had forgotten all about the incident until February 9th, 1905—one year and eight months later—when he felt unwell, irritable, and fretful; that evening he had some fear of water, and was evidently thinking about the bites. On the following day (February 10th), when sitting in the quarter guard, he told his comrades the story of the dog, mentioning his symptoms of the previous day and his fear of water. The native officer in charge did not think him really ill, but at 11 p.m. locked him up in an adjoining room, and at 6 a.m., when he opened the door, found that the man had developed symptoms of hydrophobia. His brothers were sent for, and they conveyed him to his house in the city. He was quite conscious, but, as he bit at the sight of water, his eyes were bandaged during the journey. When I saw him at 11 a.m. on February 11th he was sitting on his bed retching and spitting with a brass basin in front of him. His hands were kept down by a pillow placed in front of them. He was biting and snapping, with quiet intervals. His aged mother watched by his side and when he wanted to bite, a pillow was each time thrust toward him and he bit it. He occasionally raised his hand to his mouth as if to bite, but he never closed his teeth over it, nor did he leave any impression of teeth on the skin, although the pillow presented to him was torn to bits. His eyes were wild, ferrety, intensely congested, and the pupils dilated. In the intervals he spoke naturally, and asked his brother to give me a chair. He held up his leg and pointed out the scars of the bites, which had a blue congested look. With great vehemence he declared that he had felt pain in the